Research Highlights

Highlights from the latest articles in novel and rapid personal whole-genome sequencing: recent advances and the promise for translational medicine

C Ramana Bhasker¹ & Gary Hardiman^{2,3}

¹Department of Pediatrics, School of Medicine, University of California, San Diego, La Jolla CA 92093, USA

E-mail: rbhasker@ucsd.edu

²Department of Medicine, School of Medicine, University of California, San Diego, La Jolla CA 92093, USA

³BIOGEM, School of Medicine, University of California, San Diego, La Jolla CA 92093, USA
E-mail: ghardiman@ucsd.edu

Individual diploid genome sequence

Evaluation of: Levy S, Sutton G, Ng PC *et al.*: The diploid genome sequence of an individual human. *PLoS Biol.* 5(10), E254 (2007).

Over the past decade, with the development of high-throughput DNA sequencing protocols and advances in computational analytics, sequence assemblies have been generated that encompass the majority of the human genome [1–4]. Two versions of the human genome that have been studied to date are products of the Human Genome Sequencing Consortium [3] and Celera Genomics (MD, USA) [1], which are derived from clone-based and random whole-genome shotgun sequencing strategies, respectively. The Human Genome Sequencing Consortium assembly is a composite derived from haploids of numerous donors, whereas the Celera genome represents a consensus derived from five individuals. Variation has been reported primarily as differences in SNPs. Smaller-scale (100 base pairs) insertion/deletion sequences (indels) and large-scale structural variants have been largely ignored, which is unfortunate as both contribute to human biology and disease.

In an effort to provide a comprehensive analysis of variation in the context of human biology, the genome of J Craig Venter was sequenced using traditional Sanger sequencing. The complexities of assessing genotype–phenotype relationships are highlighted from a very preliminary glimpse of Venter's genome. His predisposition status for multifactorial

diseases is complicated. In his family there are documented cases of chronic disease, including hypertension and ovarian and skin cancer. A family history of cardiovascular disease exists, and his father died at 59 years of age of sudden cardiac arrest, so an examination of pertinent genotypes was undertaken. Venter is heterozygous for variants in the *KL* gene: F352V (r9536314) and C370S (rs9527025). It has previously been reported that these heterozygous alleles present a lower risk for coronary artery disease. However, he is also homozygous for the 5A/5A in rs3025058 in the promoter of the matrix metalloproteinase-3 (*MMP3*), a genotype associated with higher intra-arterial levels of stromelysin with a higher risk of acute myocardial infarction.

Genetic changes related to known disease risks for Venter were identified. Approximately 50% of the Caucasian population (including Venter) is heterozygous for the *GSTM1* gene, where a null mutation can increase susceptibility to the adverse effects of environmental toxins and carcinogens. Further testing (including somatic analysis) and comparison against larger datasets is, however, needed to conclusively determine if this variant contributes to this individual's reported medical conditions, such as skin cancer.

Surprisingly, inconsistencies were noted between deleted genotypes in the donor's DNA and the expected phenotype. One notable example is his *LCT* genotype, which should confer adult lactose tolerance, but is inconsistent with the lactose intolerance phenotype

For reprint orders, please contact:
reprints@futuremedicine.com

self-reported by the donor. An explanation for this is the modifying effect of other genes and their products in addition to environmental influences.

References

1. Levy S, Sutton G, Ng PC *et al.*: The diploid genome sequence of an individual human. *PLoS Biol* 5, E254 (2007).
2. Venter JC, Adams MD, Myers EW *et al.*: The sequence of the human genome. *Science* 291, 1304–1351 (2001).
3. Lander ES, Linton LM, Birren B *et al.*: Initial sequencing and analysis of the human genome. *Nature* 409, 860–921 (2001).
4. International Human Genome Sequencing Consortium: Finishing the euchromatic sequence of the human genome. *Nature* 431, 931–945 (2004).

DNA sequence of a diploid genome of a single individual

Evaluation of: Wheeler DA, Srinivasan M, Egholm M *et al.*: The complete genome of an individual by massively parallel DNA sequencing. *Nature* 452, 872–876 (2008).

The practice of medicine will soon be tailored to a patient's genetic make up based on emerging so-called 'next-generation technologies', thus heralding an era of personalized healthcare. This vision is realized through the development of highly efficient and relatively inexpensive whole-genome sequencing methodologies, carried out in miniscule picoliter wells that sequence DNA in a mere fraction of the time it took a few years ago.

A plethora of genes were sequenced from human and other species beginning in the 1970s, using methodologies developed independently by Frederick Sanger and Walter Gilbert. Two decades later, laboratories from around the world deciphered the sequence of the complete human genome by employing Sanger's chain termination method coupled to traditional laborious cloning steps. Sequencing the diploid human genome, comprising approximately 6 Gb, with conventional sequencing techniques is a stupendous task. More recently, application of massively parallel sequencing (MPS) developed by the Rothberg group is replacing traditional approaches. MPS was utilized to sequence 25 million bases of the fungi genome and the task was accomplished within an incredible span

of 4 h [1,2]. The MPS approach was employed recently to decipher the diploid genome of James Watson in an incredibly short span of 2 months, at a cost of US\$1 million, with 7.4-fold redundancy [3]. These next-generation technologies clearly herald the advent of personalized genomics [4].

MPS is achieved by attaching fragmented genomic DNA molecules to linkers facilitating binding to beads 28 µm in diameter, such that a single DNA molecule adheres to each bead. All the requisite ingredients for emulsion PCR are provided. The beads are encased in an oil droplet to separate them from other beads, creating a self-contained microenvironment resulting in over 10 million amplified copies of the original template. Optical scanning detects the newly added nucleotides to each bead at each step, utilizing a technique wherein the released ATP in the presence of luciferin triggers bioluminescence [4]. Each of the four nucleotides are sequentially added and washed to allow the detection of illuminated spots at each step.

A comparison of Watson's genomic DNA sequence to the earlier Human Genome Project reference sequence identified over 3.3 million SNPs in his DNA sequence, of which a large proportion (82%) are sites previously identified in the reference. Of these, approximately 11,000 SNPs alter protein sequence, but the relevance of these SNPs that are unique to Watson's genome (15%) is not

immediately understood. It is likely that humans, in general, harbor a large number of variations, and that these acquired changes may contribute to phenotypic characteristics, predisposition to disease and its treatment. These studies shed light upon whether a bias for these genetic changes is reflected in racial and/or ethnic groups. The compelling task of sequencing a large number of individual genomes so as to unravel the possible links to a disease state is well anticipated.

This study also permitted the most comprehensive comparison of the non-synonymous difference between two diploid human genomes, namely those of Watson and Venter. The Venter Genome sequenced using traditional approaches was also recently reported [5]. It contained approximately 2.8 million known SNPs, and approximately 0.74 million novel SNPs, which was in good agreement with the study by Wheeler *et al.* The two individuals shared 1.68 million of the SNPs, of which 5230 were nonsynonymous, accounting for 58% of the subject's nonsynonymous SNPs. Both men are each distinguished from the reference genome by 3766 and 3882 nonsynonymous SNPs, respectively, and therefore are different from each other by 7648 protein-coding changes.

A challenge exists for healthcare providers to assimilate this vast genomic information relating to all variations in an individual and prescribe an effective

and customized mode of treatment. The take-home message from the paper by Wheeler *et al.* is that it is difficult to uncover medically or biologically robust inferences from an individual's sequences. This is the first human genome sequenced by next-generation technologies, and thus serves as a pilot for personalized genome sequencing. The next step will be figuring out how best to correlate this information with phenotype, particularly in the context of disease predisposition and treatment.

References

1. Margulies M, Egholm M, Altman WE *et al.*: Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437(7057), 376–380 (2005).
2. Rogers YH, Venter JC: Genomics: massively parallel sequencing. *Nature* 437, 326–327 (2005).
3. Wheeler DA, Srinivasan M, Egholm M *et al.*: The complete genome of an individual by massively parallel DNA sequencing. *Nature* 452, 872–876 (2008).
4. Nyrén P, Pettersson B, Uhlen M: Solid phase DNA minisequencing by an enzymatic luminometric inorganic pyrophosphate detection assay. *Anal. Biochem.* 208, 171–175 (1993).
5. Levy S, Sutton G, Ng PC *et al.*: The diploid genome sequence of an individual human. *PLoS Biol.* 5, E254 (2007).

Paired-end mapping for detecting structural variation in the human genome

Evaluation of: Korbel JO, Urban A, Affourtit JP *et al.*: Paired-end mapping reveals extensive structural variation in the human genome. *Science* 318 (5849), 420–426 (2007).

The human genome is bereft with structural anomalies that compound the understanding of interindividual changes. Clearly, understanding genetic variation in a population is a prerequisite to the practice of personalized medicine. Geneticists have laboriously identified polymorphisms in genes that contribute to medical anomalies, and pharmacogeneticists have assiduously predicted the complications associated with the 'one-dose-fits-all' approach that may result in drastic consequences in patients who are poor metabolizers. The occurrence of nearly 3.3 million SNPs in the genome of an individual [1] raises the probability that a similar numbers of SNPs are likely to exist in the genomes of the rest of mankind. It was recently reported in *Science* that extensive structural variations that involve major deletions, duplications, insertions and inversions, some of these ranging from kilo- to mega-bases, exist in the human genome, besides complex rearrangements [2]. It is predicted that such variations are more widespread in the population than was earlier anticipated, and such large variations are more likely to influence genetic disease

and the phenotype of individuals when compared with SNPs that may often result in silent mutations.

The pair-end mapping (PEM) technique was developed [2] to detect structural variations in the genomes of a female of African ancestry and a second female of unknown but presumed European ancestry. Briefly, the technique involved shearing genomic DNA, selecting fragments of approximately 3 kb to which biotinylated adaptors were ligated, and circularizing the fragments. Following random cleavage of the circularized DNA, fragments containing the linkers were isolated and subjected to '454 sequencing'. Computational analysis of the sequence information generated from over 30 million 'paired ends' facilitated the mapping of structural variations in the two genomes. Interestingly, genome-wide estimates of structural variations detected for the two individuals of African and European ancestry were 761 and 887, respectively.

The two genomes did contain hot spots, for instance an 8 Mb region on chromosome 22 contained 13 structural variations, and an 18 Mb region on chromosome 7 contained 29 structural variations. Nearly 45% of the variations predicted were shared between the two individual genomes, suggesting that these variations could have an ancient origin. Although these variations could extend to

over 100 kb, the majority were less than 10 kb (65%) or 5 kb (30%) in size and, as expected, more number of genes were affected by the larger variations. The PEM method identified 407 variations that were not previously detected. A further observation was that approximately 20% of the variations were homozygous, and the actual numbers were different for the two individuals.

It is a sobering thought that so-called 'normal human genomes' harbored vast genomic differences ranging from insertions, deletions and inversions, to major chromosomal rearrangements. Not unexpectedly, deletions as called in one individual could represent an insertion in another. However, it is becoming increasingly evident that unless a large number of genomes (over 1000 as suggested by several groups) are sequenced, it will be difficult to ascribe variations in the genome by comparing one individual to another. It is therefore unlikely that we will know the composition of the 'normal genome' until this time. It is pertinent to ask whether the genome is identical in all cell types of an individual, and whether major structural variations and chromosomal rearrangements, of the kind reported in the two genomes sequenced by the PEM protocol, could exist in different cells and tissues of the same individual. It raises the question as to what types of variations one may observe in

different forms of cancer, and whether such variations also reflect interindividual differences for similar types of cancers.

■ ■ ■ ■ ■ ■ ■ ■

References

1. Wheeler DA, Srinivasan M, Egholm M *et al.*: The complete genome of an individual by massively parallel DNA sequencing. *Nature* 452, 872–876 (2008).

2. Korbelt JO, Urban AE, Affourtit JP *et al.*: Paired-end mapping reveals extensive structural variation in the human genome. *Science* 318, 420–426 (2007).

■ ■ ■ ■ ■ ■ ■ ■

Financial & competing interests disclosure

The authors have no relevant affiliations or financial involvement with any organization or

entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.

No writing assistance was utilized in the production of this manuscript.