A selected review of abstracts from the 11th Congress of the European Association of Neuro-Oncology (EANO)

Marc C Chamberlain*

**European Association of Neuro-Oncology Meeting
Turin, Italy, 9–12 October 2014**

SUMMARY The European Association of Neuro-Oncology (EANO) is the largest neuro-oncology meeting in Europe that meets biannually and reproducibly provides an exciting forum to present new brain cancer clinical trials and research data. The EANO 2014 meeting in Turin, Italy (9–12 October 2014) was comprised of 3 days of presentation, nearly 50 oral presentations and nearly 350 abstracts provides a contemporary overview of neuro-oncology that includes both metastatic diseases of the CNS as well as primary brain tumors. This summary attempts to highlight select abstracts presented at the meeting of EANO 2014 in a short review that provides a portrait of a large and multifaceted meeting.

Metastatic disease to the CNS

• Parenchymal brain metastases

Kwakman *et al.* [1] estimated the incidence of brain metastases (BMs) and patterns of care in The Netherlands concluding that basic epidemiologic, treatment and outcome data are poorly collected by tumor registries. Furthermore, there were major local differences in treatment policies and infrequent adherence to national guidelines.

Wen *et al.* [2] presented the newly proposed response assessment in neuro-oncology (RANO) criteria for response assessment in BM wherein a radiographic measurable (target) lesion is defined by a minimal contrast enhancing diameter of 10 mm (using a unidimensional methodology similar to RECIST) and that is visible on 2 or more axial images. Best overall response is a composite of target and nontarget response, corticosteroid use and clinical status. Radiographic response requires a confirmation scan at four or more weeks.

In a pathology study by Berghoff *et al.* [3], the density of tumor-infiltrating lymphocytes (TILs) as defined by CD3 or CD8 immunohistochemical positivity was correlated with peritumoral edema and overall survival (increase in TIL density, improved survival).

Galldiks *et al.* [4] reported that 18-fluoroethyl-L-tyrosine PET imaging helps differentiate radiation induced changes from true tumor progression not otherwise evident by anatomic MRI. This distinction is often relevant in BM treated with stereotactic radiotherapy (SRT).

Scoccianti *et al.* [5] demonstrated that double contrast brain MRI (using a T1-weighted 3D magnetization-prepared rapid acquisition gradient echo sequence [MPRAGE]) prior to SRT often alters therapy in that a third of patients had more BM than previously revealed by standard dose MRI.

A quality of life (QoL) study by Habets *et al.* [6] was reported in patients with BM and treated with SRT. There did not appear to be a detrimental effect on health related QoL or cognition (using a standard EORTC instrument) in patients treated with SRT.

KEYWORDS

- EANO 2014 abstracts
- primary and secondary (metastatic) brain tumors

*University of Washington, Department of Neurology/Division of Neuro-Oncology, Fred Hutchinson Cancer Research Center, Seattle Cancer Care Alliance, 825 Eastlake Avenue E, POB 19023, MS G4940, Seattle, WA 98109-1023, USA; Tel.: +1 206 288 8280; Fax: +1 206 288 2000; chambemc@u.washington.edu

• Leptomeningeal disease

Bionski *et al.* [7], in a small series of patients with non-small-cell lung cancer related-leptomeningeal metastasis (LM), compared cerebrospinal fluid (CSF) cytology with delineation of circulating tumor cells (CTC) by Veridex CellSearch technology and found the later method superior to conventional CSF cytology as well as providing quantification of CSF CTC.

Chamberlain *et al.* [8] reported on new RANO response criteria for LM. The proposal utilizes two new and as yet not validated instruments; a standardized scorable neurological examination and an imaging worksheet that has ten brain and spine domains that are scored. In conjunction with CSF CTC response, a composite score is obtained. Currently there are no standardized and agreed upon response criteria in LM.

Primary brain tumors

• Primary CNS lymphoma

Ahle *et al.* [9] presented results of a database search of 710 patients with primary CNS lymphoma (PCNSL) and found only 1% of patients present at diagnosis with an ischemic optic neuropathy (due to lymphomatous meningitis) as compared with 15% of patients presenting with a posterior uveitis.

Larrieu *et al.* [10] reported on a monocentric retrospective review of 422 patients with CNS lymphoma, both primary and secondary, in which only 2.5% had mucosa-associated lymphoid tissue (MALT) lymphoma by histology, the majority of which presented with a dural mass lesion.

Houllier [11] in a small retrospective case series showed that lenalidomide has activity in multiply recurrent PCNSL.

Diletto *et al.* [12] reported that in PCNSL treatment with hippocampal sparing whole brain irradiation, a technique designed to spare late treatment related loss of memory, was not feasible in PCNSL due to the diffuse and multifocal nature of the disease as well as frequent proximity of lymphoma to the hippocampus.

• Meningiomas

Mazzetti *et al.* [13] demonstrated by aquaporin-4 (AQP-4) immunohistochemical expression tumor infiltration into brain in an analysis of meningioma (n = 35) of all grades. These results suggest that AQP-4 may be a marker for biological behavior of meningioma.

Magistrello *et al.* [14] treated 17 patients with surgery and radiation refractory recurrent meningioma with a monthly long-acting somatostatin analogue after documenting overexpression of somatostatin receptor 2 (sstr2) by octreotide scintigraphy. Toxicity was manageable and progression free survival at 6 months (PFS-6) was 56%. These results suggest modest activity for use of somatostatin analogues in the treatment of recurrent meningioma.

Rachinger *et al.* [15] using thesstr2 analogue 68-Ga-DOTATATE as a positron emission tomography (PET) tracer showed superior detection of meningioma compared with anatomic MRI suggesting utility in surgical planning, evaluating extent of resection and radiotherapy (RT) treatment planning.

• Gliomas

Low-grade gliomas

Bertero *et al.* [16] in a small study of low-grade gliomas (LGG) clinically believed to require treatment after biopsy or partial resection were treated with a dose dense temozolomide (TMZ) regimen for 6 months to assess down staging by chemotherapy. There were no objective responses; however, one minimal response by RANO criteria was observed as well as a median 26% MRI FLAIR volume reduction. Additionally tumor infiltration as determined by MRI diffusion weighted images suggested a decrease in tumor infiltration.

Cahill *et al.* [17] showed the feasibility of performing ¹H MRI spectroscopy to interrogate isocitrate dehydrogenase (IDH) positive LGG for the oncometabolite 2-hydroxyglutamate (2HG). Quantifying 2HG may allow for another method to assess response to treatment including studies evaluating IDH small molecule inhibitors.

In another imaging study of LGG, Izquierdo *et al.* [18] demonstrated by fluid attenuation inversion recovery (FLAIR) volumetric MRI the ability to predict anaplastic transformation by a change in tumor volume assessed by comparing volumes at 6-month intervals.

Calatozzolo *et al.* [19] examined alpha thalassemia/mental retardation syndrome X-linked (ATRX) and telomerase reverse transcriptase promoter (TERTp) gene mutations in 30 LGG. Both genes code for two differing mechanisms for maintenance of telomerase length facilitating immortalization of cancer cells. ATRX mutations co-associated with IDH and p53

mutations whereas TERTp mutations associated with 1p19q codeletion.

Chan *et al.* [20] assessed TERTp mutations in 181 low grade and anaplastic gliomas. TERTp mutations were seen in 14% of astrocytomas, 67% of oligoastrocytomas and 100% of oligodendrogliomas. TERTp mutations were strongly associated with 1p19q codeletion, IDH mutations and oligodendroglial histology as well as improved survival. In nondeleted LGG, TERTp mutations were also associated with improved survival relative to TERTp wild-type tumors. These results suggest analysis of TERTp mutational status contributes to identifying prognostic subgroups in LGG.

Gleize *et al.* [21] showed that mutations in the *capicua* transcriptional repressor gene (CIC) on chromosome 19q and seen in approximately 50% of all codeleted oligodendrogliomas predicts for shorter survival and time to anaplastic transformation. These findings suggest CIC mutations may be prognostic.

Lastly, Sahm *et al.* [22] analyzed 42 oligoastrocytoma and concluded none of these mixed tumors have coexisting cell populations but rather are separable into two mutually exclusive categories based on genotype; ATRX mutated astrocytic tumors or 1p19q codeleted oligodendroglial tumors. This molecular analysis suggests that an oligoastrocytoma is not an independent entity.

Anaplastic gliomas

Upfront trials

Wick *et al.* [23] reanalyzed prognostic biomarkers in the German NOA-04 trial. The study concluded the *ATRZ* mutation is a lineage marker of astrocytic differentiation as well as a favorable prognostic indicator in anaplastic gliomas (AG). Furthermore, methylguanine methyltransferase promoter methylation is a predictive marker only in IDH wild-type AG wherein there is a survival advantage when treated with an alkylator chemotherapy compared with RT.

Figarella-Branger *et al.* [24] showed in an analysis of 2013 anaplastic oligodendroglioma (AO) enrolled in the French National network POLA, 79% of tumors were codeleted for 1p19q. Amongst these tumors, three subcategories could be defined by the presence or absence of necrosis, microvascular proliferation and the mitotic index. The least favorable outcome was seen in AO with a high mitotic index

and presence of both necrosis and microvascular proliferation.

Glioblastoma

Upfront trials

Herrlinger *et al.* [25] presented QoL data on the German GLARIUS trial that compared standard RT + TMZ followed by post-RT TMZ to RT + bevacizumab followed by post-RT bevacizumab + Irinotecan in newly diagnosed glioblastoma (GBM) that were methylguanine methyltransferase promoter unmethylated. Though improvement was seen in PFS in the experimental arm, there were no differences in overall survival or in QoL.

Reardon and Nayak *et al.* [26] presented a new RANO instrument for standardizing the neurological examination in trials involving patients with brain tumors. The instrument (currently in two industry sponsored trials but as yet not validated) utilizes eight relevant neurological domains specific to brain that is both quantifiable and clinician friendly.

Salvage trials

Duerinck *et al.* [27] reported in a small (n = 44) randomized trial comparing the vascular endothelial GFR2 oral inhibitor axitinib to bevacizumab in TMZ refractory recurrent GBM. There were no overall survival differences and axitinib was found to be more toxic. Nonetheless, the authors conclude axitinib has single agent activity in GBM.

Trevisan *et al.* [28] conducted a small (n = 54) single arm Phase II study of fotemustine + bevacizumab in recurrent GBM. Overall survival was not different than single agent bevacizumab based on a historical comparison suggesting no added advantage to partnering bevacizumab with fotemustine, a nitros urea chemotherapy. Additionally median post-bevacizumab progression survival was reported as 4.5 months with continuation of bevacizumab and only 2.2 months with best supportive care.

Taal *et al.* [29] represented the Dutch BELOB randomized Phase II trial comparing bevacizumab to lomustine to bevacizumab + lomustine. Based on a prespecified end point of overall survival at 9 months, only the combination of bevacizumab + lomustine met the criteria for further study (currently being evaluated in an EORTC randomized Phase III trial). Notably there was little to no cross over in the single agent treatment arms eliminating a confounding second salvage therapy.

Pathology

Berghoff *et al.* [30] analyzed 117 patients with GBM for expression of the immune check-point receptor (PD1) and ligand (PD-L1) using immunohistochemistry. PD1 was expressed as expected only in TILs and in 15% of tumors overall whereas PD-L1 was expressed in 44% of GBM cells. These results suggest GBM may be responsive to immune check point inhibition and clinical studies have been initiated.

Spiegel-Kreinecker *et al.* [31] assessed TERTp mutations in 126 patients with GBM and showed that in patients with TERTp mutations shorter survival suggesting another potential prognostic marker in GBM.

Labussiere *et al.* [32] performed a combined analysis of TERTp, IDH and epidermal GFR (EGFR) status in 395 GBM. The results suggest several novel molecular categories in GBM. TERTp mutation predicts for poor outcome regardless of IDH status. TERTp mutation

co-associates with IDH wild-type tumors and EGFR amplification. EGFR amplification was unassociated with survival in isolation however correlated with survival when assessed with TERTp. TERTp mutated tumors combined with EGFR amplification showed longer survival as did TERTp wild-type tumors combined with EGFR nonamplified. The authors conclude the analysis of TERTp, EGFR and IDH refines the prognostic classification of GBM.

Financial & competing interests disclosure

The author has no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.

No writing assistance was utilized in the production of this manuscript.

References

- Kwakman RJJM, Jeene PM, Dieleman E, Stalpers LJA. Estimation of the incidence of brain metastases and patterns of care in The Netherlands. *Neuro Oncol.* 16(Suppl. 2), Abstract O1.01 (2014).
- Wen PY, Lee EQ, Van Den Bent M *et al.* Response assessment in neuro-oncology (RANO) criteria for brain metastases. *Neuro Oncol.* 16(Suppl. 2), Abstract O7.01 (2014).
- Berghoff AS, Wöhrer A, Hackl M *et al.* Density of tumor infiltrating lymphocytes correlates with brain edema and overall survival in brain metastases. *Neuro Oncol.* 16(Suppl. 2), Abstract O1.03 (2014).
- Galldiks N, Ruge MI, Hamisch C *et al.* Differentiation of local recurrent brain metastasis from radiation-induced changes using O-(2-[18F]Fluoroethyl)-L-tyrosine PET. *Neuro Oncol.* 16(Suppl. 2), Abstract O7.05 (2014).
- Scoccianti S, Greto D, Bordi L *et al.* Radiosurgery and brain metastases: adequate sequence of brain MRI can significantly change the intracranial disease staging. *Neuro Oncol.* 16(Suppl. 2), Abstract O7.02 (2014).
- Habets EJJ, Dirven L, Wiggeraad R *et al.* Cognitive functioning, health-related quality of life and survival after stereotactic radiotherapy in patients with brain metastases. *Neuro Oncol.* 16(Suppl. 2), Abstract O9.05 (2014).
- Bionski M, Wittwer B, Faure G *et al.* Identification and qualification of CSF malignant cells by the CellSearch® technology in patients with lung leptomeningeal metastasis. *Neuro Oncol.* 16(Suppl. 2), Abstract P08.02 (2014).
- Chamberlain M, Junk L, Bradsma D *et al.* Leptomeningeal metastases: a RANO proposal for response criteria. *Neuro Oncol.* 16(Suppl. 2), Abstract P08.27 (2014).
- Ahle G, Hoang-Xuan K, Cassoux N *et al.* Optic nerve infiltration of primary CNS lymphoma. *Neuro Oncol.* 16(Suppl. 2), Abstract P11.03 (2014).
- Larrieu D, Choquet S, Hoang-Xuan K *et al.* MALT CNS lymphomas: a monocentric experience. *Neuro Oncol.* 16(Suppl. 2), Abstract P11.04 (2014).
- Houllier C, Choquet S, Toutou V *et al.* Lenalidomide in the treatment of relapsed primary central nervous system lymphoma (PCNSL). *Neuro Oncol.* 16(Suppl. 2), Abstract P11.05 (2014).
- Diletto B, Marchesano D, Chiesa S *et al.* May we routinely spare hippocampal region in primary central nervous system lymphoma during whole brain radiotherapy? *Neuro Oncol.* 16(Suppl. 2), Abstract O2.05 (2014).
- Mazzetti S, Calatuzzolo C, Patanè M *et al.* Novel markers if meningioma aggressiveness: a study of meningioma versus peritumoral nervous tissue. *Neuro Oncol.* 16(Suppl. 2), Abstract O1.06 (2014).
- Magistrello M, Trevisan E, Bertero L *et al.* Systemic treatment of meningiomas: is there a role for long-acting somatostatin? *Neuro Oncol.* 16(Suppl. 2), Abstract P06.08 (2014).
- Rachinger W, Stöckli V, Huang AR *et al.* [⁶⁸GA]-DOTATATE-PET imaging predicts somatostatin receptor II in meningiomas. *Neuro Oncol.* 16(Suppl. 2), Abstract O7.09 (2014).
- Bertero L, Rudà R, Castellano A *et al.* Pre-operative chemotherapy as a new strategy in treating low grade gliomas in eloquent areas. *Neuro Oncol.* 16(Suppl. 2), Abstract O8.01 (2014).
- Cahill DP, Loebel F, Bogner W *et al.* Monitoring of treatment response in IDH-mutant gliomas with in-vivo 3D magnetic resonance spectroscopy. *Neuro Oncol.* 16(Suppl. 2), Abstract O8.02 (2014).
- Izquierdo C, Majos C, Simó M *et al.* Low grade diffuse astrocytoma: prognostic factors to anaplastic transformation. *Neuro Oncol.* 16(Suppl. 2), Abstract O8.03 (2014).
- Calatuzzolo C, Patanè M, Mazzetti M *et al.* ATRX and TERT expression in relapsing low grade glioma. *Neuro Oncol.* 16(Suppl. 2), Abstract O8.06 (2014).
- Chan AK, Yao Y, Zhang Z *et al.* TERT promoter mutations contribute to subset prognostication of lower grade gliomas. *Neuro Oncol.* 16(Suppl. 2), Abstract P04.06 (2014).
- Gleize V, Alentorn A, Connen de Kérillis L *et al.* CIC mutation as a poor prognostic

- factor in 1p19q codeleted gliomas, associated to an up-regulation of proliferation pathways. *Neuro Oncol.* 16(Suppl. 2), Abstract O8.08 (2014).
- 22 Sahm F, Reuss D, Koelsche C *et al.* Oligoastrocytoma does not exist: a molecular genetic reappraisal on single-cell level. *Neuro Oncol.* 16(Suppl. 2), Abstract P04.22 (2014).
- 23 Wick W, Capper D, Sill M *et al.* Prognostic and predictive biomarker-based subgroups in the NOA-04 trial. *Neuro Oncol.* 16(Suppl. 2), Abstract O10.01 (2014).
- 24 Figarella-Branger D, Mokhtari K, Dehais C *et al.* Mitotic index, microvascular proliferation and necrosis define three groups of IP/19Q co-deleted anaplastic oligodendrogliomas associated with different genomic alterations. *Neuro Oncol.* 16(Suppl. 2), Abstract O10.02 (2014).
- 25 Herrlinger U, Schäfer N, Steinbach JP *et al.* The randomized, multicenter GLARIUS trial investigating bevacizumab/irinotecan vs standard temozolomide in newly diagnosed, MGMT-non-methylated glioblastoma patients: final survival results and quality of life. *Neuro Oncol.* 16(Suppl. 2), Abstract O10.04 (2014).
- 26 Reardon DA, Nayak L, DeAngelis LM *et al.* The neurologic assessment in neuro-oncology (NANO) scale: a tool to assess neurologic function for integration in the radiologic assessment in neuro-oncology (RANO) criteria. *Neuro Oncol.* 16(Suppl. 2), Abstract P15.3 (2013).
- 27 Duerinck J, DuFour S, Bouttens F *et al.* Randomized Phase II study of axitinib vs. standard of care in patients with recurrent glioblastoma. *Neuro Oncol.* 16(Suppl. 2), Abstract O10.07 (2014).
- 28 Trecisan E, Bertero L, Cassoni P *et al.* Phase II study of AINO (Italian Association of Neuro-Oncology) on bevacizumab and fotemustine for recurrent glioblastoma: outcome, patterns of relapse and salvage treatment. *Neuro Oncol.* 16(Suppl. 2), Abstract O10.06 (2014).
- 29 Taal W, Oosterkamp HM, Walenkamp AME *et al.* Final analysis of the BELOB trial (a randomized Phase II study on bevacizumab versus bevacizumab plus lomustine versus lomustine single agent in recurrent glioblastoma) and first radiology results. *Neuro Oncol.* 16(Suppl. 2), Abstract O10.05 (2014).
- 30 Berghoff AS, Kiesel B, Widhalm G *et al.* Analysis and PD-L1 expression in glioblastoma. *Neuro Oncol.* 16(Suppl. 2), Abstract O6.02 (2014).
- 31 Spiegl-Kreinecker S, Loetsch D, Laaber M *et al.* Mutations of the TERT promoter correlate with enhanced telomerase activation and predict a worse prognosis on primary glioblastoma patients. *Neuro Oncol.* 16(Suppl. 2), Abstract P04.24 (2014).
- 32 Labussiere M, Boiselier B, Mokhtari K *et al.* Combine analysis of TERT, EGFR, and IDH status define distinct prognostic classes of glioblastoma. *Neuro Oncol.* 16(Suppl. 2), Abstract O6.08 (2014).