Our panel of experts highlight the most important research articles across the spectrum of topics relevant to the field of CNS oncology

Expert panel: Larry Junck, University of Michigan Health System, MI, USA; Maura Massimino, Fondazione Istituto di Ricovero & Cura a Carattere Scientifico Istituto Nazionale dei Tumori, Milan, Italy

Meyer M, Reimand J, Lan X *et al.* Single cell-derived clonal analysis of human glioblastoma links functional and genomic heterogeneity. *Proc. Natl Acad. Sci. USA* 112, 851–856 (2015).

Investigators from the laboratory of Dr Peter Dirks analyzed 44 clones derived from single cells of four glioblastomas, two treatment-naïve and two recurrent, using genetic profiling, transcription profiling, cell survival analysis after drug treatment, and other techniques. They found striking differences in clones from individual tumors including heterogeneous expression of PTEN and of EGF receptor (EGFR) and its VIII variant. Cells were sensitive to temozolomide in six of eight clones from one naïve tumor and one of ten clones from a previously resistant tumor, while clones from the other two tumors were resistant. Sensitivity *in vitro* was unrelated to O6-methylguanine-DNA methyltransferase (MGMT) protein expression or promoter methylation. In the recurrent tumor with one sensitive clone, 28 known cancer genes were differentially expressed in that clone, including reduced expression of *EGFR* and *MET*, compared with resistant clones. Sensitivity to a panel of 98 other drugs also showed clone-to-clone variability within tumors. The impact of stem cell culturing on genetic stability was negligible compared with heterogeneity between clones. Using powerful techniques, the authors have shown substantial variability in gene copy number, gene amplification and expression and drug sensitivity in glioblastoma, complicating treatment of these tumors.

Written by Larry Junck

Funato K, Major T, Lewis PW, Allis CD, Tabar V. Use of human embryonic stem cells to model pediatric gliomas with H3.3K27M histone mutation. *Science* 346, 1529–1533 (2014).

Over 70% of diffuse intrinsic pontine gliomas (DIPGs) bear H3.3K27M mutations in histone H3.3, but the role of this mutation in tumorigenesis is not well understood. The authors transduced neural progenitor cells from human embryonic stem cells with combinations of a K27M-mutant form of H3.3, a constitutively active form of PDGFRA, and a small hairpin RNA against p53. The latter two genes were chosen because of their high frequency of overexpression or mutation in K27M-mutated DIPGs. *In vitro*, H3.3K27M was associated with increased H3K27 trimethylation and with increased proliferation. The H3.3K27 mutation was synergistic with the PDGFRA transgene and sh-p53 in inducing a neoplastic phenotype, as shown by radiation resistance *in vitro* and tumor formation in mouse brainstem. Screening of model cells *in vitro* with compounds that target epigenetic regulators led to preliminary evidence of efficacy of the menin inhibitor MI-2 *in vitro* and *in vivo*. The authors conclude that H3.3K27M is a driver mutation in the appropriate developmental window, that its effects are synergistic with other genetic abnormalities, and that this platform is useful for investigation of the biology and treatment of DIPGs.

Written by Larry Junck

Houillier C, Choquet S, Touitou V *et al.* Lenalidomide monotherapy as salvage

treatment for recurrent primary CNS lymphoma. *Neurology* 84, 325–327 (2015).

For patients with primary CNS lymphoma (PCNSL) whose tumors progress after high-dose methotrexate-based chemotherapy, treatment options are unsatisfactory. Lenalidomide, an immunomodulatory drug derived from thalidomide, has been shown to be effective against multiple myeloma and certain types of systemic lymphomas. The authors treated six patients with relapsed PCNSL, median age 74 years, median Karnofsky performance status 60, with oral lenalidomide, 25 mg/day on days 1–21 of a 28-day schedule. Responses by MRI were complete in two (duration 24+ and 3 months) and partial in one; three patients had progressive disease. One patient died suddenly of unknown cause, but there were no other grade III–IV toxicities. Although the number of patients is small and two of the three responses were of short duration, this study is of value because it demonstrates activity of a drug from a class not previously known to be active against PCNSL.

Written by Larry Junck

Mistry M, Zhukova N, Merico D *et al.* BRAF mutation and CDKN2A deletion define a clinically distinct subgroup of childhood secondary high-grade glioma. *J. Clin. Oncol.* doi:10.1200/JCO.2014.58.3922 (2015) (Epub ahead of print).

While a great majority of adult low-grade glioma transforms during time into high-grade subtypes, this occurrence is exceedingly rare in children and is more feared, especially by nonspecialized physicians, than described in really documented series, when not including patients already pretreated with radiotherapy or affected by procancer syndrome. Recent next-generation sequencing efforts have clearly defined the different mutations characterizing pediatric and adolescent glioma with respect to the adult counterpart. Pediatric primary high-grade glioma (HGG) and pediatric low-grade glioma (PLGG), are characterized by somatic mutations in histone H3.3, ATRX and TP53 genes in the first case, and almost always by alterations causing activation of the MAPK/ERK pathway, in the second case. In this wide analysis of Canadian patients during an almost thirty year- time span, of 886 patients treated for PLGG, 26 patients (2.9%) fulfilled the criteria for malignant transformation, the majority

being pilocytic astrocytoma at first diagnosis. Five patients, being already pretreated with irradiation, that is a well-known mutation agent, and three having an established Li-Fraumeni syndrome, should have been excluded by the report giving a total of 18 ‘spontaneously’ transformed PLGG. Exome sequencing and array CGH were performed on available samples followed by detailed genetic analysis of the entire secondary HGG cohort demonstrating that the most recurrent alterations were BRAF V600E and CDKN2A deletion. sHGGs exhibited a higher somatic mutation load than primary pediatric HGGs data available from literature. Analysis of patient-matched low- and high-grade samples revealed moreover that BRAF mutations identified in the sHGG were also identified in 100% of the corresponding PLGGs and that 80% of sHGG CDKN2A deletions were also identified in the corresponding PLGGs. The BRAF V600E mutant PLGGs had also a longer latency to transformation and a better prognosis than wild-type transformed PLGG thus also demonstrating a unique clinical profile. Since both BRAF V600E and CDKN2A-deleted tumors may be eligible for targeted therapies, the authors recommend that preventive measures, including a more aggressive surgical approach and consideration of medical therapies even when a stable tumor is observed, may be able to mitigate the devastating transformation event. It will be important also to test in the future how many incompletely resected tumors that harbor both alterations will not transform with long clinical follow-up.

Written by Maura Massimino

Eisenstat DD, Pollack IF, Demers A *et al.* Impact of tumor location and pathological discordance on survival of children with midline high-grade gliomas treated on Children’s Cancer Group high-grade glioma study CCG-945. *J. Neurooncol.* 121, 573–581 (2015).

The CCG-945 study compared the regimen known as ‘8-drugs-in-1-day’ (‘8-in-1’), given both pre- and postirradiation, to the chemotherapy-containing arm in CCG-943 that was vincristine, prednisolone and lomustine. This study, being the widest randomized for malignant gliomas so far published, represents a good reservoir of patients where it is possible to study different prognostic factors. In this paper the

authors come back to tumor location, that was midline in 60/172 eligible children, to describe this subgroup patients outcome and difficulties in precise pathological diagnosis. At the end of their analyses, including also some biological markers (MGMT, p53 and Ki67 expression), when compared with the overall group, the midline location had poorer survival, discordant diagnoses were evident in more than 30% of patients, MGMT overexpression was correlated to very poor prognosis and rate of complete resections was very low. All these elements contributed to an awful outcome.

Written by Maura Massimino

Khajuria RK, Blankenburg F, Wuithschick I et al. Morphological brain lesions of pediatric cerebellar tumor survivors correlate with inferior neurocognitive function but do not affect health-related quality of life. *Childs Nerv Syst*. doi:10.1007/s00381-015-2635-4 (2015) (Epub ahead of print).

The authors describe the extent of morphological brain injury in pediatric cerebellar tumor survivors – 17 after pilocytic astrocytoma (PA) and 17 after medulloblastoma (MB) – and its correlation with neurocognitive function and health-related quality of life (HrQoL). The posterior fossa harbors the cerebellum, containing almost half of all neurons of the central nervous system therefore any damage in this site can compromise fundamental functions like motor, executive, attention and working memory that constitute the performance in intelligence. Moreover, also behavioral problems like social withdrawal and anxiety can arouse after cerebellar damage. In this report describing different morphological alterations appreciated through MRI according to the treatment received – mostly surgery only by patients with PA, and surgery, craniospinal irradiation and chemotherapy by patients with MB – the authors come to very interesting conclusions about neurocognitive outcome and

HrQoL. MB patients had lower results in all intelligence tests, attention, verbal memory and visuo-spatial integration, but also PA patients had lower scores in performance IQ, alertness and fine motor functions, decreased verbal memory concerning learning and interference when the volume of the cerebellar lesion increased. Despite these objective differences from normal pairs, both MB and PA patients had better results in HrQL if compared with a reference group of children with chronic illnesses. Both cerebellar tumor patient groups were found to suffer from significantly reduced attention, deficits in visual motor integration, auditory learning and reduced IQ. Almost all neurocognitive deficits were more pronounced in MB patients than in PA patients, due to the sum of chemotherapy and radiotherapy administered after surgery. In PA patients, where cerebellar loss was more profound, radiologic parameters allowed to suggest that cerebellar damage in PA patients is involved in their cognitive and affective development, in other words, higher-order function. PA patients had also lower scores compared with healthy peers. Suffering from a malignant disease and from the long duration of treatment, one would have expected a clearly significant reduction in HrQoL. Psychological support, rehabilitation measurements and sufficient time to adapt to the disease between time of diagnosis and time of evaluation could influence this parameter in a positive way.

Written by Maura Massimino

Financial & competing interests disclosure

The authors have no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.

No writing assistance was utilized in the production of this manuscript.