About the Editor

Christopher R Chapple

Christopher R Chapple is a Consultant Urological Surgeon at Sheffield Teaching Hospitals NHS Foundation Trust, UK. He has a long-standing interest in the assessment of lower urinary tract dysfunction and functional reconstruction. He has published more than 350 articles in peer-reviewed literature dealing with urodynamic assessment of the lower urinary tract, pharmacotherapy and surgical reconstruction of the lower urinary tract. He is currently Editor of the journal *Neurourology & Urodynamics*, and serves on the editorial board of several major journals. He is a member of many international societies and also a member of the American Association of Genito-Urinary Surgeons. His research interests continue in the field of functional assessment of the urinary tract, pharmacology of the lower urinary tract, functional restoration of normal function and he has a particular interest in the etiology and treatment of overactive bladder symptom complex and associated detrusor overactivity.

For reprint orders, please contact: reprints@futuremedicine.com

Foreword

Contemporary management of overactive bladder

Christopher R Chapple

It is a great pleasure to introduce this book, which deals with the subject of overactive bladder. I would like to emphasize the enormous amount of work that all of my colleagues have put into the production of this book and I hope that you will find it a very useful adjunct to your clinical practice.

In the five chapters that comprise this work, we explore all of the pertinent issues relating to the clinical management of overactive bladder in contemporary practice.

Nadir Osman and I review the epidemiology of overactive bladder, emphasizing that following on from its initial description in the 2002 standardization report from the International Continence Society [1], this symptom complex is now well-recognized as an important clinical entity that produces significant morbidity with implications for patients' quality-of-life.

Robinson and Cardozo in their chapter, clearly emphasize the need to adequately evaluate patients. They review existing guidelines and the potential role of frequency volume charts and urodynamic assessment are considered in detail. Having excluded significant uropathology with urinalysis, it is important to realize that frequency volume charts should be used in all patients prior to instituting therapy to ensure that the therapy is directed in the most appropriate fashion. A frequency volume chart will, for instance, identify whether a patient can improve symptoms by modifying fluid intake, or whether pharmacotherapy or more invasive treatment is

necessary. In cases that fail to respond to conservative management, such as lifestyle modification and pharmacotherapy, then more detailed assessment is appropriate prior to proceeding on to second-line therapeutic options, which are more invasive and may have more significant consequences for patients.

It is clear that overactive bladder symptom complex is a very common condition, affecting populations across the world and leading to an enormous interference with quality of life as a consequence of the associated morbidity that these symptoms cause. Bhide and Khullar emphasize this, looking at lifestyle modification and how this can help in the management of these very troublesome and intrusive symptoms, which is often ignored by clinicians. It is clear from this comprehensive overview, that attention to many factors associated with lifestyle can have a very significant impact on the storage symptoms associated with overactive bladder.

In their chapter, Gomelsky and Dmochowski review the use of the current 'gold standard' pharmacotherapy for overactive bladder symptom complex, namely antimuscarinics. These agents have been in widespread clinical use for over 30 years and are efficacious, but are limited in their usefulness as a consequence of efficacy and tolerability issues. Nevertheless, they have brought considerable benefit to many patients, albeit the evidence for their long-term use in real-life clinical practice is very limited and rather deficient. The widely used agents are very comprehensively reviewed by these authors in this very practical chapter. It will be of interest to see what the development of β_3 agonists contributes to the management of overactive bladder in real-life clinical practice. The first of these agents, mirabegron, has been introduced into routine clinical practice in Japan, the application for its use has been accepted by the US FDA and it is currently being evaluated by the EMA.

Sajadi, Goldman and Staskin comprehensively review the use of sacral nerve stimulation and percutaneous tibial nerve stimulation, both of which have been introduced into clinical practice as second-line treatment for patients who fail to respond to lifestyle changes and pharmacotherapy. This very practical overview contrasts these therapies and emphasizes the pros and cons of therapy; in particular, pointing out that there are significant cost issues with this therapy. Clearly in the future, more detailed cost-benefit analyses need to be carried out in order to help us to better define how to most effectively treat our patients.

A new class of second-line therapy is also currently being introduced, namely the use of botulinum toxin treatment injected intravesically. There

is a lot of debate over how this contrasts both clinically and in terms of efficacy and also cost–benefit, to electrical stimulation techniques. Currently, licensing authorization has been granted by the FDA and a number of other countries across the world for its utilization, using the agent onabotulinumtoxinA in the management of neurogenic bladder overactivity, and a Phase III trial program has been completed for the use of this agent in treating patients with idiopathic overactive bladder symptom syndrome. This agent has the benefit of being easy to administer, albeit requiring cystoscopically guided injections, but does last for up to 6–8 months following each treatment. Patients need to be warned that certainly when used for idiopathic overactive bladder symptom complex, it is not yet licensed for this use. The major adverse events relate to retention in a proportion of patients (at a dose of 100 units, being less than ~5% based on our current knowledge base) and a number of patients do experience symptoms suggestive of a urinary tract infection, although this may not always be confirmed on culture despite the presence of symptoms. Patients, therefore, also need to be prepared to use intermittent self catheterization if necessary following application of this treatment.

Financial & competing interests disclosure

CR Chapple is a Consultant to AMS and Lilly; Consultant and Researcher to ONO; Consultant, Researcher and Speaker to Allergan, Astellas, Pfizer and Recordati. The author has no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Reference

- 1 Abrams P, Cardozo L, Fall M *et al.* The standardisation of terminology of lower urinary tract function: report from the Standardisation Subcommittee of the International Continence Society. *Neurourol. Urodyn.* 21(2), 167–178 (2002).