About the Editors

Nicola Dalbeth

Nicola Dalbeth is a Consultant Rheumatologist and Associate Professor of Medicine at the University of Auckland (New Zealand). She is a Principal Investigator in the Auckland Bone and Joint Research Group. She leads a clinical and laboratory research program in gout, focusing on the mechanisms of inflammation and joint damage in chronic gout.

Fernando Perez-Ruiz

Fernando Perez-Ruiz is a Senior Clinician at the Hospital Universitario Cruces (Biscay, Spain). His clinical research over the last 20 years has been focused on treatment, outcome measures, audits and recommendations for gout.

Naomi Schlesinger

Naomi Schlesinger is Chief of the Division of Rheumatology and Rheumatology Fellowship Program Director at the University of Medicine and Dentistry of New Jersey, Robert Wood Johnson Medical School (USA), where she is also Professor of Medicine in the Department of Medicine. She is a noted authority in the field of gout, having published papers regarding the diagnosis, treatment and better understanding of the pathogenesis of gout. She is the author of over 190 scientific articles, abstracts, book chapters and reviews. She is the past President of the New Jersey Rheumatology Association. In addition she is Co-Chair of the American College of Rheumatology Crystal Study Group, as well as the Co-Chair of the American College of Rheumatology Abstract Review: Metabolic and Crystal Arthropathies. She has a special interest in evidence-based medicine and serves as a cofacilitator of the Acute Gout Review Group for Cochrane International Collaboration as well as a member of the Outcome Measures in Rheumatology Gout Special Interest Organizing Group.

For reprint orders, please contact: reprints@futuremedicine.com

Foreword

Gout

Nicola Dalbeth, Fernando Perez-Ruiz
& Naomi Schlesinger

In the last decade, we have witnessed exciting advances in the understanding and treatment of gout. Gout is now the most common inflammatory arthritis, affecting almost 4% of the US population, with increasing rates in the last two decades [1]. The impact of poorly controlled gout on health-related quality of life, disability, work, participation and healthcare utilization has been highlighted, emphasizing the need for improved management of this condition. We have also seen new understanding of the pathophysiology of gout, particularly the genetics of hyperuricemia, environmental risk factors and the underlying immunological basis of this inflammatory disease. The complex relationship between hyperuricemia/gout and comorbid conditions has been explored. Imaging methods, particularly ultrasonography and dual energy computed tomography have allowed us to understand the mechanisms of disease in new ways. Large international collaborations have led to important progress in definitions, outcome measures for clinical trials and the genetic basis of gout [2–4].

The application of these scientific discoveries is leading to new therapeutic strategies to combat both acute gout flares and prevent the consequences of poorly controlled disease. The identification that activation of the NLRP3 inflammasome and subsequent release of mature IL-1 β is central to the initiation phase of acute gout inflammation [5] has led to the identification of anti-IL-1 biologic therapy for treatment and prophylaxis

of acute flares [6]. Two new urate-lowering agents have been approved by the US FDA in the last 5 years, febuxostat and pegloticase [7,8], and a number of other novel urate-lowering therapies are in Phase III clinical trials. In addition to these new agents, the optimal use of ‘veteran’ drugs such as colchicine and allopurinol has been clarified. Gout management guidelines have been published by the major rheumatology societies, including the European League Against Rheumatism, British Society for Rheumatology and American College of Rheumatology. Central to all of these guidelines is the importance of ‘treating to serum urate target’, a key strategy to address the underlying basis of disease and prevent the consequences of poorly controlled gout.

This book highlights many of the advances in understanding mechanisms of disease, and provides a practical framework for gout management, outlining the therapeutic principles of gout management and providing detailed information about specific agents. Current controversies are explored, particularly issues around the relationship between serum urate and comorbid conditions, and management of ‘asymptomatic’ hyperuricemia.

Despite the major advances described in this book, the quality of gout treatment is frequently poor. Understanding barriers to effective gout care and strategies to address these barriers is required to translate progress into therapeutic benefits for the patient. We hope that this book will provide a useful resource for researchers and clinicians, and ultimately lead to improved care of patients with gout.

Financial & competing interests disclosure

N Dalbeth has received consulting fees from Takeda, Ardea, Novartis, Fonterra and Metabolex; speaker’s fees from Novartis and Savient; and grant support from Fonterra. F Perez-Ruiz discloses consultancies/speakers bureau/educational materials/advisory board for Ardea, Menarini, Metabolex, Novartis and Savient. N Schlesinger reports having received a grant, travel expenses and payment for advisory board membership from Novartis Pharma, payment for advisory board membership and educational presentations from Takeda and Savient, and payment for advisory board membership from Savient, URL Pharma and Enzyme Rx. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

References

- 1 Zhu Y, Pandya BJ, Choi H. Prevalence of gout and hyperuricemia in the US general population: the National Health and Nutrition Examination Survey 2007–2008. *Arthritis Rheum.* 63(10), 3136–3141 (2011).
- 2 Gaffo AL, Schumacher HR, Saag KG *et al.* Developing a provisional definition of flare in patients with established gout. *Arthritis Rheum.* 64(5), 1508–1517 (2012).
- 3 Schumacher HR, Taylor W, Edwards L *et al.* Outcome domains for studies of acute and chronic gout. *J. Rheumatol.* 36(10), 2342–2345 (2009).
- 4 Kolz M, Johnson T, Sanna S *et al.* Meta-analysis of 28,141 individuals identifies common variants within five new loci that influence uric acid concentrations. *PLoS Genet.* 5(6), e1000504 (2009).
- 5 Martinon F, Pétrilli V, Mayor A, Tardivel A, Tschopp J. Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* 440(7081), 237–241 (2006).
- 6 So A, De Meulemeester M, Pikhak A *et al.* Canakinumab for the treatment of acute flares in difficult-to-treat gouty arthritis: results of a multicenter, Phase II, dose-ranging study. *Arthritis Rheum.* 62(10), 3064–3076 (2010).
- 7 Becker MA, Schumacher HR Jr, Wortmann RL *et al.* Febuxostat compared with allopurinol in patients with hyperuricemia and gout. *N. Engl. J. Med.* 353(23), 2450–2461 (2005).
- 8 Sundry JS, Baraf HS, Yood RA *et al.* Efficacy and tolerability of pegloticase for the treatment of chronic gout in patients refractory to conventional treatment: two randomized controlled trials. *JAMA* 306(7), 711–720 (2011).