Future MICROBIOLOGY

December 2017 Volume 12 Issue 16

Editorial

- 1445 Insights into the evolution of the mutational resistome of *Pseudomonas aeruginosa* in cystic fibrosis
C López-Causape & A Oliver

Short Communication

- 1449 Protective efficacy of antidiarrheal agents in a permeability model of *Escherichia coli*-infected CacoGoblet® cells
B de Servi & F Ranzini

Research Articles

- 1457 Matrix-assisted laser desorption/ionization–time of flight mass spectrometry: protocol standardization and database expansion for rapid identification of clinically important molds
S Paul, P Singh, SM Rudramurthy, A Chakrabarti & AK Ghosh
- 1467 γ -irradiation from radiotherapy improves the virulence potential of *Candida tropicalis*
EM da Silva, B Kischkel, CS Shinobu-Mesquita, PS Bonfim-Mendonça, ESB Mansano, MA da Silva, JF Barbosa, A Fiorini, L Hernandez, MC Furlaneto & TIE Svidzinski

- 1487 Carvacrol is highly disruptive against coagulase-negative staphylococci in *in vitro* biofilms
V Gaio, CA Lima, F Oliveira, A França & N Cerca

- 1499 Histone acetyltransferase encoded by *NGG1* is required for morphological conversion and virulence of *Candida albicans*
D-D Li, BB Fuchs, Y Wang, X-W Huan, D-D Hu, Y Sun, D Chai, Y-Y Jiang & E Mylonakis

- 1513 Comparative genomics of type 1 IncC plasmids from China
L Ma, Z Yin, D Zhang, Z Zhan, Q Wang, X Duan, H Gao, Q Liang, Y Zhao, J Feng, Y Zhao, E Dai & D Zhou

Special Report

- 1525 Susceptibility testing and reporting of new antibiotics with a focus on tedizolid: an international working group project
MH Wilcox, N Dmitrieva, AC Gales, I Petukhova, S Al-Obeid, F Rossi & JM Blondeau

Review

- 1535 The hidden pathogenic potential of environmental fungi
GR de S Araújo, W de Souza & S Frases

Future MICROBIOLOGY

Editorial Board

The Editorial Board is drawn from the leading forces in research and academia

Senior Editors

Richard Calderone, Georgetown University Medical Center, USA
B Brett Finlay, University of British Columbia, Canada

Editorial Board

Andrew PW, University of Leicester, UK
Anné J, KU Leuven, Belgium
Armstrong GD, University of Calgary Health Sciences Centre, Canada
Blomfield I, University of Kent, UK
Blondeau JM, University of Saskatchewan, Canada
Bonomo RA, Case Western Reserve Univ., USA
Casadevall A, Albert Einstein Coll. Med., USA
Cauda R, Catholic University of Rome, Italy
El-Sayed NM, University of Maryland, USA
Evans TJ, University of Glasgow, UK
Fairlamb AH, University of Dundee, UK
Galabov AS, The Stephan Angeloff Institute of Microbiology, Bulgaria
Garau J, University of Barcelona, Spain
Goldberg JB, Emory Univ. School of Medicine, USA
Hagan P, Scottish Funding Council, UK
Høiby N, University of Copenhagen, Denmark
Hong SP, Gene Matrix Inc., Korea
Hsueh PR, National Taiwan Univ. Hosp., Taiwan
Karch H, Univ. Münster, Germany
Kaufmann SH, Max-Planck-Inst. Infect. Biol., Germany

Kaye KM, Brigham and Women's Hospital, USA
Maloy S, San Diego State University, CA, USA
Mitsuya H, National Cancer Institute, MD, USA
Nordmann P, University of Fribourg, Switzerland
Pennington TH, University of Aberdeen, UK
Pfaller MA, Univ. of Iowa Carver Coll. of Medicine, USA
Read R, University of Sheffield, UK
Roberts IS, University of Manchester, UK
Rossolini GM, University of Siena, Italy
Rubens CE, University of Washington, WA, USA
Salmond GPC, University of Cambridge, UK
Shapiro DS, Univ. of Nevada School of Medicine, USA
Singh S, CCMB, India
Stephens DS, Emory University Hospital, GA, USA
Struelens MJ, Univ. Libre de Bruxelles, Belgium
Swaminathan S, Univ. of Utah School of Medicine, USA
Tsakris A, University of Athens, Greece
Vila Estapé J, University of Barcelona, Spain
Waters AP, University of Glasgow, UK
Yen-Lieberman B, Cleveland Clinic, OH, USA
Zhou D, Beijing Institute of Microbiology and Epidemiology, China
Zinner SH, Harvard Medical School, MA, USA

Editorial Staff

Chairman
James Drake
Managing Director
Phil Garner

Commissioning Department

Editorial Director
Laura Dormer
Managing Commissioning Editor
Adam Price-Evans
Commissioning Editors
Alice Greenway
Rhiannon Finnie
Drug Evaluations Editor
Johanna Todd

Production Department

Head of Production
Kathryn Berry
Deputy Head of Production
Zara Robinson
Production Editors
Sharon Salt
Kelvin Adu-Darko
Editorial Assistant
Francis Davies

Enquiries

Advertising

Dionne Murray, Business Development Manager
d.murray@future-science-group.com

Editorial

Laura Dormer, Editorial Director
l.dormer@futuremedicine.com

Subscriptions

Dominik March, Subscription Sales Manager
d.march@futuremedicine.com

Reprints

Sam Cavana, Reprint Sales Manager
s.cavana@futuremedicine.com

Permissions

Adriana Gonzalez, Accounts Administrator
a.gonzalez@future-science-group.com

Disclaimer: Whilst every effort is made by the Publisher and Editorial Board to ensure that no inaccurate or misleading data, opinions or statements appear in this journal, they wish to make it clear that the data and opinions appearing herein are the responsibility of the contributor concerned. Accordingly, the Publisher, Editorial Board and their respective employees, officers and agents accept no liability whatsoever for the consequences of any inaccurate or misleading data, opinions or statements.

Copyright: Conditions of sale: *Future Microbiology* may be circulated only to those members of staff who are employed at the site at which the subscription is taken out. Readers are reminded that, under internationally agreed copyright legislation, photocopying of copyright materials is prohibited other than on a limited basis for personal use. Thus making copies of any article published in *Future Microbiology* is a breach of the law and can be prosecuted.

Indexing

Biobase, BIOSIS Previews, BIOSIS Reviews Reports and Meetings, Biotechnology Citation Index®, CAB abstracts, Chemical Abstracts, Current Awareness in Biological Sciences (CABS), EMBASE/Excerpta Medica, EMBiology, Journal Citation Reports/Science Edition, MEDLINE/Index Medicus, Science Citation Index Expanded (SciSearch®), Scopus®

Impact factor: 3.374 (2016)

Future Microbiology has inevitably been impacted by the genomic revolution. Research into the extreme diversity of microbial organisms will lead to the elucidation of new biologic pathways and gene products, and thus potential therapeutic strategies to combat or prevent infection. The availability of both human and microbial genome sequences will enable scientists to better understand exactly how microbes interact with the hosts they infect, and how an individual's genetic make-up influences their susceptibility to pathogens.

Accessible 'at-a-glance' formats are important in an increasingly time-constrained clinical community. The article structure has been optimized to make information immediately understandable and accessible.

Future Microbiology delivers essential information. Key advances in the field are reported and analyzed by international experts, providing an authoritative, but accessible, forum for this increasingly important and vast area of research.

Subscription options

Institutional subscriptions

Future Microbiology is available in print, electronic or print and electronic formats, and pricing will depend on your organization type (academic, corporate, hospital, etc). Please contact sales@futuremedicine.com for more details.

Global e-access licenses are available on request and attract considerable discounts from standard site license fees. For further details on global access licenses, please contact: sales@futuremedicine.com

Consortia pricing

Future Microbiology welcomes discussion with all consortia, and offers flexible packages and discounted prices. If you have specific questions or would like a quote please contact sales@futuremedicine.com for more details.

Personal subscriptions

Personal subscriptions are currently available to all *Future Medicine* journals. Payment must be made from a personal credit card registered to a home address. Print subscriptions will only be sent to a personal address. Please contact sales@futuremedicine.com for our personal order form.

Reprints

Article reprints are available through our reprint service. Please contact Sam Cavana: s.cavana@futuremedicine.com

Coverage includes:

- Microbe–host interactions
- Overviews highlighting optimal therapeutic and diagnostic approaches, along with potential future options
- The use of microbes in the treatment of disease and genetic engineering
- Summaries evaluating newly approved antimicrobial agents
- Pharmacoeconomics and cost–benefit issues in microbiology
- New and re-emerging microbes impacting human health
- Microbial genome research, and the implications to higher organisms
- Microbes and cancer
- Epidemiologic studies and trends
- The problem of drug resistance, and potential methods to overcome this
- Defense against the use of microbes in bioterrorism
- Real-world evidence and outcomes research

Ordering information

Please contact your local sales representative to place an order:

Worldwide

Future Science Ltd
Unitec House, 2 Albert Place,
London, N3 1QB, UK
Tel.: +44 (0)20 8371 6090
Fax: +44 (0)20 8343 2313
subscriptions@future-science.com

North America

sales.us@future-science.com

Latin America and the Caribbean

dotLib
Tel.: +55 (21) 3431 3430
info@dotlib.com

China

Charlesworth China
Tel.: +86 106 779 1601
sales@charlesworth.com.cn

Japan

USACO Corporation
Tel.: +81 3 3505 3257
Fax: +81 3 3505 6283
marketing@usaco.co.jp

Republic of Korea

Accucoms, Inc. (J Brighton)
Tel.: +82 10 3296 3611
seonghwi@accucoms.com

Hong Kong, Taiwan and Macau

FlySheet Information Services Co. Ltd.
Tel.: +886 2 26581258
emedical@flysheet.com.tw

Ordering information (cont.)

Far East (excluding China, Japan & Korea)

iGroup Asia Pacific
Tel.: +852 2572 7228
Fax: +852 2575 8822
info.hk@igroupnet.com

India

Moidore Publishing Group
Tel.: +91 8826008169
info@moidorepubgrp.com

JOURNAL POLICIES

Future Medicine titles endorse the Recommendations for the Conduct, Reporting, Editing and Publication of Scholarly Work in Medical Journals, issued by the International Committee for Medical Journal Editors, and the Code of Conduct and Best Practice Guidelines for Journal Editors, produced by the Committee on Publication Ethics, and GPP3. This information is also available at www.futuremedicine.com

Manuscript submission & processing

Future Medicine titles publish a range of article types, including solicited and unsolicited reviews, perspectives and original research articles. Receipt of all manuscripts will be acknowledged within 1 week and authors will be notified as to whether the article is to progress to external review. Initial screening of articles by internal editorial staff will assess the topicality and importance of the subject, the clarity of presentation, and relevance to the audience of the journal in question. If you are interested in submitting an article, or have any queries regarding article submission, please contact the Editorial Director (contact information can be found on our website at: www.futuremedicine.com). For new article proposals, the Editorial Director will require a brief article outline and working title in the first instance. We also have an active commissioning program whereby the Commissioning Editor, under the advice of the Editorial Board, solicits articles directly for publication.

External peer review: Through a rigorous peer-review process, Future Medicine titles aim to ensure that reviews are unbiased, scientifically accurate and clinically relevant. All articles are peer reviewed by three or more members of the International Editorial Board or other specialists selected on the basis of experience and expertise. Review is performed on a double-blind basis – the identities of peer reviewers and authors are kept confidential. Peer reviewers must disclose potential conflicts of interests that may affect their ability to provide an unbiased appraisal (see Conflict of Interest Policy below). Peer reviewers provide general comments to the editor and both general and specific comments to the author(s).

Where an author believes that an editor has made an error in declining a paper, they may submit an appeal. The appeal letter should clearly state the reasons why the author(s) considers the decision to be incorrect and provide detailed, specific responses to any comments relating to the rejection of the review. Further advice from members of the journal's Editorial Board and/or other external experts will be sought regarding eligibility for re-review.

Revision: Most manuscripts require some degree of revision prior to acceptance. Authors should provide two copies of the revised manuscript – one of which should be highlighted to show where changes have been made. Detailed responses to reviewers' comments, in a covering letter/email, are also required. Manuscripts may be accepted at this point or may be subject to further peer review. The final decision on acceptability for publication lies with the journal editor.

Post-acceptance

Accepted manuscripts will undergo production in-house. This will involve type-setting, copy-editing, proof-reading and re-drawing of any graphics. Authors will receive proofs of their article for approval and sign off and will be asked to sign a transfer of copyright agreement, except in circumstances where the author is ineligible to do so (e.g., government employees in some countries).

Embargo policy

Following the acceptance of articles for publication, authors (and their institutions, etc.) are welcome to publicize the publication; authors wishing to do so, should advise the editor of the details beforehand. No publicity relating to publication in a Future Medicine journal should be carried out while the manuscript is under consideration. However, prior publicity linked to presentations at meetings does not jeopardize publication in a Future Medicine journal. In cases where data may be of overwhelming public health importance, the above policy may be waived; should this be the case, the appropriate authorities responsible for public health should decide whether to disseminate information to physicians and the media in advance and should be responsible for this decision. The journal editor should be informed if these circumstances apply. Any queries relating to publicity of manuscripts should be directed to the journal editor.

Disclosure & conflict of interest policy

Authors must state explicitly whether financial and/or nonfinancial relationships exist that potentially conflict with the subject matter or materials discussed in the manuscript and any such potential conflict of interest (including sources of funding) should be summarized in a separate section of the published article. Authors must disclose whether they have received writing assistance and identify the sources of funding for such assistance. Authors declaring no conflict of interest are required to publish a statement to that effect within the article.

Authors must certify that they have disclosed relationships in which they (or a close family member): is employed, is a contractor, provides services, or has otherwise collaborated in commercial or scientific pursuits – even in the absence of direct monetary remuneration. Stock holdings and issued or pending patents of an author or family member should also be disclosed. This list is not exclusive of other forms of financial involvement. A 36-month disclosure window should be used. Details of relevant conflicts of interests (or the lack of) must be declared in the 'Disclosure' section of the manuscript for all listed authors.

External peer reviewers must disclose any conflicts of interest that could bias their opinions of the manuscript, and they should disqualify themselves from reviewing specific manuscripts if they believe it appropriate. Should any such conflict of interest be declared, the journal editor will judge whether the reviewer's comments should be recognized or will interpret the reviewer's comments in the context of any such declaration.

Ethical conduct of research

For studies involving data relating to human or animal experimental investigations, appropriate institutional review board approval is required and should be described within the article. For those investigators who do not have formal ethics review committees, the principles outlined in the Declaration of Helsinki should be followed. For investigations involving human subjects, authors should explain how informed consent was obtained from the participants involved.

Patients' rights to privacy

Patients have a right to privacy that should not be infringed without informed consent. Identifying information should not be included unless the information is essential for scientific purposes and the patient (or parent or legal guardian) gives written informed consent for publication. Informed consent for this purpose requires that the patient be shown the manuscript to be published. When informed consent has been obtained it should be indicated in the manuscript.

In attempting to maintain patient anonymity, identifying details should be omitted where they are not essential. However, patient data should never be amended or falsified. Informed consent should be obtained whenever there is any doubt that anonymity can be assured.

Use of personal communications & unpublished data

Where an individual is identified within a review as a source of information in a personal communication or as a source for unpublished data, authors should include a signed statement of permission from the individual(s) concerned and specify the date of communication.

Clinical trial registration

Future Medicine titles prefer to publish clinical trials that have been included in a clinical trials registry that is accessible to the public at no charge, is electronically searchable, is open to prospective registrants and is managed by a not-for-profit organization, such as www.clinicaltrials.gov (sponsored by the United States National Library of Medicine). Where a clinical trial registration number is available, this should be included at the end of the abstract and also listed the first time the authors use a trial acronym to refer to the trial they are reporting in the manuscript. Unregistered clinical trials should be declared as such, and the reason for nonregistration should be provided. Whilst referees will take registration status into account, all well designed and presented trials and corresponding data will be considered for publication.

Errata/corrigenda

Mistakes by either editor or author should be identified wherever possible and an erratum or corrigendum published at the earliest opportunity. We will attempt to contact the author of the original article to confirm any error, and publish an appropriate erratum or corrigendum at the earliest opportunity.

Duplicate publication/submission & plagiarism

All manuscripts submitted to Future Medicine titles are considered for publication on the understanding that they have not been published previously elsewhere or are under consideration for publication elsewhere. The journal may, however, consider republication of a paper previously published in a language other than English, subject to prominent disclosure of the original source and with any necessary permission. Authors will be asked to certify that the manuscript represents valid work and that neither this manuscript nor one with substantially similar content under their authorship has been published or is being considered for publication elsewhere. Where specific findings from a particular study have been previously published (in Future Medicine titles or elsewhere), Future Medicine titles will not consider manuscripts reporting the same findings, except where: the results are substantially reanalyzed, reinterpreted for a different audience, or translated into another language; the primary publication is clearly acknowledged and cited and the trial registration number (where available) of the original research is included; and the publication is clearly presented as an analysis derived from the primary publication results or marked as a translation, with appropriate permission obtained from the previous publisher and copyright laws upheld. All submitted articles will be evaluated using plagiarism detection software, which compares the submitted manuscript with full text articles from all major journals databases and the internet. The use of published or unpublished ideas, words or other intellectual property derived from other sources without attribution or permission, and representation of such as those of the author(s) is regarded as scientific misconduct and will be addressed as such.

Scientific misconduct & retraction

If misconduct by authors or reviewers is suspected, either pre- or post-publication, action will be taken. An explanation will be sought from the party or parties considered to be involved. If the response is unsatisfactory, then an appropriate authority will be asked to investigate fully. Future Medicine will make all reasonable attempts to obtain a resolution in any such eventuality and correct the record or archive as necessary (publishing a retraction of the article as required).

Compliance with funder open or public access policies

Future Medicine is supportive of open and public access policies mandated by various funding bodies. In the first instance, please check whether your article type is covered by your funder mandate (i.e., in some cases, funder policies only cover research articles, rather than review articles). Please advise the Editor on submission if your article was funded by any of the bodies below. Where open access publication is a requirement of your funding, our article processing charge will need to be paid (in many cases, this will be covered by your funder, and you should check the specific details of this on the funder website). As standard for our Open Access option, articles are published under a CC BY-NC-ND license, and this will be used in the first instance; however, where the funder requires it, articles can be published under a different Creative Commons license (i.e., the CC BY license in the case of The Wellcome Trust).

Self-archive policy

Future Medicine is supportive of self-archiving. Should you be interested in archiving your work, please be aware of the policies outlined on our website at www.futuremedicine.com/page/authors.jsp.

Insights into the evolution of the mutational resistome of *Pseudomonas aeruginosa* in cystic fibrosis

Carla López-Causapé¹ & Antonio Oliver^{*,1}

¹Servicio de Microbiología & Unidad de Investigación, Hospital Universitario Son Espases, Instituto de Investigación Illes Balears (IdISBa), Palma de Mallorca, Spain

* Author for correspondence: antonio.oliver@ssib.es

“The analysis of WGS mutational resistomes has proven to be useful for understanding the evolutionary dynamics of classical resistance mechanisms and to depict new ones for the majority of antimicrobial classes”

First draft submitted: 6 September 2017; Accepted for publication: 12 September 2017; Published online: 25 October 2017

Keywords: antibiotic resistance • bacterial evolution • chronic infections • cystic fibrosis • hypermutation • *Pseudomonas aeruginosa* • resistome

Chronic respiratory infection (CRI) by *Pseudomonas aeruginosa* is the main driver of morbidity and mortality in cystic fibrosis (CF) patients [1]. CRI results from an intense adaptation process, where bacterial evolution is tested against host immune responses and years of aggressive antimicrobial treatments [2]. Once established, CRI can seldom be eradicated despite intensive antimicrobial treatments, and therefore our therapeutic goals resignedly move from attempting to cure the infection to minimizing its long-term impact through chronic suppressive therapy [3]. The plasticity of *P. aeruginosa* genome for antimicrobial resistance acquisition, the greatly enhanced mutation supply rate provided by frequent hypermutable variants (mutators) and the highly structured environment determined by the characteristic biofilm growth and the anatomy of the respiratory tract make bacterial evolution and genetic diversification a hallmark of CF CRI [2,4]. While the enhanced evolution of antimicrobial resistance in CF, frequently linked to mutator phenotypes, was noted many years ago [5], it is with the introduction of whole-genome sequencing (WGS) that we are starting to understand its real dimensions [6].

The term resistome was first used to account for the set of primary antibiotic resistance genes that could be eventually transferred to human pathogens [7]. Soon after the concept of intrinsic resistome was introduced to include all chromosomal genes that are involved in intrinsic resistance, and whose presence in strains of a bacterial species is independent of previous antibiotic exposure and is not due to horizontal gene transfer (HGT) [8]. Finally, the term mutational resistome was more recently implemented to account for the set of mutations involved in the modulation of antibiotic resistance levels in the absence of HGT [9]. Recent WGS data obtained from *in vitro* assays on the evolution of antibiotic resistance and clinical isolates, and in particular sequential CF isolates, provide new insights into the evolutionary dynamics and mechanisms of *P. aeruginosa* antibiotic resistance. However, in too many cases, the documented genomic variations fail to provide causative relations in the absence of phenotypic information. The analysis of WGS mutational resistomes has proven to be useful for understanding the evolutionary dynamics of classical resistance mechanisms and to depict new ones for the majority of antimicrobial classes, including β -lactams, aminoglycosides, fluoroquinolones and polymyxins.

Regarding β -lactams, the analysis of WGS mutational resistomes has confirmed the major role of classical resistance mutations such as those leading to the overexpression of the chromosomal β -lactamase AmpC (such as DacB [PBP4], AmpD and/or AmpR mutations) or the inactivation of the carbapenem porin OprD. However, the analysis of mutational resistomes of *in vitro* evolved strains and sequential CF isolates have identified other key mutations, such as those occurring in β -lactam targets (essential PBPs), particularly involving mutations in *fisI* which encodes PBP3, an essential high molecular class B penicillin binding protein (PBP) with transpeptidase

activity. Indeed, data from CF patients [9,10] as well as from *in vitro* studies [11] have recently demonstrated that PBP3 is under strong mutational pressure, with specific mutations contributing to β -lactam resistance development. Among them are particularly relevant and frequent mutations affecting amino acids R504 or F533, located within the protein domains responsible for the formation and stabilization of the inactivating complex β -lactam–PBP3. Moreover, PBP3 mutations seem to play a role in the emergence of resistance to novel β -lactam– β -lactamase inhibitor combinations, such as ceftolozane/tazobactam [9]. Another relevant mutational β -lactam resistance mechanism is the selection of large (>200 kb) deletions affecting specific parts of the chromosome. Although the basis of the conferred resistance phenotype still needs to be further clarified, these mutants can be recognized by the characteristic brown pigment (pyomelanine) caused by the deletion of one of the affected genes, *hmgA*, coding for a homogentisate-1,2-dioxygenase. This type of deletion has been documented in both, *in vitro* evolved β -lactam-resistant mutants and CF isolates [11,12]. However, the deletion of *hmgA* is not responsible for the resistance phenotype, which may be linked to the deletion of another of the affected genes, *galU*, coding for a UDP-glucose pyrophosphorylase required for lipopolysaccharide core synthesis. Indeed, analysis of transposon mutant libraries has shown that inactivation of *galU* increases ceftazidime and meropenem minimum inhibitory concentrations [13,14]. Finally, another emerging mutational β -lactam resistance mechanism is the structural modification of AmpC [10,11].

With respect to aminoglycosides, results from analysis of mutational resistomes of CF isolates point to the underlying strong evolutionary pressure of *mexZ* and the relevance of *MexXY* overexpression for resistance development [15–17]. Moreover, recent *in vitro* studies and findings from CF isolates have revealed that high-level aminoglycoside resistance requires the acquisition of additional mutations; among them, those in *FusA1* seem to be particularly frequent and relevant [9,16,18–19]. Likewise, the fluoroquinolone resistome frequently includes mutations in efflux pump regulators, among which *nfxB*, leading to the overexpression of *MexCD-OprJ*, is particularly noteworthy in the CF setting. However, high-level ciprofloxacin resistance generally involves one or several mutations in the quinolone resistance determining regions of *GyrA/B* and/or *ParC/E* [9]. Regarding polymixin resistance, findings from WGS studies of *in vitro* evolved strains and CF isolates have shown that development of high-level colistin resistance requires the acquisition of multiple mutations, including those in the two-component regulators (*PmrAB*, *PhoPQ* or *ParRS*) involved in the addition of 4-amino-4-deoxy-L-arabinose to lipid A from the lipopolysaccharide [9,20]. Finally, in addition to the resistance mechanisms to classical antipseudomonal agents, the CF mutational resistome may also include resistance to other used antibiotics such as the frequent mutations of domain V of 23S rRNA – conferring macrolide resistance [21].

The complexity of the CF isolates resistomes is further enhanced when the inpatient genetic diversity of CF *P. aeruginosa* populations is introduced. Certainly, to understand the resistance dynamics and evolution, future steps should endeavor to analyze the mutational resistomes in CF at the whole population level, as opposed to the analysis of single isolated colonies. Indeed, full understanding of the evolution of the mutational resistome requires a longitudinal and transversal analysis of *P. aeruginosa* populations in the CF patient. Moreover, recent evidence suggests that interpatient transmission should also be considered when analyzing the evolution of the mutational resistome, especially when introducing mutator lineages of epidemic clones [9].

Beyond addressing a relevant scientific question, the analysis of mutational resistomes would be useful for therapeutic strategy design and monitoring the efficacy of administered antibiotic treatments. Obviously, the evolution of the mutational resistome is a direct consequence of antimicrobial exposure. As such, it is not surprising that exposure to one antibiotic drives evolution of the mutational resistome for that antibiotic. However, the complexity of the actual resistance profile is further increased by the specificity and interactions among different resistance mechanisms. A classic example is cross resistance (or collateral resistance), which implies that exposure to one antibiotic drives also the development of resistance to a different one. Typically this is caused by the developed resistance mechanism (such as efflux pump overexpression) affecting simultaneous different antibiotics. Indeed, potential development of cross resistance is a major issue to consider when using antibiotic combinations [22].

Perhaps less obvious is collateral susceptibility, which implies that exposure to one antibiotic increases the susceptibility to a different one. This might be achieved through two mechanisms. One possible mechanism is that exposure to one antibiotic directly causes increased susceptibility to a different one, for example, mutations in the β -lactamase AmpC increases cephalosporin hydrolysis while reducing that of penicillins or carbapenems [23]. The second possibility is that the development of a resistance mechanism impairs the activity of another existing resistance mechanism, for example, competition between the different efflux pumps, the overexpression of one may impair the expression of another [24]. Indeed, in the CF setting it is very frequent that the overexpression of efflux pump *MexXY*, involved in aminoglycoside resistance, is linked to the impaired expression of efflux pump *MexAB*,

involved in the resistance to a broad range of antibiotics including most β -lactams [25]. Thus, the evolution of the mutational resistome for a given antibiotic is not only dependent on the exposure to this antibiotic, but it is also conditioned by the simultaneous or even previous exposures to other antibiotics. An illustrative example is provided in a recent *in vitro* study that demonstrated, for a broad range of antibiotic classes, that the history of exposure and resistance development to a given antibiotic, conditions the dynamics and mechanisms of resistance development when exposed to a second one [18].

Moreover, knowledge of the interactions between resistance mechanisms could be useful in the design of sequential treatments that minimize the risk of resistance development. Such is the case for a recent *in vitro* study showing the effectiveness of aztreonam–tobramycin sequential treatment, based on the antagonism between the resistance mechanisms for each of the antibiotics; overexpression of the efflux pump MexAB (aztreonam) or MexXY (tobramycin) which compete for the same outer membrane channel (OprM) [26].

Finally, it should be noted that a hallmark of *P. aeruginosa* CRI in CF is the biofilm mode of growth. Indeed, biofilm growth, in addition to providing a compact structured environment likely facilitating the evolution of mutational resistance [27], they also add further complexity due to the major differences in the mechanisms involved when compared with conventional planktonic growth that needs to be considered [28]. Likewise, persistence of *P. aeruginosa* in the CF lung despite intensive antimicrobial treatments relays in the acquisition of a vast number of adaptive mutations that extend far beyond classical antibiotic resistance mutations [29].

In summary, the comprehensive analysis of the mutational resistomes of *P. aeruginosa* in CF CRI is expected to become a useful tool for optimizing therapeutic strategies and monitoring the efficacy of administered antibiotic treatments in the near future.

Financial & competing interests disclosure

The authors are supported by the Ministerio de Economía y Competitividad of Spain, Instituto de Salud Carlos III – co-financed by European Regional Development Fund ‘A way to achieve Europe’ ERDF, through the Spanish Network for the Research in Infectious Diseases (RD12/0015 and RD16/0016). The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

References

1. Lyczak JB, Cannon CL, Pier GB. Lung infections associated with cystic fibrosis. *Clin. Microbiol. Rev.* 15(2), 194–222 (2002).
2. Oliver A, Mena A, Macià MD. Evolution of *Pseudomonas aeruginosa* pathogenicity: from acute to chronic infections. In: *Evolutionary Biology of Bacterial and Fungal Pathogens*. Baquero F, Nombela C, Cassell GH, Gutiérrez JA (Eds). ASM Press, WA, USA, 433–444 (2008).
3. López-Causapé C, Rojo-Molinero E, Macià MD, Oliver A. The problems of antibiotic resistance in cystic fibrosis and solutions. *Expert. Rev. Respir. Med.* 9(1), 73–88 (2015).
4. Fernández-Barat L, Ciofu O, Kragh KN *et al.* Phenotypic shift in *Pseudomonas aeruginosa* populations from cystic fibrosis lungs after 2-week antipseudomonal treatment. *J. Cyst. Fibros.* 16(2), 222–229 (2017).
5. Oliver A, Cantón R, Campo P, Baquero F, Blázquez J. High frequency of hypermutable *Pseudomonas aeruginosa* in cystic fibrosis lung infection. *Science* 288(5469), 1251–1254 (2000).
6. Feliziani S, Marvig RL, Luján AM *et al.* Coexistence and within-host evolution of diversified lineages of hypermutable *Pseudomonas aeruginosa* in long-term cystic fibrosis infections. *PLoS Genet.* 10(10), e1004651 (2014).
7. D’Costa VM, McGrann KM, Hughes DW, Wright GD. Sampling the antibiotic resistome. *Science* 311(5759), 374–377 (2006).
8. Fajardo A, Martínez-Martín N, Mercadillo M *et al.* The neglected intrinsic resistome of bacterial pathogens. *PLoS ONE* 3(2), e1619 (2008).
9. López-Causapé C, Sommer LM, Cabot G *et al.* Evolution of the *Pseudomonas aeruginosa* mutational resistome in an international cystic fibrosis clone. *Sci. Rep.* 7(1), 5555 (2017).
10. Diaz Caballero J, Clark ST, Coburn B *et al.* Selective sweeps and parallel pathoadaptation drive *Pseudomonas aeruginosa* evolution in the cystic fibrosis lung. *MBio.* 6(5), e00981–15 (2015).
11. Cabot G, Zamorano L, Moyà B *et al.* Evolution of *Pseudomonas aeruginosa* antimicrobial resistance and fitness under low and high mutation rates. *Antimicrob. Agents Chemother.* 60(3), 1767–1778 (2016).
12. Hocquet D, Petitjean M, Rohmer L *et al.* Pyomelanin-producing *Pseudomonas aeruginosa* selected during chronic infections have a large chromosomal deletion which confers resistance to pyocins. *Environ. Microbiol.* 18(10), 3482–3493 (2016).

13. Alvarez-Ortega C, Wiegand I, Olivares J, Hancock RE, Martínez JL. Genetic determinants involved in the susceptibility of *Pseudomonas aeruginosa* to beta-lactam antibiotics. *Antimicrob. Agents Chemother.* 54(10), 4159–4167 (2010).
14. Dötsch A, Becker T, Pommerenke C *et al.* Genomewide identification of genetic determinants of antimicrobial drug resistance in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 53(6), 2522–2531 (2009).
15. Prickett MH, Hauser AR, McColley SA *et al.* Aminoglycoside resistance of *Pseudomonas aeruginosa* in cystic fibrosis results from convergent evolution in the *mexZ* gene. *Thorax* 72(1), 40–47 (2017).
16. Greipel L, Fischer S, Klockgether J *et al.* Molecular epidemiology of mutations in antimicrobial resistance loci of *Pseudomonas aeruginosa* isolates from airways of cystic fibrosis patients. *Antimicrob. Agents Chemother.* 60(11), 6726–6734 (2016).
17. Marvig RL, Sommer LM, Molin S, Johansen HK. Convergent evolution and adaptation of *Pseudomonas aeruginosa* within patients with cystic fibrosis. *Nat. Genet.* 47(1), 57–64 (2015).
18. Yen P, Papin JA. History of antibiotic adaptation influences microbial evolutionary dynamics during subsequent treatment. *PLoS Biol.* 15(8), e2001586 (2017).
19. Feng Y, Jonker MJ, Moustakas I, Brul S, Ter Kuile BH. Dynamics of mutations during development of resistance by *Pseudomonas aeruginosa* against five antibiotics. *Antimicrob. Agents Chemother.* 60(7), 4229–4236 (2016).
20. Jochumsen N, Marvig RL, Damkiær S *et al.* The evolution of antimicrobial peptide resistance in *Pseudomonas aeruginosa* is shaped by strong epistatic interactions. *Nat. Commun.* 7, 13002 (2016).
21. Mustafa MH, Khandekar S, Tunney MM *et al.* Acquired resistance to macrolides in *Pseudomonas aeruginosa* from cystic fibrosis patients. *Eur. Respir. J.* 49(5), pii:1601847 (2017) (Epub ahead of print).
22. Vestergaard M, Paulander W, Marvig RL *et al.* Antibiotic combination therapy can select for broad-spectrum multidrug resistance in *Pseudomonas aeruginosa*. *Int. J. Antimicrob. Agents.* 47(1), 48–55 (2016).
23. Cabot G, Bruchmann S, Mulet X *et al.* *Pseudomonas aeruginosa* ceftolozane-tazobactam resistance development requires multiple mutations leading to overexpression and structural modification of AmpC. *Antimicrob. Agents Chemother.* 58(6), 3091–3099 (2014).
24. Mulet X, Moyá B, Juan C *et al.* Antagonic interactions of *Pseudomonas aeruginosa* antibiotic resistance mechanisms in planktonic but no biofilm growth. *Antimicrob. Agents Chemother.* 55(10), 4560–4568 (2011).
25. Vettoretti L, Plésiat P, Muller C *et al.* Efflux unbalance in *Pseudomonas aeruginosa* isolates from cystic fibrosis patients. *Antimicrob. Agents Chemother.* 53(5), 1987–1997 (2009).
26. Rojo-Molinero E, Macià MD, Rubio R *et al.* Sequential treatment of biofilms with aztreonam and tobramycin is a novel strategy for combating *Pseudomonas aeruginosa* chronic respiratory infections. *Antimicrob. Agents Chemother.* 60(5), 2912–2922 (2016).
27. Macià MD, Pérez JL, Molin S, Oliver A. Dynamics of mutator and antibiotic-resistant populations in a pharmacokinetic/pharmacodynamic model of *Pseudomonas aeruginosa* biofilm treatment. *Antimicrob. Agents Chemother.* 55(11), 5230–5237 (2011).
28. Ciofu O, Rojo-Molinero E, Macià MD, Oliver A. Antibiotic treatment of biofilm infections. *APMIS.* 125(4), 304–319 (2017).
29. Smith EE, Buckley DG, Wu Z *et al.* Genetic adaptation by *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients. *Proc. Natl Acad. Sci. USA* 103(22), 8487–8492 (2006).

Protective efficacy of antidiarrheal agents in a permeability model of *Escherichia coli*-infected CacoGoblet® cells

Barbara de Servi*,¹ & Francesco Ranzini¹

¹VitroScreen Srl, Via Mosè Bianchi 103, 20149 Milano, Italy

* Author for correspondence: Tel.: +39 02 8907 7608; barbara.deservi@vitroscreen.com

Aim: To compare the protective efficacy of gelatine tannate/probiotic with other antidiarrheal agents in *Escherichia coli*-inoculated CacoGoblet® cells. **Methods:** Four test compounds – gelatine tannate plus inactivated probiotic, diosmectite, probiotic mixture and *Saccharomyces boulardii* – were added to *E. coli*-infected CacoGoblet cells. After 1 and 24 h, transepithelial electrical resistance was measured and a lucifer yellow assay performed. **Results:** Gelatine tannate/probiotic markedly increased transepithelial electrical resistance by 123.1% (at 1 h) and 149.5% (at 24 h), and produced paracellular flux values of 0.41% (1 h) and 1.34% (24 h), which were considerably less than the *E. coli*-invasion value (2.41%). **Conclusion:** The protective efficacy of gelatine tannate/probiotic against *E. coli*-induced reduction of membrane integrity manifests early and is maintained for 24 h.

First draft submitted: 17 October 2016; Accepted for publication: 27 July 2017; Published online: 25 October 2017

Keywords: acute gastroenteritis • diarrhea • *Escherichia coli* • gelatin tannate • inactivated probiotics • mucoprotectors

Worldwide, diarrhea is a major cause of morbidity and mortality. Despite marked enhancements in health technology, medical management and the use of oral rehydration solutions in recent decades, it remains a significant cause of death in children ≤5 years of age [1,2]. Pathogenic microorganisms can adhere to the intestinal mucosa and influence cellular tight-junction function, leading to diarrhea. Such organisms also penetrate the mucosa, becoming internalized and proliferating inside intestinal cells, thereby worsening diarrhea [3]. Rotavirus, for instance, is the principal cause of acute gastroenteritis in infants and children, and is associated with a major clinical and economic burden [4].

In vitro, Caco-2 cells provide a well-established reference model of intestinal mucosal functionality [5–8]. The model is widely used for the investigation of drug effects on human intestinal epithelial function, and is acknowledged by the US FDA as an appropriate test medium for *in vitro* toxicology studies [9].

Tight junctions, which are intercellular junctions composed of a sticky filament network, regulate polyphenol paracellular intestinal transport *in vivo* and the pathway can be followed *in vitro* in appropriate conditions in Caco-2 models. However, co-cultured Caco-2 cells and human mucosal goblet cells, as in the CacoGoblet® medium (ReadyCell SL, Barcelona, Spain), are now being used more widely than Caco-2 cells alone to provide an enhanced physiological model of human intestinal permeability for evaluating antidiarrheal agents [10,11]. Advantages of the CacoGoblet medium over Caco-2 cells are listed in Table 1.

Gelatine tannate is an antidiarrheal agent that forms a mechanical barrier protecting the intestinal mucosa in patients with acute gastroenteritis [12–15]. Additional evidence now suggests that gelatine tannate also has direct film-forming and antiadhesive properties in the intestinal mucosa, thus inhibiting growth of some bacterial species and preserving the intestinal mucous layer [12,15–16]. Health-promoting bacteria (probiotics), such as some lactobacilli, bifidobacteria and streptococci, have mild luminal effects in the intestine: antitoxic activity against specific pathogens; antimicrobial activity; preservation of tight-junction integrity; and modulation of the intestinal flora [17–20]. Heat-killed probiotics, such as dead cells of *Enterococcus faecalis* and bifidobacteria, have shown favorable anti-inflammatory effects, including enhanced production of TNF- α , increased neutrophil phagocytosis

Table 1. CacoGoblet® medium is a better physiological model of human intestinal functionality than Caco-2 cells.

	Human intestine	CacoGoblet medium	Caco-2 cells
Composition: absorptive cells; mucus-secreting cells, %	80%; 10–30%	50%; 50%	100%; 0%
Presence of mucus	Yes	Yes	No
Paracellular permeability	More permissive epithelium	More permissive epithelium	Very tight epithelium
TEER (Ω/cm^2)	20–110	2000–3000	80–120

TEER: Transepithelial electrical resistance.

and general stimulation of the gastrointestinal immune system [21]. Such heat-killed probiotics are considered to be at least as effective, and markedly safer, than live probiotics [22], and in the clinical setting no significant differences were identified between live and heat-killed probiotics regarding reduced duration and frequency of diarrhea [23]. It has therefore been suggested that combination of gelatine tannate with probiotics may lead to: a faster onset of probiotic action, after resolution of diarrhea by gelatine tannate; and synergistic activity to restrict the intestinal adhesion, internalization and proliferation of pathogenic bacteria.

Thus, the current study was designed to compare the protective efficacy of gelatine tannate plus inactivated probiotic (Tasectan Duo®; Novintethical, SA, Lugano, Switzerland) with that of various other antidiarrheal agents in an *in vitro* model of *Escherichia coli* invasion of mucus-secreting CacoGoblet medium.

Materials & methods

Study medium

CacoGoblet is a ready-to-use medium for *in vitro* evaluation of intestinal absorption. The kit provides a 21-day cell barrier formed by differentiated co-cultured Caco-2 and human goblet mucus secreting cells (HT29-H and HT29-MTX) plated on HTS Transwell® (Sigma-Aldrich, Milan, Italy) permeable supports. CacoGoblet was delivered to the laboratory on day 20 of differentiation (with an exclusive and proprietary shipping medium stable at room temperature). The complete medium was charged and transepithelial electrical resistance (TEER) measured by an internal procedure (see ‘TEER measurement’).

Escherichia coli inoculum

The *E. coli* (ATCCC 8739 wild-type with standard virulence) inoculum was prepared in a microbiological Good Laboratory Practice-certified laboratory. The inoculum comprised apical addition of 10 μl of stock culture (1×10^9 colony-forming units [cfu]/ml) on a CacoGoblet cell monolayer to obtain a final concentration of 1×10^8 cfu/ml.

Test compounds

All test compounds were dissolved in 1% sodium bicarbonate solution. Four test compounds were evaluated:

- Gelatine tannate plus inactivated probiotic (comprising 2.5% inactivated lactobacilli; Tasectan Duo) 5 mg/ml.
- Diosmectite (Smecta®; Ipsen, Merelbeke, Belgium) 5 mg/ml.
- Probiotic mixture (*Lactobacillus* spp., *Bifidobacterium bifidum*, *Streptococcus thermophilus*) 12.5 mg/ml.
- *Saccharomyces boulardii* 12.5 mg/ml.

Test controls

Negative test controls comprised untreated CacoGoblet medium alone, and the test medium in 1% sodium bicarbonate.

Test reagents

For the lucifer yellow paracellular passage assay (see ‘The lucifer yellow assay’), the principal reagents used were lucifer yellow (Sigma-Aldrich), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES; Lonza Group Ltd, Basel, Switzerland), 2-(N-morpholino)ethanesulfonic acid (MES; Sigma-Aldrich), and Hank’s balanced salt solution (HBSS; Sigma-Aldrich).

Table 2. Percentage transepithelial electrical resistance recovery at 1 and 24 h after incubation of test compounds with *Escherichia coli*-infected CacoGoblet®.

	TEER recovery (%)	
	1 h	24 h
Gelatine tannate/probiotic 5 mg/ml	123.1	149.5
Diosmectite 5 mg/ml	68.8	-4.5
Probiotic mixture 12.5 mg/ml	107.2	17.9
<i>Saccharomyces boulardii</i> 12.5 mg/ml	16.1	11.4

TEER: Transepithelial electrical resistance.

Study procedure

Transepithelial electrical resistance measurement

TEER is a direct measure of barrier function. It indicates the overall strength of a tissue based on structure and thickness, and measures barrier integrity at tight junctions between enterocytes. A standard value for TEER on CacoGoblet is 80–120 Ω/cm^2 , as measured by a volt-ohm-meter (Millicell® ERS, Merck, Vimodrone, Italy).

TEER measurement was performed according to the internal standard operating procedure, by using the appropriate electrode in accordance with supplier instructions. A single measurement of TEER was made before and after *E. coli* inoculation (10^8 cfu/ml) of CacoGoblet for 30 min, as this time has previously been shown to reversibly reduce TEER. A single TEER measurement was also made for untreated CacoGoblet. Each of the four test compounds, or 1% sodium bicarbonate (control), was applied (without washing) to *E. coli*-infected CacoGoblet for 1 and 24 h; a TEER measurement was made at each time-point for each test compound.

The lucifer yellow assay

Lucifer yellow is a fluorescent dye that does not readily penetrate cell membranes and, as such, is used to evaluate paracellular permeability. When intercellular tight junctions are unbroken, lucifer yellow has a very low permeability; however, if junctions are damaged, the flow of lucifer yellow is much greater.

The lucifer yellow paracellular passage assay was performed after each application of test compound or 1% sodium bicarbonate control to the *E. coli*-infected CacoGoblet medium: that is, at times 0, 1 and 24 h. The medium was rinsed with HBSS, and the basolateral cellular chamber was filled with 0.9 ml of 1% HBSS–HEPES; the apical chamber was filled with 0.3 ml of 100 μM lucifer yellow in 1% HBSS–MES. Transport of lucifer yellow was assessed as movement of the reagent into the basolateral chamber during a 30-min incubation period at 37°C. Apical fluorescence (RFU-AP) and basolateral fluorescence (RFU-BL) readings were obtained using a Tecan Infinite® M200 spectrofluorimeter (Tecan, Männedorf, Switzerland) with 428 nm excitation and 535 nm emission. Flux of lucifer yellow (%) was calculated from the following formula:

$$\frac{\text{RFU} - \text{BL}}{\text{RFU} - \text{AP}(t = 0)} \times 100$$

No specific statistical analyses were conducted, except for calculation of standard deviation.

Results

Transepithelial electrical resistance measurement

Before addition of each of the four test compounds, 30-min inoculation of CacoGoblet medium with *E. coli* led to a 41.0–42.4% decrease in TEER to 72.93 Ω/cm^2 (Figure 1). Gelatine tannate/probiotic markedly increased TEER by 123.1% (at 1 h) and 149.5% (at 24 h) above the 30-min *E. coli*-invasion value (Table 2). This indicated that gelatine tannate/probiotic produced complete recovery, to above the baseline value, of stress induced by *E. coli* inoculation of CacoGoblet cells.

Diosmectite increased TEER by 68.8% (at 1 h) above the *E. coli*-invasion value, but this effect was not maintained (-4.5% at 24 h), and no recovery to above the baseline value was noted. The probiotic mixture increased TEER to above the baseline and *E. coli*-invasion values (+107.2% at 1 h), but again, this effect was not maintained (+17.9%

Figure 1. Measurement of transepithelial electrical resistance (mean $\pm$ standard deviation) in *Escherichia coli*-infected CacoGoblet[®] medium.

Times shown are before (0 min) and after (30 min) *E. coli* inoculation of CacoGoblet cells, and after a subsequent 1 and 24 h of incubation with each of the four test compounds.

TEER: Transepithelial electrical resistance.

at 24 h). TEER values after incubation of *E. coli*-infected CacoGoblet cells with 12.5 mg/ml *S. boulardii* were similar to the baseline and *E. coli*-invasion values (+11.4–16.1%; Table 2).

Lucifer yellow paracellular flux

Flux measured in the *E. coli*-invaded medium, without test compounds, was 3.00% at 1 h and 2.41% at 24 h. The value for untreated medium was 0.51%. These values were less than the threshold of 10% considered to confirm maintenance of monolayer integrity, and appropriateness of the model for determining changes in paracellular flux in response to test compounds.

Gelatine tannate/probiotic produced a paracellular flux value of 0.41% at 1 h, which was similar to that for untreated medium (0.51%). The value at 24 h (1.34%) was also considerably less than the *E. coli*-invasion value (2.41%; Figure 2), thus suggesting major protective efficacy for gelatine tannate/probiotic in counteracting *E. coli* invasion. Diosmectite had only a marginal effect in countering the increase in paracellular flux produced by *E. coli* inoculation, as indicated by flux values of 1.9% (at 1 h) and 2.30% (at 24 h). The probiotic mixture had a marked initial effect (flux value 0.54% at 1 h), but this was not maintained, since the value at 24 h (6.44%) was considerably greater than the *E. coli*-invasion value (2.41%). The same was true for 12.5 mg/ml *S. boulardii*: 1.70% (1 h) and 4.16% (24 h).

Discussion

This study has clearly shown that in an *E. coli*-invasion model in CacoGoblet medium, the protective efficacy of gelatine tannate/probiotic against *E. coli*-induced reduction of membrane integrity manifests early and is maintained for 24 h. That is, marked increases in TEER to 138.9 and 153.1 Ω/cm^2 were noted after 1 and 24 h, respectively. The mean pretreatment TEER was 72.9 Ω/cm^2 , and importantly, values $<70 \Omega/\text{cm}^2$ correspond to severe modification of tissue permeability that allows paracellular flux of almost all molecules and bacteria. The lucifer yellow assay is more sensitive than TEER for assessing tight-junction integrity and function. In this study, gelatine tannate/probiotic produced lucifer yellow paracellular flux values at 1 and 24 h that were considerably less than the *E. coli*-invasion value, thus suggesting major protective efficacy for gelatine tannate/probiotic in counteracting *E. coli* invasion.

Tight junctions are markedly dynamic structures that undergo rapid reorganization after biological and physical stimuli. Depending on the severity of bacterial challenge, and highlighting validity of the *E. coli*-invasion model in

Figure 2. Results from the lucifer yellow paracellular flux assay.

Times shown are after *Escherichia coli* inoculation of CacoGoblet® cells, and after subsequent 1 and 24 h of incubation with each of the four test compounds. Values shown are mean $\pm$ standard deviation.

LY: Lucifer yellow.

the present study, recovery of tight-junction function and tissue integrity [24] can be used to classify the protective efficacy of various test compounds. The observed recovery of barrier integrity and tight-junction fence properties in response to gelatine tannate/probiotic in the current trial may have been due to positive interaction of the combination with mucus at the epithelial surface in the CacoGoblet medium.

Regarding other compounds evaluated in this study, diosmectite moderately restored barrier integrity and fence properties in the CacoGoblet medium after 1 h, as indicated by a TEER increase and slight decrease in lucifer yellow paracellular flux; however, these protective effects were no longer evident at 24 h. The same was true for the probiotic mixture tested. *S. boulardii* had no effect on paracellular flux, as indicated by TEER measurement, although a marginal effect was noted at 1 h in the lucifer yellow assay, but not at 24 h in the latter assay.

In the clinical setting, the rationale for combination of gelatine tannate with inactivated probiotic is the potential for synergy. That is, both gelatine tannate and probiotics have a protective effect on the intestinal wall due to an ability to improve barrier function and prevent adhesion, internalization and proliferation of enteropathogenic bacteria [25]. Indeed, gelatine tannate has a strong protective effect against adhesion of *Salmonella* Typhimurium and *E. coli* in Caco-2 and CacoGoblet models. These results are in good agreement with the barrier-stabilizing properties of gelatine tannate, quantified as reduction of paracellular flux by TEER and lucifer yellow measurements, and corroborate preliminary data about physical barrier formation by gelatine tannate. Moreover, selected health-promoting bacteria are known to improve microbial balance and may prove beneficial in reducing the duration of acute infectious diarrhea. Health-promoting bacteria can survive in the gastrointestinal tract, adhere to mucosal surfaces and metabolize available energy sources from dietary compounds. Bacteria such as lactobacilli, bifidobacteria and streptococci have a luminal action: antitoxic activity against specific pathogens; antimicrobial activity; preservation of tight junctions; and modulation of intestinal flora, or enhancement of conditions favoring growth of the endogenous bacterial flora [17–20]. Health-promoting bacteria also have a trophic effect on the mucosa, and anti-inflammatory mucosal effects.

Gelatine tannate may reduce the time required for probiotics to become active. This is based on the assumption that gelatine tannate will resolve diarrhea within the first 12 h of starting treatment, thereby permitting health-promoting bacteria to sit inside the bowel and exert intraluminal antidiarrheal activity earlier than if probiotics were administered alone. Furthermore, synergistic activity can be expected from combined administration of gelatine tannate and probiotics. Such activity might comprise combined competition with pathogenic bacteria, thereby preventing mucosal adhesion, internalization and proliferation of such organisms.

Conclusion

Although further research is needed to clearly define the mechanism of gelatine tannate/probiotic action in the clinical setting, preliminary data suggest that gelatine tannate has a protective film-forming effect through interaction with tight junctions and mucoadhesive properties: interaction with mucins permits gelatine tannate to adhere to apical epithelial cells. In this *in vitro* trial, the strong increase in TEER and strong decrease in lucifer yellow flux produced by gelatine tannate/probiotic suggested early and maintained restoration of tight-junction function and membrane integrity in CacoGoblet cells. Thus, at 1 and 24 h, gelatine tannate/probiotic had strong protective efficacy against *E. coli* damage in CacoGoblet medium.

Future perspective

The precise mechanism of gelatine tannate/probiotic action warrants further investigation, since this combination has potential broad clinical utility in the management of diarrhea of divergent causes (e.g., infectious gastroenteritis, antibiotic therapy and chemotherapy) in population groups of all ages.

Summary points

- Pathogenic microorganisms can adhere to the intestinal mucosa and influence cellular tight-junction function, leading to diarrhea.
- CacoGoblet[®] medium provides an enhanced physiological model of human intestinal permeability for evaluating antidiarrheal agents.
- This study compared the protective efficacy of gelatine tannate plus inactivated probiotic with that of various other antidiarrheal agents in *Escherichia coli*-inoculated CacoGoblet medium.
- Gelatine tannate/probiotic increased transepithelial electrical resistance by approximately 125–150%, and reduced lucifer yellow paracellular flux by about 1–2%.
- The protective efficacy of gelatine tannate/probiotic against *E. coli*-induced reduction of membrane integrity manifests early and is maintained for 24 h.

Financial & competing interests disclosure

The authors are employees of VitroScreen, Italy; an independent *in vitro* research laboratory used by Novintethical to perform the study. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed. The authors thank Content Ed Net, Madrid, Spain, for providing writing and editorial assistance, with funding from Novintethical, SA.

Ethical conduct of research

The authors state that they have obtained appropriate institutional review board approval or have followed the principles outlined in the Declaration of Helsinki for all human or animal experimental investigations. In addition, for investigations involving human subjects, informed consent has been obtained from the participants involved.

References

1. De Servi B, Ranzini F, Meloni M. Antidiarrhoeal agents and paracellular permeability of *E. coli*-infected Caco-Goblet intestinal model. Novintethical www.vitroscreen.com/pdf/2013%20POSTER%20CACO-GOBLET-GASTROENTEROLOGY.pdf
2. Esteban Carretero J, Durban Reguera F, Lopez-Argueta Alvarez S, Lopez Montes J. A comparative analysis of response to vs. ORS + gelatin tannate pediatric patients with acute diarrhea. *Rev. Esp. Enferm. Dig.* 101(1), 41–48 (2009).
3. Zhao H, Zhang H, Wu H *et al.* Protective role of 1,25(OH)₂ vitamin D₃ in the mucosal injury and epithelial barrier disruption in DSS-induced acute colitis in mice. *BMC Gastroenterol.* 12, 57 (2012).
4. Allegrini A, Costantini M. TasectanTM (gelatine tannate) for the treatment of acute gastroenteritis in children: preliminary results from a study conducted to evaluate its efficacy and safety. http://tasectan.ie/wp-content/uploads/2012/10/10.1.-Poster-TASECTANT-Belgium_March-20111.pdf
5. Ferruzza S, Rossi C, Scarino ML, Sambuy Y. A protocol for *in situ* enzyme assays to assess the differentiation of human intestinal Caco-2 cells. *Toxicol. In Vitro* 26(8), 1247–1251 (2012).
6. Ferruzza S, Rossi C, Scarino ML, Sambuy Y. A protocol for differentiation of human intestinal Caco-2 cells in asymmetric serum-containing medium. *Toxicol. In Vitro* 26(8), 1252–1255 (2012).

7. Prieto P, Hoffmann S, Tirelli V *et al.* An exploratory study of two Caco-2 cell models for oral absorption: a report on their within-laboratory and between-laboratory variability, and their predictive capacity. *Altern. Lab. Anim.* 38(5), 367–386 (2010).
8. Angelis ID, Turco L. Caco-2 cells as a model for intestinal absorption. *Curr. Protoc. Toxicol.* Chapter 20, Unit 20–26 doi:10.1002/0471140856.tx2006s47 (2011) (Epub ahead of print).
9. Kim JS, Mitchell S, Kijek P, Tsume Y, Hilfinger J, Amidon GL. The suitability of an *in situ* perfusion model for permeability determinations: utility for BCS class I biowaiver requests. *Mol. Pharm.* 3(6), 686–694 (2006).
10. Pontier C, Pachot J, Botham R *et al.* HT29-MTX and Caco-2/TC7 monolayers as predictive models for human intestinal absorption: role of the mucus layer. *J. Pharm. Sci.* 90(10), 1608–1619 (2001).
11. Hilgendorf C, Spahn-Langguth H, Regardh CG, Lipka E, Amidon GL, Langguth P. Caco-2 versus Caco-2/HT29-MTX co-cultured cell lines: permeabilities via diffusion, inside- and outside-directed carrier-mediated transport. *J. Pharm. Sci.* 89(1), 63–75 (2000).
12. Ruszczynski M, Urbanska M, Szajewska H. Gelatin tannate for treating acute gastroenteritis: a systematic review. *Ann. Gastroenterol.* 27(2), 121–124 (2014).
13. Bueno L, Sekkal S, Theodoru V, Dattilo M. OP249. Undissociated gelatine tannate reduces intestinal leakiness and mucosa inflammation by forming a protective biofilm: results from *in vitro* and *in vivo* studies. *United European Gastroenterol. J.* 1(Suppl. 1), A75–A76 (2013).
14. Lopetuso LR, Scaldaferri F, Bruno G, Petito V, Franceschi F, Gasbarrini A. The therapeutic management of gut barrier leaking: the emerging role for mucosal barrier protectors. *Eur. Rev. Med. Pharmacol. Sci.* 19(6), 1068–1076 (2015).
15. Michalek D, Kolodziej M, Konarska Z, Szajewska H. Efficacy and safety of gelatine tannate for the treatment of acute gastroenteritis in children: protocol of a randomised controlled trial. *BMJ Open* 6(2), e010530 (2016).
16. Scaldaferri F, Lopetuso LR, Petito V *et al.* Gelatin tannate ameliorates acute colitis in mice by reinforcing mucus layer and modulating gut microbiota composition: emerging role for ‘gut barrier protectors’ in IBD? *United European Gastroenterol. J.* 2(2), 113–122 (2014).
17. Thomas DW, Greer FR. Probiotics and prebiotics in pediatrics. *Pediatrics* 126(6), 1217–1231 (2010).
18. Putaala H, Salusjarvi T, Nordstrom M *et al.* Effect of four probiotic strains and *Escherichia coli* O157:H7 on tight junction integrity and cyclo-oxygenase expression. *Res. Microbiol.* 159(9–10), 692–698 (2008).
19. Alemka A, Clyne M, Shanahan F, Tompkins T, Corcionivoschi N, Bourke B. Probiotic colonization of the adherent mucus layer of HT29MTXE12 cells attenuates *Campylobacter jejuni* virulence properties. *Infect. Immun.* 78(6), 2812–2822 (2010).
20. Ahrne S, Hagslatt ML. Effect of lactobacilli on paracellular permeability in the gut. *Nutrients* 3(1), 104–117 (2011).
21. Adams CA. The probiotic paradox: live and dead cells are biological response modifiers. *Nutr. Res. Rev.* 23(1), 37–46 (2010).
22. Kataria J, Li N, Wynn JL, Neu J. Probiotic microbes: do they need to be alive to be beneficial? *Nutr. Rev.* 67(9), 546–550 (2009).
23. Indriyani A, Juffrie M, Setyati A. Effects of live versus heat-killed probiotics on acute diarrhea in young children. *Paediatr. Indones* 52, 249–254 (2012).
24. Sawada N, Murata M, Kikuchi K *et al.* Tight junctions and human diseases. *Med. Electron Microsc.* 36(3), 147–156 (2003).
25. Resta-Lenert S, Barrett KE. Live probiotics protect intestinal epithelial cells from the effects of infection with enteroinvasive *Escherichia coli* (EIEC). *Gut* 52(7), 988–997 (2003).

Matrix-assisted laser desorption/ionization–time of flight mass spectrometry: protocol standardization and database expansion for rapid identification of clinically important molds

Saikat Paul¹, Pankaj Singh¹, Shivaprakash M Rudramurthy¹, Arunaloake Chakrabarti¹ & Anup K Ghosh^{*,1}

¹Department of Medical Microbiology, Postgraduate Institute of Medical Education & Research, Chandigarh-160012, India

* Author for correspondence: Tel.: +91 172 275 5156; Fax: +91 172 274 4401; anupkg3@gmail.com

Aim: To standardize the matrix-assisted laser desorption ionization–time of flight mass spectrometry protocols and expansion of existing Bruker Biotyper database for mold identification. **Materials & methods:** Four different sample preparation methods (protocol A, B, C and D) were evaluated. **Results:** On analyzing each protein extraction method, reliable identification and best log scores were achieved through protocol D. The same protocol was used to identify 153 clinical isolates. Of these 153, 123 (80.3%) were accurately identified by using existing database and remaining 30 (19.7%) were not identified due to unavailability in database. On inclusion of missing main spectrum profile in existing database, all 153 isolates were identified. **Conclusion:** Matrix-assisted laser desorption ionization–time of flight mass spectrometry can be used for routine identification of clinically important molds.

First draft submitted: 9 June 2017; Accepted for publication: 27 July 2017; Published online: 27 October 2017

Keywords: α -cyano-4-hydroxycinnamic acid matrix • acetonitrile • database • formic acid • log score • main spectrum profiles • MALDI–TOF MS • molds • MSPs • sabouraud dextrose agar • trifluoroacetic acid

Invasive fungal infection (IFI) is a serious life-threatening infection associated with high morbidity and mortality. Despite available effective antifungal therapy, rapid diagnosis is crucial to reduce the mortality [1]. However, the diagnosis of fungal infection still relies on conventional techniques (microscopy, histopathology and culture), which have several limitations including low sensitivity, long turnaround time and requirement of expert taxonomist to identify the fungus [2]. In addition, with the changing epidemiology and expansion of spectrum of fungi causing disease, the situation has become more complicated for conventional diagnosis [3]. Although, molecular techniques based on DNA including PCR, real-time PCR, hybridization techniques and Sanger sequencing have improved the identification of fungi, these methods are expensive, labor intensive and prone to contamination due to multistep procedures [4–6].

In recent years, proteomic-based approaches, specifically matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI–TOF MS)-based identification emerge as a rapid diagnostic techniques in clinical microbiology [7–9]. In comparison to sequencing, the efficiency of MALDI–TOF MS is equivalent to DNA sequencing and the consumable cost is minimum [7]. Nevertheless, the identification of filamentous fungi by MALDI–TOF MS still lags behind due to lack of a common universal protein extraction protocol and limited database availability [10–12].

Several studies reported the identification of filamentous fungi by MALDI–TOF MS [11–22]. However, most of these studies have used heterogeneous protein preparation techniques and targeted only commonly isolated fungal genera such as *Aspergillus*, *Fusarium*, dermatophytes, etc. In addition, the MALDI spectra can be influenced by various factors including culture conditions, age of culture, extraction methods, the type of matrix and the spectrometer used [9,11,13–14,17–19]. Moreover, the pigment production by some fungi like *Aspergillus* and *Fusarium*

can also affect the spectra profile [15,18]. Due to these limitations, standardized consensus protocol has not been established for mold identification by MALDI-TOF MS, till now.

In this study, we standardized and compared the various protein extraction protocols and added new main spectrum profiles (MSPs) of some rare filamentous fungi to the Bruker Biotyper database (version 4.0). Initially, the protein extraction protocols for MALDI-TOF MS-based identification were standardized on reference strains. The best protocol was further evaluated by using sequentially confirmed isolates and the MSPs of the unidentified isolates were incorporated in the existing database for further improvement in identification.

Materials & methods

Fungal strains

For MALDI-TOF MS-based identification, *Aspergillus flavus* (ATCC 204304), *Rhizopus oryzae* (CBS 131512) and *Fusarium solani* (NCCPF 58A0057) were used for initial standardization. A total of 153 isolates obtained from our National Culture Collection of Pathogenic Fungi [23] were used for validation. All these isolates were confirmed by conventional and as well as PCR sequencing of internal transcribed spacer (*ITS*) and 28S ribosomal RNA gene's *D1/D2* regions.

Protein extraction protocols

For the optimization of protein extraction protocol, four different methods were assessed. All these were similar to the Bruker-recommended protocol with considerable modifications. Acetonitrile (ACN), formic acid (FA), trifluoroacetic acid (TFA), α -cyano-4-hydroxycinnamic acid (HCCA) matrix and double-distilled water (DDW) were the key components of each protocol. Fungal isolates used in protocol A, B and C were cultured on Sabouraud dextrose agar (SDA) and incubated at 28°C for 24–48 h. The four protocols were as follows:

Protocol A

Mycelial growth was taken in 10 μ l of 70% ethanol and directly spotted on MALDI plate (Bruker Daltonik GmbH, Bremen, Germany) in triplicates and air dried. A 1 μ l 98% FA was added to each spot and air dried. Each spot was overlaid with 1 μ l matrix solution (50 μ l ACN, 47.5 μ l DDW, 2.5 μ l TFA and 1 mg HCCA) followed by MALDI-TOF MS identification.

Protocol B

Fungal growth was crushed in liquid nitrogen to make fine powder. Then, 50–80 μ l of FA and ACN were added and mixed properly. A 30 μ l emulsified sample was transferred to a microcentrifuge tube and spin at 10,000 r.p.m. for 5 min. A 3–5 μ l supernatant was spotted (by adding multiple sequential drops in each spot) onto a MALDI target plate and air dried. Each spot was covered with 1 μ l matrix solution, air dried and analyzed in MALDI-TOF MS.

Protocol C

Fungal growth was suspended in a 1.5 ml DDW, vortexed and centrifuged at 13,000 r.p.m. for 10 min. 1 ml supernatant was removed and 500–600 μ l absolute ethanol was added followed by incubation for 2–5 min at room temperature. After centrifugation, supernatant was removed and dried the pellet by vacuum concentrator (Savant™ DNA SpeedVac™ concentrator, Thermo Fisher Scientific, MA, USA). Then, 60–90 μ l of 98% FA and ACN were added to suspend the pellet and vortexed for 10–15 min with 0.5–0.75 mm acid-washed glass beads (Sigma-Aldrich, Saint-Quentin-Fallavier, L'Isle D'Abeau Chesnes, France). After centrifugation, 3–5 μ l samples were spotted followed by addition of matrix solution and MALDI-TOF MS analysis.

Protocol D

Fungal growth was inoculated in 5 ml of Sabouraud dextrose broth in 15 ml centrifuge tube and spin in a test tube rotator (Rotospin Tarsons, West Bengal, India) at 50 r.p.m. for 12–24 h at 28°C. The tubes were kept in static condition for 10 min for sedimentation of hyphal flakes. By using blunt-end pipette tips, 1.5 ml of suspension was transferred to a microcentrifuge tube and centrifuged at maximum speed for 5 min. The media was completely removed and 1.5 ml sterile DDW was added for washing the pellet. After centrifugation at 13,000 r.p.m., 1.2 ml supernatant was removed and replaced with 700 μ l absolute ethanol and left for 5 min. The ethanol content was completely removed by pipetting and vacuum concentrator. The dried pellet was emulsified in 50–80 μ l FA and

incubated at room temperature for 5 min followed by addition of 50–80 μ l ACN. The emulsion was vortexed for 5–10 min with 0.5 mm acid-washed glass beads for proper lysis. The rest of the steps were similar and were previously described in above protocols.

Mass spectrum acquisition & analysis

All the MALDI–TOF MS assays were carried out on MALDI Microflex LT mass spectrometer (Bruker Daltonik, GmbH, Bremen, Germany) by using the flex control software (Flex control 3.4, Bruker Daltonik, GmbH). Ionized spectra from the individual spot were acquired in a linear positive mode at a laser frequency of 60 Hz within a mass range of 2–20 kDa. The instrument settings for automatic run were as follows: ion source 1: 20 kV; ion source 2: 18.05 kV, and lens: 9.0 kV. The spectrum of each sample was generated by randomly heating six different positions of a single spot (40 laser short steep) and produced a total of 240 laser shorts. The instrument was calibrated by using the Bruker Bacterial Test Standard *Escherichia coli* (BTS 8255343). Subsequently MALDI BioTyper (MALDI Biotyper 3.1, Bruker Daltonik, GmbH) was used for automated smoothing, normalization, baseline subtraction and peak selection of different spectra. Most significant peaks were selected to create a profile for each sample, which was then compared with the MALDI BioTyper database (version 4.0) and filamentous fungal library (version 1.0). In these databases, MALDI Biotyper database v4.0 contains 775 entries for different fungal species, whereas filamentous fungal library v1.0 consists of 365 entries for only filamentous fungi. Results were expressed as log score ranging from a value of 0–3 by using pattern matching algorithm. As per the manufacturer's instructions a score more than or equal to two is reliable for species identification; a score between 1.7 and 1.99 reliably identifies the genus of any isolate; and a score below 1.7 is considered unreliable for identification. Any species misidentified or unavailable in the manufacturer's database was resolved by PCR sequencing [7].

Calculation of log score

The log score is calculated by MALDI Biotyper software. In brief, raw mass spectra was converted into peak lists followed by matching of these peaks with reference database using pattern matching algorithm. The matching algorithm computes three essential factors of the reference and unknown spectra. The log score is calculated by multiplication of these three factors (each factor can reach a maximum value of one), which includes number of signal matches in the unknown spectrum against reference spectrum, number of signal matches in the reference spectrum against the unknown spectrum and similarity of the intensity profile of matching signals. These three values are multiplied together and result is normalized by 1000. The MALDI log score was the log of this result. The maximum log score value can be three ($\log_{1000} = 3$). All log scores are ranked from highest to lowest. The higher log score value results in a species-level identification.

Development of new in-house database/spectrum of new projections

Fungal strains confirmed by PCR sequencing (*ITS* and *D1/D2* region) were included in Bruker MALDI Biotyper database. The sample preparation was performed according to protocol D. Each sample was spotted at eight positions onto the target plate and measured three-times per spot. Finally, 24 spectra were acquired for each isolate. For the automatic measurement, 24 spectra were analyzed with flexControl method (e.g., MBT_FC.par) and autoExecute method (e.g., MBT_autoX.axe). Initially, Flex Analysis (version 3.4) was used for manual analysis of the quality and consistency of mass spectra by smoothing and baseline subtraction. After analysis, minimum 20 best spectra were opened in MALDI Biotyper software simultaneously by selecting the respective file, and processing (adjust mass, smooth, subtract baseline, normalize, pick peaks and batch processing) of acquired spectra was performed and saved to the MALDI Biotyper database as the MSP of the new species. For quality control, all newly created MSPs were checked against the existing Bruker Biotyper database v4.0.

Statistical analysis

All statistical analysis was done by using SPSS (IBM SPSS statistics 22) software. The log score of reference isolates by different protein extraction method was presented as median with interquartile range, whereas log score of validation set of isolates as mean $\pm$ standard deviation. Kruskal–Wallis test was used for comparing protocol D with other methods. The statistical analysis was executed by employing two-sided tests with $p \leq 0.05$ indicating significance level.

Figure 1. Comparison of main spectrum profiles of four different protein extraction methods. Mass spectra of *Aspergillus flavus* ATCC-204304 obtained with the different protein extraction procedures. a.u.: Arbitrary unit; intens.: Intensity.

Results

Among four different protein extraction methods, protocols C and D correctly identified the reference strains with log score >1.7 , whereas protocols A and B give no reliable identification with log score <1.7 . On comparing log score between protocols C and D, the statistical analysis revealed significant log score difference in between protocols C and D (*Aspergillus flavus*: 1.8 vs 2.3, $p < 0.001$; *Fusarium solani*: 1.8 vs 2.3, $p < 0.001$; *Rhizopus arrhizus*: 1.8 vs 2.2, $p < 0.001$). As the best log score was achieved through protocol D, this protocol was adopted for the remaining study. The log score and MALDI spectra of different protein extraction methods are presented in Figures 1 & 2.

Out of 153 isolates, 80.3% (123) were successfully identified by using existing Bruker database (Table 1 & Figure 3). The remaining 19.6% (30) isolates gave no reliable identification due to the lack of reference spectra in Bruker database. We updated MALDI Biotyper for those 30 isolates of 12 different missing species including *Aspergillus nidulans* (4), *Aspergillus tetrazonus* (3), *Aspergillus sclerotium* (1), *Aspergillus conicus* (1), *Apophysomyces elegans* (1), *Apophysomyces variabilis* (10), *Apophysomyces ossiformis* (1), *Mucor indicus* (3), *Apophysomyces trapeziformis* (1), *Fusarium napiforme* (1), *Cunninghamella bertholletiae* (3) and *Lichtheimia ramosa* (1). MALDI-TOF MS correctly identified all these 30 isolates using the new MSP profiles incorporated into the database with log score ≥ 1.98 (Table 2). Distribution of species-specific log scores showed that the majority of strains (122, 79.7%) could be successfully identified between scores of 1.7 and 2.4 (Table 1).

On comparing, MALDI-TOF MS with PCR sequencing, MALDI-TOF MS identified hyaline fungi with 100% accuracy with receiver operating characteristic (ROC) cut-off log score ≥ 2.0 for *Paecilomyces lilacinus*, *Fusarium moniliforme*, *Fusarium proliferatum*, *Aspergillus clavatus*, *Aspergillus tetrazonus*, *Paecilomyces variotii*, *Fusarium oxysporum*, *Cunninghamella elegans*, *Cunninghamella bertholletiae*, *Mucor indicus* and *Apophysomyces elegans*. Spectral cut-off computed analysis showed 100% accuracy with ROC cut-off log score of ≥ 1.70 for *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus niger*, *Aspergillus terreus*, *Aspergillus oryzae*, *Aspergillus variabilis*, *Fusarium solani*, *Fusarium incarnatum*, *Rhizopus oryzae*, *Rhizopus microsporus*, *Rhizomucor pusillus* and *Lichtheimia corymbifera* (Table 3). The overall predictive power, accuracy and efficiency of MALDI-TOF MS were comparable to sequencing.

Figure 2. Comparison of four different protein extraction methods using log score.

Among four different protein extraction methods, protocols C and D correctly identified the reference strains with log score >1.7, whereas protocols A and B give unreliable identification with log score <1.7. The statistical analysis revealed significant log score difference in between protocol D and other protocols.

** p-value less than 0.01.

**** p-value less than 0.0001.

Fungal species (12) that could not be reliably identified by MALDI-TOF MS had log scores varying from 0.87 to 1.61. These species were identified by PCR sequencing (*ITS2* and *D1/D2* region) and MSP were included in Bruker Biotyper database. After addition of MSP of those species in existing Bruker database, all species were identified by MALDI-TOF MS (Figure 3). While comparing, the existing Bruker database with Bruker plus in-house database, 76 (49.7%) versus 104 (68%) isolates correctly identify to species level, 47 (30.7%) versus 49 (32%) isolates to genus level and 30 (19.6%) versus 0 (0%) isolates gave unreliable identification (Table 2 & Figure 4).

Discussion

In this study, four methods of sample preparation were evaluated. Initially, optimum extraction procedure was standardized. For this, the fungi were cultured in SDA, the most common medium used for culturing fungi in clinical laboratories. The direct-on-plate method (protocol A) was faster, consumes fewer resources but the best match log score (LS) was significantly lower. Among off-plate method, liquid nitrogen (protocol B) also gave an unreliable identification with low LS values. The inadequate protein extraction may be the reason of low sensitivity of both these methods [10,19]. Only protocol C (solid media) and D (liquid media) were associated with both reliable identification and high log score. As maximum log score was achieved through protocol D, we used subsequently the same techniques. On comparing protocols C and D, heterogeneous fungal growth and pigment production by hyaline fungi were observed on solid media [15,18]. On the contrary, uniform growth without any pigment production was observed in liquid media. As the pigments may interfere with MALDI mass spectra [18], this may be the reason of maximum sensitivity of protocol D. In addition, Lau *et al.* (2013) also reported cross identification among database, while using solid media [11]. The difference in protein extraction protocol and database diversity may be the possible reason for the difference in sensitivity between protocols C and D [24]. Moreover, sporulating and nonsporulating fungi behave similarly in liquid medium, whereas in solid media fungal growth may contain spores/mycelia or a mixture of spores and mycelia, which may alter spectra. In protocol C, agar contamination may be also a problem during picking growth from solid media, whereas in liquid media only minute flakes are formed. So there is no chance of media contamination. During protein extraction, the minute flakes of liquid media are also easy to break in comparison to solid media fungal growth. Thus, choice of protocol D is most rational due to its sensitivity, reproducibility and easy to perform techniques.

Table 1. Species distribution of isolates across various matrix-assisted laser desorption/ionization Biotyper log scores for validation set by existing Bruker database.

S. No.	Species	Number of isolates	No. of isolates with different log score		
			<1.7	1.7–1.99	2.0–3.0
1	<i>Aspergillus clavatus</i>	3	0	0	3
2	<i>Aspergillus conicus</i>	1	1	0	0
3	<i>Aspergillus flavus</i>	21	0	12	9
4	<i>Aspergillus fumigatus</i>	13	0	8	5
5	<i>Aspergillus nidulans</i>	4	4	0	0
6	<i>Aspergillus niger</i>	15	0	2	13
7	<i>Aspergillus oryzae</i>	4	0	2	2
8	<i>Aspergillus sclerotiorum</i>	1	1	0	0
9	<i>Aspergillus terreus</i>	11	0	3	8
10	<i>Aspergillus tetrazonus</i>	3	3	0	0
11	<i>Apophysomyces elegans</i>	1	1	0	0
12	<i>Apophysomyces ossiformis</i>	1	1	0	0
13	<i>Apophysomyces trapeziformis</i>	1	1	0	0
14	<i>Apophysomyces variabilis</i>	10	10	0	0
15	<i>Cunninghamella bertholletiae</i>	3	3	0	0
16	<i>Cunninghamella elegans</i>	2	0	0	2
17	<i>Fusarium incarnatum</i>	5	0	5	0
18	<i>Fusarium moniliforme</i>	4	0	1	3
19	<i>Fusarium napiforme</i>	1	1	0	0
20	<i>Fusarium oxysporum</i>	2	0	0	2
21	<i>Fusarium proliferatum</i>	3	0	0	3
22	<i>Fusarium solani</i>	8	0	4	4
23	<i>Lictheimia corymbifera</i>	4	0	2	2
24	<i>Lictheimia ramosa</i>	1	1	0	0
25	<i>Mucor indicus</i>	3	3	0	0
26	<i>Paecilomyces lilacinus</i>	7	0	0	7
27	<i>Paecilomyces variotii</i>	3	0	0	3
28	<i>Rhizomucor pusillus</i>	2	0	1	1
29	<i>Rhizopus microsporus</i>	5	0	3	2
30	<i>Rhizopus oryzae</i>	11	0	4	7
Total		153	30	47	76

S. No.: Serial number.

For identification by MALDI-TOF MS, the database is equally important [7]. In the present study, after the standardization of protocol with reference strains, the protocol is further validated on clinical strains. The existing Bruker database failed to identify 30 of 153 isolates. Those 30 isolates included *A. nidulans*, *A. tetrazonus*, *A. sclerotium*, *A. conicus*, *A. elegans*, *A. variabilis*, *A. ossiformis*, *M. indicus*, *A. trapeziformis*, *F. napiforme*, *C. elegans*, *C. bertholletiae* and *L. ramosa*, which are clinically relevant. Similarly, a study by Lau *et al.* (2013) also reported low-mold identification efficiency of Bruker database in comparison to NIH database [11]. The low fungal diversity of Bruker database may be the main reason of unreliable mold identification. In few decades, the fungal epidemiology is also completely changed; rare fungi are becoming common [25]. Among these isolates, the incidence of mucorales is rapidly increasing in India with an increase in susceptible population (diabetic patients) [26]. As mucorales causes rapid angioinvasion, the misidentification may be critical for the morbidity and mortality of patients. Similarly, in our study, the under-representation of those species in the existing Bruker database was the reason for non-identifications [7,14]. On addition of MSP in Bruker database, all those fungal isolates could be identified by protocol D.

The database can also be a critical factor in genus or species level identification. Bruker recommended the log score >2.0 for species level identification. On the contrary, we observed that some fungi can be identified up to

Figure 3. Comparison of MALDI-TOF MS with DNA sequencing before and after database expansion.

A donut chart depicting the species-specific distribution and performance of MALDI-TOF MS using existing Bruker database (middle ring), DNA sequencing (inner ring) and MALDI-TOF MS using Bruker plus in-house database for the validation set of fungal strains. Numbers indicate the number of strains identified by the respective techniques.

MALDI-TOF MS: Matrix-assisted laser desorption ionization–time of flight mass spectrometry; MSP: Main spectrum profile.

species level even at log score <2.0. In this study, fungal species such as *A. flavus*, *A. fumigatus*, *A. niger*, *A. terreus*, *F. solani*, *F. moniliforme*, *F. incarnatum*, *F. proliferatum*, *R. oryzae*, *R. microsporus*, *R. pusillus* and *L. corymbifera* identified up to species level at log score >1.7. Whereas, *P. lilacinus*, *F. moniliforme*, *A. clavatus*, *A. nidulans*, *P. variotii*, *F. oxysporum*, *C. bertholletiae*, *M. indicus* and *A. elegans* identified up to species level at log score ≥ 2 . The under-representation of these fungal species in Bruker database may be the reason of high log-score cut off.

In this study, protocol D (modified version of Bruker protocol for filamentous fungi) identified isolates with significantly higher log score compared with other protocols. Further, this protocol was used for the expansion of existing Bruker database. The most important finding of this study was the inclusion of some rare but clinically important filamentous fungi (*A. tetrazonus*, *A. conicus* and four different species of *Apophysomyces*) in the existing database. The reliable identification of fungi depends on the complexity of MALDI-TOF MS database. However, this type of study is useful for the community, if databases are made publicly available. So, a global MALDI database with multiple entries for each species from different geographical regions can be created to resolve this problem. In addition, the database should be regularly updated to include rare clinical fungal species. The main limitation of

Table 2. Matrix-assisted laser desorption/ionization Biotyper log score before and after in-house database creation.

S. No.	Species	Number of isolates	Bruker database mean log score $\pm$ SD	Bruker plus in-house database mean log score $\pm$ SD	p-value
1	<i>Aspergillus nidulans</i>	4	1.54 $\pm$ 0.06	2.41 $\pm$ 0.19	<0.002
2	<i>Aspergillus tetrazonus</i>	3	1.17 $\pm$ 0.11	2.52 $\pm$ 0.07	<0.001
3	<i>Aspergillus sclerotiorum</i>	1	0.9 $\pm$ 0.0	2.5 $\pm$ 0.0	–
4	<i>Aspergillus conicus</i>	1	1.1 $\pm$ 0.0	2.46 $\pm$ 0.0	–
5	<i>Fusarium napiforme</i>	1	1.4 $\pm$ 0.0	2.5 $\pm$ 0.0	–
6	<i>Apophysomyces elegans</i>	1	0.87 $\pm$ 0.0	2.6 $\pm$ 0.0	–
7	<i>Apophysomyces variabilis</i>	10	0.96 $\pm$ 0.11	2.15 $\pm$ 0.16	<0.001
8	<i>Apophysomyces ossiformis</i>	1	1.03 $\pm$ 0.0	2.57 $\pm$ 0.0	–
9	<i>Apophysomyces trapeziformis</i>	1	1.01 $\pm$ 0.0	2.48 $\pm$ 0.0	–
10	<i>Mucor indicus</i>	3	0.97 $\pm$ 0.07	2.37 $\pm$ 0.22	<0.001
11	<i>Cunninghamella bertholletiae</i>	3	1.04 $\pm$ 0.06	2.36 $\pm$ 0.24	<0.008
12	<i>Lichtheimia ramosa</i>	1	1 $\pm$ 0.0	2.4 $\pm$ 0.0	–

S. No.: Serial number; SD: Standard deviation.

Table 3. Comparison of MALDI-TOF MS-based identification with DNA sequencing of selected mycelial fungi.

S. No	Species	Number of isolates	Correctly identified (n)	AUC (95% CI)	ROC	κ (95% CI)
1	<i>Aspergillus flavus</i>	21	21	1.00 (0.99–1.00)	≥ 1.79	1.00 (1.00–1.00)
2	<i>Aspergillus niger</i>	15	15	1.00 (0.99–1.00)	≥ 1.98	1.00 (1.00–1.00)
3	<i>Aspergillus fumigatus</i>	13	13	1.00 (0.99–1.00)	≥ 1.85	1.00 (1.00–1.00)
4	<i>Aspergillus terreus</i>	11	11	1.00 (0.99–1.00)	≥ 1.87	1.00 (1.00–1.00)
5	<i>Rhizopus oryzae</i>	11	11	1.00 (0.99–1.00)	≥ 1.89	1.00 (1.00–1.00)
6	<i>Apophysomyces variabilis</i>	10	10	1.00 (0.99–1.00)	≥ 1.70	1.00 (1.00–1.00)
7	<i>Fusarium solani</i>	8	8	1.00 (0.99–1.00)	≥ 1.89	1.00 (1.00–1.00)
8	<i>Paecilomyces lilacinus</i>	7	7	1.00 (0.99–1.00)	≥ 2.0	1.00 (1.00–1.00)
9	<i>Fusarium incarnatum</i>	5	5	1.00 (0.99–1.00)	≥ 1.79	1.00 (1.00–1.00)
10	<i>Rhizopus microsporus</i>	5	5	1.00 (0.99–1.00)	≥ 1.92	1.00 (1.00–1.00)
11	<i>Aspergillus oryzae</i>	4	4	1.00 (0.99–1.00)	≥ 1.95	1.00 (1.00–1.00)
12	<i>Fusarium moniliforme</i>	4	4	1.00 (0.99–1.00)	≥ 2.04	1.00 (1.00–1.00)
13	<i>Lichtheimia corymbifera</i>	4	4	1.00 (0.99–1.00)	≥ 1.97	1.00 (1.00–1.00)
14	<i>Fusarium proliferatum</i>	3	3	1.00 (0.99–1.00)	≥ 2.1	1.00 (1.00–1.00)
15	<i>Aspergillus clavatus</i>	3	3	1.00 (0.99–1.00)	≥ 2.0	1.00 (1.00–1.00)
16	<i>Aspergillus tetrazonus</i>	3	3	1.00 (0.99–1.00)	≥ 2.18	1.00 (1.00–1.00)
17	<i>Mucor indicus</i>	3	3	1.00 (0.99–1.00)	≥ 2.18	1.00 (1.00–1.00)
18	<i>Cunninghamella bertholletiae</i>	3	3	1.00 (0.99–1.00)	≥ 2.18	1.00 (1.00–1.00)
19	<i>Paecilomyces variotii</i>	3	3	1.00 (0.99–1.00)	≥ 2.0	1.00 (1.00–1.00)
20	<i>Cunninghamella elegans</i>	2	2	1.00 (0.99–1.00)	≥ 2.1	1.00 (1.00–1.00)
21	<i>Fusarium oxysporum</i>	2	2	1.00 (0.99–1.00)	≥ 2.0	1.00 (1.00–1.00)
22	<i>Rhizomucor pusillus</i>	2	2	1.00 (0.99–1.00)	≥ 1.9	1.00 (1.00–1.00)

AUC: Area under curve; CI: Confidence interval; MALDI-TOF MS: Matrix-assisted laser desorption ionization–time of flight mass spectrometry; ROC: Receiver–operating characteristics; S. No.: Serial number.

this study was a less number of uncommon filamentous fungi used to expand the database. As the number of these species will be increased in MALDI database, the performance of MALDI-TOF MS will be improved.

Figure 4. Comparison of existing Bruker database with Bruker plus in-house database.
Genus and species level identification by Bruker database and Bruker plus in-house database.

Conclusion

In conclusion, our study demonstrates that MALDI–TOF MS-based identification can be used for filamentous fungi by protocol D. The expansion of MALDI database with multiple entries from different centers can further enhance the capability of identification of filamentous fungi.

Summary points

- Matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI–TOF MS) is an emerging diagnostic technique in clinical microbiology.
- MALDI–TOF MS-based identification is well established for bacteria and yeast. However, MALDI–TOF MS-based filamentous fungi identification still far lags behind.
- Protocol D can be used as a standardized consensus protocol for MALDI–TOF MS-based identification of filamentous fungi.
- Regular expansion of database is very crucial for reliable identification of emerging pathogenic fungi.

Acknowledgements

The authors duly acknowledge the Department of Science and Technology, Government of India for financial supports.

Financial & competing interest's disclosure

The study was financially supported by Department of Science and Technology, Government of India. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Ethical conduct of research

The authors state that they have obtained appropriate institutional review board approval or have followed the principles outlined in the Declaration of Helsinki for all human or animal experimental investigations. In addition, for investigations involving human subjects, informed consent has been obtained from the participants involved.

References

1. Badiee P, Hashemizadeh Z. Opportunistic invasive fungal infections: diagnosis & clinical management. *Indian J. Med. Res.* 139(2), 195–204 (2014).
2. Cuenca-Estrella M, Bernal-Martinez L, Buitrago MJ *et al.* Update on the epidemiology and diagnosis of invasive fungal infection. *Int. J. Antimicrob. Agents.* 32(Suppl. 2), S143–S147 (2008).
3. Richardson M, Lass-Flörl C. Changing epidemiology of systemic fungal infections. *Clin. Microbiol. Infect.* 14(Suppl. 4), 5–24 (2008).
4. Hsiue H-C, Huang Y-T, Kuo Y-L, Liao C-H, Chang T-C, Hsueh P-R. Rapid identification of fungal pathogens in positive blood cultures using oligonucleotide array hybridization. *Clin. Microbiol. Infect.* 16(5), 493–500 (2010).
5. Leaw SN, Chang HC, Sun HF, Barton R, Bouchara J-P, Chang TC. Identification of medically important yeast species by sequence analysis of the internal transcribed spacer regions. *J. Clin. Microbiol.* 44(3), 693–699 (2006).
6. Marcos JY, Pincus DH. Fungal diagnostics: review of commercially available methods. *Methods Mol. Biol.* 968, 25–54 (2013).
7. Ghosh AK, Paul S, Sood P *et al.* Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry for the rapid identification of yeasts causing bloodstream infections. *Clin. Microbiol. Infect.* 21(4), 372–378 (2015).
8. Lagacé-Wiens PRS, Adam HJ, Karlowsky JA *et al.* Identification of blood culture isolates directly from positive blood cultures using MALDI-TOF mass spectrometry and a commercial extraction system – analysis of performance, cost and turnaround time. *J. Clin. Microbiol.* 50(10), 3324–3328 (2012).
9. Van Veen SQ, Claas ECJ, Kuijper EJ. High-throughput identification of bacteria and yeast by matrix-assisted laser desorption/ionization-time of flight mass spectrometry in conventional medical microbiology laboratories. *J. Clin. Microbiol.* 48(3), 900–907 (2010).
10. Cassagne C, Cella A-L, Suchon P, Normand A-C, Ranque S, Piarroux R. Evaluation of four pretreatment procedures for MALDI-TOF MS yeast identification in the routine clinical laboratory. *Med. Mycol.* 51(4), 371–377 (2013).
11. Lau AF, Drake SK, Calhoun LB, Henderson CM, Zelazny AM. Development of a clinically comprehensive database and a simple procedure for identification of molds from solid media by matrix-assisted laser desorption/ionization-time of flight mass spectrometry. *J. Clin. Microbiol.* 51(3), 828–834 (2013).
12. Sleiman S, Halliday CL, Chapman B *et al.* Performance of matrix-assisted laser desorption/ionization-time of flight mass spectrometry for identification of *Aspergillus*, *Scedosporium*, and *Fusarium* spp. in the Australian Clinical Setting. *J. Clin. Microbiol.* 54(8), 2182–2186 (2016).
13. Cassagne C, Ranque S, Normand A-C *et al.* Mould routine identification in the clinical laboratory by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *PLoS ONE* 6(12), e28425 (2011).
14. De Carolis E, Posteraro B, Lass-Flörl C *et al.* Species identification of *Aspergillus*, *Fusarium* and *Mucorales* with direct surface analysis by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Clin. Microbiol. Infect.* 18(5), 475–484 (2012).
15. Dong H, Kempfner J, Marchetti-Deschmann M, Kubicek CP, Allmaier G. Development of a MALDI two-layer volume sample preparation technique for analysis of colored conidia spores of *Fusarium* by MALDI linear TOF mass spectrometry. *Anal. Bioanal. Chem.* 395(5), 1373–1383 (2009).
16. Schrödl W, Heydel T, Schwartze VU *et al.* Direct analysis and identification of pathogenic *Lichtheimia* species by matrix-assisted laser desorption/ionization-time of flight analyzer-mediated mass spectrometry. *J. Clin. Microbiol.* 50(2), 419–427 (2012).
17. Valentine NB, Wahl JH, Kingsley MT, Wahl KL. Direct surface analysis of fungal species by matrix-assisted laser desorption/ionization mass spectrometry. *Rapid Commun. Mass Spectrom.* 16(14), 1352–1357 (2002).
18. Hettick JM, Green BJ, Buskirk AD *et al.* Discrimination of *Aspergillus* isolates at the species and strain level by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry fingerprinting. *Anal. Biochem.* 380(2), 276–281 (2008).
19. Riat A, Hinrikson H, Barras V, Fernandez J, Schrenzel J. Confident identification of filamentous fungi by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry without subculture-based sample preparation. *Int. J. Infect. Dis.* 35, 43–45 (2015).
20. Erhard M, Hipler UC, Burmester A, Brakhage AA, Wöstemeyer J. Identification of dermatophyte species causing onychomycosis and tinea pedis by MALDI-TOF mass spectrometry. *Exp. Dermatol.* 17(4), 356–361 (2008).
21. Marinach-Patrice C, Lethuillier A, Marly A *et al.* Use of mass spectrometry to identify clinical *Fusarium* isolates. *Clin. Microbiol. Infect.* 15(7), 634–642 (2009).
22. Ranque S, Normand A-C, Cassagne C *et al.* MALDI-TOF mass spectrometry identification of filamentous fungi in the clinical laboratory. *Mycoses* 57(3), 135–140 (2014).
23. National Culture Collection of Pathogenic Fungi. www.nccpf.com
24. Schulthess B, Ledermann R, Mouttet F *et al.* Use of the Bruker MALDI Biotyper for identification of molds in the clinical mycology laboratory. *J. Clin. Microbiol.* 52(8), 2797–2803 (2014).
25. Enoch DA, Yang H, Aliyu SH, Micallef C. The changing epidemiology of invasive fungal infections. *Methods Mol. Biol.* 1508, 17–65 (2017).
26. Chakrabarti A, Singh R. Mucormycosis in India: unique features. *Mycoses* 57(Suppl. 3), 85–90 (2014).

γ -irradiation from radiotherapy improves the virulence potential of *Candida tropicalis*

Eliane M da Silva^{‡,1}, Brenda Kischkel^{‡,1}, Cristiane S Shinobu-Mesquita^{‡,1}, Patrícia S Bonfim-Mendonça^{‡,1}, Elaine SB Mansano^{‡,1}, Marcos A da Silva^{‡,2}, Jéssica F Barbosa^{‡,2}, Adriana Fiorini^{‡,3}, Luzmarina Hernandez^{‡,1}, Marcia C Furlaneto^{‡,4} & Terezinha IE Svidzinski^{‡,*}

¹Department of Clinical Analysis & Biomedicine/Postgraduate Program in Biosciences & Pathophysiology, Centro de Ciências da Saúde, Universidade Estadual de Maringá, Maringá, Paraná, Avenida Colombo, 5790, Maringá, PR, CEP 87020-900, Brazil

²Sant'ana Radiation Oncology Center of Maringá, Paraná, Brazil

³Department of Microbiology, Universidade Federal do Paraná, Setor Palotina, Paraná, Brazil

⁴Department of Microbiology, Universidade Estadual de Londrina, Londrina, Paraná, Brazil

* Author for correspondence: terezinha.svidzinski@gmail.com

‡ Authors contributed equally

Aim: To evaluate if radiation used in radiotherapy can cause changes in the virulence potential of *Candida tropicalis* ATCC 750. **Materials & methods:** *C. tropicalis* was exposed *in vitro* to identical dose and scheme of irradiation would be used in patients with head and neck cancer. Some virulence parameters were analyzed before and after irradiation. **Results:** Colony morphologies were irreversibly affected by irradiation. Increase in growth rate, filamentation, adhesion on cell lines and phagocytosis process were also observed. Overall the irradiated *C. tropicalis* cells became more efficient at causing systemic infection in mice. **Conclusion:** γ -radiation induced important changes in *C. tropicalis* increasing its virulence profile, which could directly affect the relationship between yeasts and hosts.

First draft submitted: 9 July 2017; Accepted for publication: 30 August 2017; Published online: 7 November 2017

Keywords: *Candida tropicalis* • head and neck neoplasms • radiation • virulence factors

Future
Medicine

Radiotherapy is one of the most common treatments for cancer, and can be used as a single or adjuvant cancer therapy [1,2]. As well as its effects in killing and controlling malignant cells, γ -irradiation can also cause injury to normal cells, including changes in protein expression [3,4].

Studies with cancer patients have shown that opportunistic yeasts can colonize sites where radiation is applied, such as oral mucosa and therefore could be a start of local or systemic infections in weakness host [5–8]. Patients with head and neck cancer submitted to radiotherapy can display a higher risk of oral colonization by *Candida* species [9–11]. However, there are few studies addressing the impact of γ -radiation directly on yeast cells. A pioneering study reported that yeasts after irradiation exhibited changes in cell morphology, sensibility to antifungal agents and increased pathogenicity [12], but this was made with a unique application of irradiation. A further study carried out with five *Candida* species submitted to irradiation identified morphological and physiological changes such as accelerated growth and greater production of pseudohyphae and blastospores, which are structures involved in the tissue-invasion process [4]. Other *in vitro* studies have also demonstrated important alterations after irradiation [13], surprisingly fungal species do not die due to irradiation effects, on the contrary, they resist and suffer changes such as faster growth and increased biomass – shown in melanized and nonmelanized fungi after exposure to γ -radiation [14–16]. It is possible that these changes would impact on the yeast virulence profile. Morphological changes were also observed in the germ tubes of *Candida albicans* isolated from patients during head and neck tumor radiotherapy. This study demonstrated that yeasts colonizing the buccal mucosa became more virulent during the course of radiotherapy treatment [17]. Recent study with clinical isolates of *C. albicans* from oral cavity of cancer patients, showed increased production of phospholipase and proteinase after irradiation [18]. Despite these findings, there are no in-depth studies about the possible changes in *Candida non-albican* (CNA) strains, exposed to γ -radiation. This information is relevant, as CNAs have globally increased in cancer patients.

In particular, *Candida tropicalis* is the most common cause of candidemia among CNAs, and is one of the most frequent fungal infections in intensive care units and internal medicine wards [19], usually associated with high mortality rates. In Latin America countries, *C. tropicalis* is the third most common *Candida* species isolated from candidemia [20]. In Brazil, this species was responsible for 18% of all candidemia episodes according to a multicenter surveillance study of 16 hospitals distributed across 5 regions of the country [21].

Studies have indicated that during the radiotherapy process there is increase of colonization by *C. tropicalis* [9,22]; but little is known about the reasons for this increase. In order to investigate if the radiation used in radiotherapy can cause changes in the virulence potential of *C. tropicalis*, we conducted experiments with a suspension of *C. tropicalis* cells, which was submitted to equal doses of the radiation used in the radiotherapy of a patient with head and neck cancer.

Materials & methods

Ethics statement

The experimental protocol was approved by the Ethics Committee in Animal Experimentation of the State University of Maringá (CEEUA 110/2013).

Microorganisms

C. tropicalis ATCC 750 was used throughout this study. A preinoculum was prepared from a pure colony on Sabouraud Dextrose Agar (SDA; Difco, MD, USA) following growth in Sabouraud Dextrose Broth (SDB; Difco) and the yeast cells were diluted to 2.0×10^3 colony-forming units (CFU).

Irradiation

Tubes containing the yeasts suspension in SDB were exposed *in vitro* to telecobalt irradiation. The experimental conditions such as the calculated area and parameters of irradiation were the same as employed in cancer radiotherapy. The conditions were: an irradiation area of 9×26 cm to the right and left sides, with a spine angle of 270 and 90°, respectively. The source-to-skin distance was 80 cm; the target depth was 6.5 cm; and the percentage depth dose was 72.8%. The irradiation rate was 50.93 cGy/min, lasting 2 m 25 s in 40 applications of 180 cGy/day for each side, totaling 7200 cGy. The yeast was therefore irradiated once a day, 5 days a week under a daily fraction dose of 180 cGy for 8 weeks. Every Friday the suspension of irradiated yeasts during the week was evaluated regarding its purity by culture on chromogenic medium, next, the suspension on liquid SDB was centrifuged, and the supernatant

discarded, following it was replaced with equal volume of fresh SDB. The control group corresponded to the nonirradiated yeasts maintained under the same storage and transport conditions used for the irradiated yeasts.

Morphological analysis

After irradiation, *C. tropicalis* was subcultured into yeast extract peptone dextrose broth (YPD; HiMedia Laboratories, Mumbai, India) at 35°C overnight. Afterwards, the cells were diluted and plated on YPD agar at a concentration of approximately 2×10^2 CFU/ml incubated at 25°C for 96 h. 5000 colonies of each sample (irradiated and nonirradiated) were counted. The colonies were examined by portable USB digital microscope 20–800×. The irreversibility was observed after ten subculture of colony type and analyses by the same USB digital microscope [23].

Scanning electron microscopy

The colonies from morphological analysis were used for scanning electron microscopy (SEM), in accordance with França *et al.* [24]. Critical-point drying was performed using a Balzers CPD-010 (Balzers Instruments, Balzers, Liechtenstein) with carbonic gas. Metallization in gold was performed using a Balzers SCD-030 (Balzers Instruments). The samples were then observed with a Shimadzu SS-550 Super scan device (Shimadzu, Tokyo, Japan) at Complexo de Centrais de Apoio à Pesquisa (COMCAP), Universidade Estadual de Maringá.

Filamentation assay

The assays were performed in accordance with Dambrosio *et al.* [17] described on topic frequency and size of *C. albicans* germ tubes. The experiment was performed in triplicate and repeated weekly by three-times.

Growth curve

C. tropicalis were inoculated in YPD broth at 35°C overnight. The growth curve assay was based on Klepser *et al.* [25], with some modifications: cells from the overnight culture were diluted at 1×10^4 yeasts in 30 ml of YPD broth and incubated at 37°C with shaking at 130 r.p.m. Absorbance was measured at 600 nm every hour and a volume of 1 µl of each test suspension was plated on SDA. Colony counting was performed 24 h later. The experiment was independently carried out on two occasions.

Antifungal susceptibility test

The minimal inhibitory concentration (MIC) of each antifungal was determined by broth microdilution according to documents M27-A2 and M27-A3 of the Clinical and Laboratory Standards Institute [26,27]. The antifungal drugs used in this study were amphotericin B (Bristol-Myers Squibb, OH, USA), fluconazole (Pfizer Ireland Pharmaceuticals, Dublin, Ireland), and micafungin (Astellas, Takaoka, Japan). The readings were made on an Expert plus – ASYS® (Biochrom Ltd – Cambridge, UK) microplate reader at 450 nm. MIC was determined as the lowest concentration capable of inhibiting 50% of fungal growth in the presence of azoles and micafungin, and any visible growth in the presence of amphotericin B. The experiments were performed in triplicate. *Candida parapsilosis* ATCC 22019 was employed as control.

Viability test against fluconazole

C. tropicalis were cultivated in YPD broth at 35°C overnight, cells from this culture were counted in Neubauer chamber (Sigma-Aldrich, MO, USA) and next adjusted for a final volume to contain 2×10^4 yeasts in 25 ml of RPMI 1640 (Merck KGaA, Darmstadt, Germany) medium with MOPS buffer (3-(N-morpholino)propanesulfonic acid). Fluconazole at 500 µg/ml was added and the samples were incubated under shaking at 130 r.p.m., at 37°C. Each 2 h 10 µl of sample were mixed in 10 µl Trypan Blue, counted by microscopy to verify the cell yeast viability. The experiment was performed in triplicate and repeated weekly by three-times.

Adherence & biofilm on abiotic surface

The adhesion and biofilm assays were performed in accordance with Porman *et al.* [28] with some modifications. After growth in YPD broth and incubation at 35°C overnight, cells were inoculated in YPD agar for 24 h. Colonies were resuspended in 3 ml of YPD broth and incubated at 35°C overnight. The inoculum was adjusted to 2 OD₆₀₀ in YPD broth and 1 ml was transferred to each well in 24-well polystyrene microplates. Subsequently, the plates were incubated at 25°C for 2 and 24 h without shaking for adhesion and biofilm assay, respectively. The assays

were performed in three plates simultaneously and three methods were applied to evaluate adhesion and biofilm formation ability, with one plate for each method as described below. The experiment was performed on three independent occasions weekly.

Evaluation of biomass density

Each well was washed three-times with 1 ml of sterile water. The plates were scraped to remove the adherent cells, then the cells were resuspended in 1 ml of sterile water and optical density was determined by spectrometry at 600 nm.

Analysis by number of CFU: following scraping as described above, 10 µl of yeast suspension was plated on SDA and the culture was incubated for 24 h at 35°C. Colony counting and statistical tests were performed.

Biofilm analysis by crystal violet staining

After three washes, the plates were drying at room temperature for 45 min. After that, each well was stained with crystal violet aqueous solution (0.4%), for 45 min. Each well was washed three-times with 1 ml of sterile water, and bleached with 700 µl of 95% ethanol. Following 100 µl of each well was transferred to a 96-well plate. Optical density was determined by spectrometry at 595 nm. The experiment was performed in triplicate and repeated weekly by three-times.

Biofilm analysis by XTT

The kinetics of biofilm formation was determined by the tetrazolium salt reduction 2,3-Bis(2-Methoxy-4-Nitro-5-Sulphophenyl)-5-[(Phenylamino)Carbonyl]-2H-Tetrazolium Hydroxide (XTT) colorimetric method. After two washes of the biofilm formed on the plate, 100 µl saline solution, 0.85% saline with 100 µg/ml XTT and 10 µg/ml phenazine methosulphate (PMS) were added to each well and incubated at 35°C for 3 h protected from light. The absorbance of the colorimetric product was determined in 492 nm. The experiment was performed in triplicate and repeated weekly by three-times.

Hydrophobicity

Cellular surface hydrophobicity (CSH) assay was performed in accordance with Gibbons and Etherden [29] with some modifications. Colonies from YPD agar were resuspended in phosphate urea magnesium sulfate (PUM) buffer and homogenized by gentle agitation. The cells were washed three-times, followed by centrifugation. They were then resuspended in 3 ml of PUM buffer, transferred to glass test tubes and homogenized. Optical density was measured at 530 nm in a spectrophotometer. After the addition of 400 µl of n-hexadecane, the cells were incubated in a water bath at 30°C for 10 min. Tubes were mixed in a vortex for two 30-s periods with a 5-s interval. Following phase separation, the optical density was measured at 530 nm. Samples were considered hydrophobic when >50% of cells migrated from the PUM buffer toward the n-hexadecane. The experiment was repeated weekly on three independent occasions.

Adherence to buccal epithelial cells

Adherence assays were performed in accordance with Queiroz *et al.* [30] with some modifications. The yeasts were incubated in RPMI 1640 supplemented with fetal bovine serum at 37°C under agitation at 120 r.p.m. for 60 min to stimulate filamentation. Buccal epithelial cells (BEC) were obtained from healthy volunteers by scraping the buccal mucosa with a sterile spatula and dispensed in 10 ml of PBS (pH 7.2). Following washing procedures, the cells were stained by gram and analyzed under a microscope to rule out the possible presence of yeast from the originally colonization of BEC. All the volunteers were free of oral lesions, and none received antibiotics or antifungals. BEC at a concentration of 2×10^5 cell/ml were incubated with the yeast at a concentration of 2×10^6 yeast/ml in RPMI for 30 min. The cells were washed five-times with PBS and the suspension was placed on glass microscope slides and dried at room temperature. Gram staining was then performed, and the percentage of BEC with yeast adhered and number of yeasts per BEC were evaluated. A total of 100 cells were considered and the experiment was conducted on three independent assays weekly.

Adherence to TR146 cell

Human buccal epithelium TR146 corresponds to a tumor cell described by Rupniak *et al.* [31], from a buccal mucosa that had infiltrated a lymph node. These cells were obtained from Sigma-Aldrich Corporation (10032305).

The adherence assay was performed according to Murciano *et al.* [32]. Briefly, the TR146 cells were cultivated under 5% CO₂ in 24-well microplates following the culture instructions for these cells. After obtaining the monolayer, *C. tropicalis* before and after irradiation was added to the wells, which were then incubated for 90 min at 37°C. After incubation, the nonadherent yeast were removed by aspiration and the plates were scraped to release the adhered yeasts, which were inoculated in plates containing SDA and incubated at 30°C for 24 h. The resulting colonies were then counted. The experiments were carried out in triplicate and repeated on three different weeks.

***In vitro* phagocytosis evaluation**

Peritoneal phagocytes were isolated from female Swiss mice in accordance with Ray and Dittel [33]. Briefly, after counting, approximately 5×10^5 cells were transferred to a 24-well polystyrene culture plate and incubated at 37°C with an atmosphere of 5% CO₂ for 2 h. The cells were washed once, resuspended in 500 µl of RPMI, and incubated at 37°C with 5% CO₂ for 24 h. Each well was washed with PBS to remove the nonadhered cells, and 500 µl of new RPMI was added. *C. tropicalis* were inoculated into YPD broth and incubated at 35°C overnight. Cells from the overnight culture were washed and diluted to a ratio of approximately five yeast cells for one phagocyte cell in 500 µl of RPMI and added to each well. The culture plate was incubated for 2 h at 35°C. Each well was washed with PBS once and panoptic staining was used to stain each cover slip. After drying overnight, the cover slip was fixed and the phagocytosis index (PI) was measured by light microscopy. The phagocytic activity was determined by counting the number of yeasts ingested per phagocyte. A total of 200 phagocytes were counted per cover slip. Experiments were performed in duplicate on two independent assays weekly.

Animal infection

Systemic infection was performed in accordance with Xie *et al.* [34] with modifications. 8-week-old male and female Swiss mice were intravenously injected with 100 µl of PBS containing approximately 1×10^7 CFU/ml via the tail vein. Two male and two female mice were used per culture (before and after irradiation). The mice were sacrificed on the third day post infection. The right kidneys were collected, weighed and macerated with 2 ml of PBS, which was diluted at 1:100 and plated on SDA and incubated at 35°C for 24 h. Some colonies of each sample were plated again on selective media to guarantee purity then frozen at -80°C in SDB with the addition of 50% of glycerol. The experiments were performed on two independent weekly assays.

Histological study of mice kidneys

Immediately after the sacrifice of the animals, the left kidney was collected and preserved in paraformaldehyde 4% for 24 h for fixation. After fixing, the kidneys were maintained in 70°C alcohol, which was changed every 24 h, and processed for inclusion in paraffin. Semiserial sections of 5-mm thickness were made. The slides were stained with hematoxylin and eosin for histopathological evaluation, and with silver according to Grocott–Gomori and counter stained with light green to locate the yeast in the kidney. The slides were observed and photographed in a Nikon Eclipse 80i trinocular light microscope with a Nikon camera (DS Fi1C; Nikon Instruments Inc Melville, NY, USA), coupled to a computer using the Nis-Element software package (Tokyo, Japan).

Representational difference analysis

Irradiated and nonirradiated *C. tropicalis* yeast was recovered from the kidneys of infected animals and submitted to representational difference analysis (RDA). Cells were previously grown in YPD broth at 35°C overnight and posteriorly in YPD agar for 24 h. The inoculum was adjusted to approximately 0.1 OD₆₀₀ in YPD broth and incubated at 35°C for 6 h. Total RNA was extracted using an RNeasy Mini kit (Qiagen, Germany) according to the manufacturer's instructions. Contaminating DNA was removed with RNase-free DNase I (Invitrogen – Thermo Fisher Scientific, MA, USA). The RNA concentration and purity ratio were determined by UV absorbance at 260 and 280 nm using a nano-spectrophotometer Nanok (Kasvi – São José do Pinhais, PR, Brazil). Total RNA integrity was analyzed by electrophoresis in 1.4% agarose gel. The first- and second-strand cDNAs were synthesized in accordance with Siqueira *et al.* [35]. cDNA–RDA was performed according to Gress *et al.* [36] with some modifications in rounds of subtractions and amplifications as follow: cDNA was digested with *Sau3AI* restriction enzyme; the products of PCR amplification were purified using a PCR Purification Kit (Qiagen); one tester-driver hybridizations were performed and reverse cDNA–RDA experiments were carried out as a control, which was submitted to the same conditions. After purification, the final RDA products were cloned into the pGem-T-Easy Vector (Promega, WI, USA), and transformed into *Escherichia coli* DH5α competent cells. Plasmidial DNA was

extracted with a PureLink™ Pro Quick96 Plasmid Purification Kit (Invitrogen) according to the manufacturer's instructions.

Samples were then sequenced using the AB 3500 Genetic Analyzer (Applied Biosystems; Thermo Fisher Scientific) automatic sequencer. Plasmidial DNA (125 ng) was labeled with 2.5 pmol of the primer (M13 Forward 5' GTTTTCCCAGTC 3' and M13 Reverse 5' CAGGAAACAGCTATGACC 3') and 0.5 µl of BigDye Terminator v3.1 Cycle Sequencing Standard (Applied Biosystems) in a final volume of 10 µl. Labeling reactions were performed in an LGC XP Cyclor thermocycler (LGC, Teddington, Middlesex, UK) with an initial denaturation step of 96°C for 3 min followed by 25 cycles of 96°C for 10 s, 55°C for 5 s and 60°C for 4 min. Labeled samples were purified by isopropanol precipitation 75% followed by rinsing with ethanol 60%. Precipitated products were suspended in 10 µl formamide Hi-Fi (Applied Biosystems), denatured at 95°C for 5 min, ice cooled for 5 min and electro injected in the automatic sequencer. Sequencing data were collected using the Data Collection 2 (Applied Biosystems) software package programmed with the following parameters: Dye Set 'Z'; Mobility File 'KB_3500_POP7_BDTv3.mob'; BioLIMS Project '3500_Project1'; Run Module 1 'FastSeq50_POP7_50cm_cfv_100'; and Analysis Module 1 'BC-3500SR_Seq_FASTA.saz'. The resulting sequences were checked by the Phred program module [37] and were assembled to obtain the final set of contigs and singlets using the CAP3 program, then compared with the GenBank [38] nonredundant (nr) database from the National Center for Biotechnology Information using the BLASTX algorithm [39,40]. Identification and protein function was analyzed in UniProt [41] and InterPro [42]. A search for functional categories was performed using the Blast2GO program [43].

Statistical analysis

Statistical analysis was performed using Prism 6.0 software (GraphPad, CA, USA) and significance was determined using the Mann–Whitney pair-wise test, unpaired t-test, the Wilcoxon nonparametric test or ANOVA datasets. The statistical significances were considered when values of $p < 0.05$.

Results

Morphological analysis

Morphological differences were observed between colonies before and after irradiation. Nearly 5000 colonies of each sample were counted and analyzed by digital stereoscopic microscope. Before irradiation, the yeast showed three phenotypic morphologies (Figure 1), described as ring (96.84%); flocculate (2.40%) and wrinkled (0.76%). After irradiation more variation among the phenotypic morphologies was observed (Figure 2): the ring type has decreased, but it continued predominant (91.49%), while the other morphologies were grouped (7.28%), wrinkled (0.92%) and smooth (0.31%). In the smooth type a noticeable loss of aerial mycelia was observed, resulting in colonies presenting new morphological aspects.

Reversibility test, after irradiation, showed colonies that lost the aerial mycelia did not return to the normal aspect; thus all smooth colonies retained their smooth shape after ten subcultures.

C. tropicalis colonies before and after irradiation were analyzed by SEM, around 50 fields, each one was observed. Nonirradiated yeasts presented few filamentation and abundant blastoconidia (Figure 3B). The contrary occurred after irradiation, that is, a decrease in the blastoconidia and an increase of the filamentation forms (Figure 3D). Besides, the smooth colonies presented only blastoconidia forms (Figure 3F).

Filamentous growth

After irradiation, *C. tropicalis* were more effective at forming hyphae *in vitro* (Figure 4A), as well as the size of the filaments was bigger (Figure 4B).

Growth curve

The *C. tropicalis* growth curves were compared before and after irradiation using both complex (rich) and chemically defined (poor) media, YPD and RPMI, respectively (Figure 5A & B). The yeasts presented a higher growth rate after irradiation, and this behavior was more evident in the first 8 h of culture, independent of the culture medium.

Antifungal susceptibility & viability test

There was no yeast growth at any concentration of amphotericin B. The MIC of fluconazole observed before and after irradiation was 1 µg/ml, while for micafungin it was 0.03 µg/ml before irradiation and 0.06 µg/ml after irradiation. Additionally, fluconazole was used to verify the effect on the time kill curve of the yeasts. The results

Figure 1. Different morphologies observed by digital stereoscopic microscope in *Candida tropicalis* ATCC 750 before γ -irradiation.

Colonies grown on yeast extract peptone dextrose medium agar for 96 h at 25°C; about 5000 colonies of each sample (before and after irradiation) were observed by digital stereoscopic microscope. Three phenotypes were detected in yeast before irradiation, which appeared at the following frequencies: **(A)** rings (96.84%); **(B)** flocculate (2.40%) and **(C)** wrinkled (0.76%). All morphologies before irradiation presented whole aerial mycelia at the peripheral margin of the colonies.

indicated that after irradiation, the yeasts survived at fluconazole presence more effectively than before irradiation (Figure 6).

Adherence to abiotic surface & hydrophobicity

A decrease in the capacity of the yeast to form biomass after irradiation was observed (Figure 7A). On the other hand, the optical density (OD_{600}) and determination of CFU revealed no significant difference ($p > 0.05$), as can be seen in Figure 7B and C. Despite statistical differences not being verified, it is possible these results indicate some changes in extracellular matrix production, a feature which may be important for biofilm formation. Regarding

Figure 2. Morphologies observed by digital stereoscopic microscope in *Candida tropicalis* ATCC 750 after γ -irradiation.

The yeasts were irradiated at a rate of 50.93 cGy/min for 2 min 25 s at a 40-fraction dose of 180 cGy/day for each side, totaling 7200 cGy. The cultures were therefore irradiated once a day, five-times a week, for 8 weeks. After irradiation, most yeast morphologies retained (A) ring (91.49%), while on the other hand the minor morphology frequency showed apparent disorganization in the morphology of the colony, generating a variety of morphologies: (B) others (7.28%). Few colonies presented noticeable loss of aerial mycelia: (C) smooth (0.31%) (D) wrinkled (0.92%).

CSH, no statistical difference was observed, indicating that the irradiation probably does not induce changes in the electric charge of the cell wall (Figure 7D).

Biofilm & XTT reduction

Regarding biofilm density (OD₆₀₀) and CFU determination, there was no difference after 24 h (Figure 8A and B). However, the violet crystal method revealed some significant differences in the samples before and after irradiation ($p = 0.0079$), indicating a reduction in the capacity of the irradiated yeasts to form biomass (Figure 8C). The results of XTT reduction assay revealed the same pattern in the irradiated and nonirradiated yeasts (Figure 8D).

Adherence to BEC

C. tropicalis was more adherent to BEC after irradiation (mean = 2.243) than before (mean = 1.767). This difference was significant ($p \leq 0.0001$), with a high mean number of adhered yeasts per cell (Figure 9).

Adherence to TR146 cells

The CFU number of *C. tropicalis* that adhered to TR146 cells decreased from 1×10^5 CFU (before irradiation) to $3\text{--}4 \times 10^4$ after irradiation (Figure 10); this difference was statistically significant ($p = 0.0281$).

In vitro phagocytosis

Irradiated yeast cells were more resistant to murine peritoneal phagocytes activity; the PI was significantly lower after irradiation ($p = 0.0045$), PI = 173.9 (before), and PI = 124.2 after irradiation (Figure 11).

Figure 3. *Candida tropicalis* ATCC 750 ultrastructure of colonies before (A & B) and after (C & D) γ -irradiation, and without aerial mycelium (E & F).

After 7200 cGy of γ -irradiation, the yeasts were grown in yeast extract peptone dextrose agar medium for 96 h. The colonies were fixed in glutaraldehyde, frozen in liquid nitrogen and lyophilized. The ultrastructure was observed with a scanning electron microscope at 500 $\times$ (A, C & E) and 1500 $\times$ (B, D & F). Irradiation changed the organization of the yeasts into colonies. Before irradiation, the colony appeared dense and bushy (A) and many yeast forms and little filamentation (B). After irradiation, the colony was more widely spread (C) with an evident increase in filamentation compared with before the irradiation (D). The smooth colonies were almost wholly yeast forms, with very little filamentation (E).

Animal infection

The yeasts were effective at causing systemic infection in mice as before as after irradiation; however, the irradiated yeasts were significantly more aggressive than the nonirradiated yeasts ($p = 0.0168$), as shown in Figure 12A. This was confirmed by histopathology (Figure 12C), where a higher number of irradiated than nonirradiated yeasts can be seen in the renal tissue (Figure 12B). Additionally, the yeasts were more capable of developing hyphae after radiation (Figure 12D).

The yeasts were able to reach the mice kidneys both before and after irradiation, evidenced by changes that suggest an inflammatory response. In addition, it is clear that the yeasts were more aggressive after irradiation,

Figure 4. *Candida tropicalis* ATCC 750: filamentation on RPMI-1640 medium supplemented with fetal bovine serum 10% for 3 h at 37°C.

(A) After irradiation, the filamentation of the yeasts was more effective, and the difference was statistically significant using ANOVA test ($p = 0.0003$). (B) The pseudohyphae were longer after irradiation than before, and this difference was statistically significant using unpaired t test ($p \leq 0.0001$). Data from three independent replicated experiments were analyzed.

Figure 5. *Candida tropicalis* ATCC 750: growth curve in liquid medium during 12 h under shaking at 37°C.

(A) Yeast before irradiation in comparison with the same strain after irradiation, in RPMI medium. The samples after irradiation had a faster growth until 8 h of incubation. The same behavior was observed in the yeast extract peptone dextrose medium, with more effective growth after irradiation (B).

CFU: Colony-forming unit; YPD: Yeast extract peptone dextrose broth.

disrupting the renal tissue, causing hyperemia and increasing the influx of phagocytic cells (Figure 13C–F), while such changes were subtler in the kidneys of mice infected with nonirradiated yeast (Figure 13B).

Additionally, animals infected with irradiated yeast were more severe on symptomatic clinical aspect than those infected with nonirradiated yeasts. The principal signals or symptoms were lethargy, loss appetite and prostration. There was no difference in the infection rate of males and females in terms of clinical, microbiological or histopathological parameters (data not shown).

RDA

A total of 15 bacterial clones containing the cDNA from the RDA subtraction assay were considered. From these, seven cDNA sequences were generated, five genes sequences were not expressed by yeasts after irradiation, and

Figure 6. Viability test of *Candida tropicalis* ATCC 750 before and after irradiation, both exposed to fluconazole at 500 $\mu\text{g/ml}$, stained by Trypan Blue and counted at microscopy. After irradiation, the yeast was more effective to survive than before irradiation. Data were analyzed from three independent replicated experiments to compose the graph. CFU: Colony-forming unit.

Figure 7. Evaluation of adhesion and cell surface hydrophobicity of *Candida tropicalis* ATCC 750 before and after irradiation.

Crystal violet staining showed a significant decrease in adhesion after irradiation ($p = 0.0087$) (A). There was no significant difference in adhesion measured by OD 600 nm (B) or colony-forming units (C). The cell surface hydrophobicity assay did not exhibit a statistical difference (D). Standard deviation is indicated with error bars. Data from three independent replicated experiments were analyzed using Mann–Whitney test. CFU: Colony-forming unit; OD: Optical density.

another two genes sequences were expressed only after irradiation (Figure 14). Based on the yeast expression profiles revealed by RDA assay, it could be inferred that the response of the yeast during the process of infection was different before and after irradiation, demonstrating that the γ -radiation may interfere with protein expression during systemic infection.

Discussion

The present study investigated the effect of γ -radiation on *C. tropicalis*, without any interference of the host immune system, medication or other external factors. It was demonstrated for the first time that *C. tropicalis*

Figure 8. Biofilm formation of *Candida tropicalis* ATCC 750 and 2,3-bis(2-methoxy-4-nitro-5-sulfophenyl)-5-[(phenylamino)carbonyl]-2H-tetrazolium hydroxide reduction assay.

The yeasts were incubated in yeast extract peptone dextrose at 25°C for 24 h without shaking. The difference in the number of colony-forming units adherent cells of yeasts before and after irradiation was not significant (A & B). Crystal violet staining (C) showed a significant difference $p = 0.0079$. There was no significant difference in 2,3-bis(2-methoxy-4-nitro-5-sulfophenyl)-5-[(phenylamino)carbonyl]-2H-tetrazolium hydroxide assay (D), showing that the irradiation did not affect the metabolism of the yeast. Data from three independent replicated experiments were analyzed using Mann–Whitney test. Standard deviation is indicated with error bars. CFU: Colony-forming unit; OD: Optical density.

Figure 9. Adherence on buccal epithelial cells by *Candida tropicalis* ATCC 750 before and after irradiation.

The yeast were incubated in RPMI supplemented with fetal bovine serum at 37°C for 1 h and shaken at 120 r.p.m. Cells were counted under light microscope after the incubation period. After irradiation, the adherence was more effective, and the difference was statistically significant $p \geq 0.0001$. Data from three independent replicated experiments were analyzed using unpaired t test.

Figure 10. Adherence on TR146 cells by *Candida tropicalis* ATCC 750.

The yeasts were incubated in RPMI supplemented with fetal bovine serum at 37°C for 1 h and shaken at 120 r.p.m. The counting of adhered yeast was performed by colony-forming units counts. Data from three independent replicated experiments were analyzed using unpaired t test. After irradiation, the adherence was less effective, and the difference was statistically significant. CFU: Colony-forming unit.

Figure 11. Phagocytosis assay of *Candida tropicalis* ATCC 750 before and after irradiation.

After irradiation *C. tropicalis* ATCC 750 was much more resistant to *in vitro* phagocytosis by mice peritoneal macrophages. Data from two independent replicated experiments were analyzed using Mann–Whitney test and the difference was significant ($p = 0.0045$).

colony morphology was irreversibly affected by radiation. Moreover, the yeast-cell multiplication capacity was also significantly accelerated. The ability to infect animals and to escape phagocytosis was highly improved after irradiation ($p = 0.0045$). Interestingly, though the radiation dose used (7200 cGy) is cidal for mammalian cancer cells, it did not kill *C. tropicalis*. Instead yeast becomes more virulent than before irradiation. Recently, it has been shown that low doses of γ -radiation are able to induce important changes in fungi molecules, such as melanin, increasing protein synthesis and ensuring more resistance to environmental conditions [15,16]. But in cancer treatment patients the mechanistic details remain to be discovered.

In current study the first observed change was the colony morphologies of the yeasts, with visible disorganization of the macroscopic aspects. After irradiation, the frequency of the ring type morphological colony was decreased due to the increase of the other morphological types (Figures 1 & 2). Moralez *et al.* [23] demonstrated that this morphological diversity is common in colonies of yeasts obtained from clinical isolates. It is well known that some factors such as antifungal [23,44–45] as well as exposure and UV radiation can cause changes in yeast colony morphology [46]. Thus it is possible that these morphological changes were produced in response of the *C. tropicalis* to physical aggression by γ -irradiation, and a reflection of the adaptation process, equally was observed in *C. albicans* [12].

The morphology reversibility test revealed that all colonies that maintained the aerial mycelium returned to the ring morphology (parental type) after two or three subcultures. However, the loss of the aerial mycelium was irreversible (Figure 2C), irrespective of whether it was totally or partially lost. The smooth morphology, after ten subcultures, did not return to the parental morphology. Despite few colonies undergoing permanent changes, it was clear that many factors may be involved in such modifications, like the expression of genes in relation to filamentation forms, which are of great importance for other virulence factors [23,24].

After irradiation the yeasts improved their filamentation capacity when yeasts were grown in liquid medium (Figure 4), like has been reported by other authors [4,47]. According to Figure 3C, we can to infer that yeasts colonies with aerial mycelium have preserved the filamentation capacity, but apparently occurred a change regarding hyphae organization, and it would affect the appearance morphological.

Figure 12. Murine experimental candidemia by evaluation of yeast in the kidneys of animals infected with *Candida tropicalis* ATCC 750 before and after irradiation.

Four mice per group were infected intravenously with *C. tropicalis* ATCC 750 (1×10^7 colony-forming units/ml) before irradiation and after irradiation. The mice were sacrificed on day 3 after infection. The number of yeasts recovered from the kidneys of animals infected with irradiated yeast was much higher than from animals infected with yeast that had not been irradiated. (A) Data from these experiments were analyzed using Mann–Whitney test and the difference was significant. The kidney sections, stained by Grocott–Gomori showed yeasts that before irradiation were able to weakly reach the kidneys (B), although this was the only yeast profile found in >50 sections examined. On the other hand, kidneys of mice infected with irradiated yeasts showed yeasts in all microscope fields and in great numbers, as in (C), and also exhibited filamentous forms (D). CFU: Colony-forming unit.

The yeasts became more activated after irradiation, since the number of CFU in the growth curve was significantly increased. In both YPD and RPMI culture media (complex and defined, respectively), the irradiated yeasts exhibited the same behavior (Figure 5A & B). This observation indicates that the radiation affected the rate of growth of *C. tropicalis*, probably due to metabolic changes, regardless of the culture medium. Similarly growth increase was also observed in *C. neoformans* as well as in others melanized fungi, exposed to ionizing γ -radiation [16].

Although not statistically significant, it was observed that irradiated yeasts survived more effectively in presence of fluconazole (Figure 6). We can speculate that since the growing rate of the yeasts was accelerated, they became more able to resist antifungal action.

C. tropicalis has a high biofilm-formation capacity, with great biomass production ability, which is one of the most important virulence features of this species [48–50]. In this study with *C. tropicalis*, it was found that this characteristic appears not to be affected by irradiation, as there were no important differences in adhesion on abiotic surfaces, CSH and biofilm formation, which were evaluated by different methods (Figures 7 & 8). The exception was biomass, which was evaluated by violet crystal staining, and showed a significant decrease in both adhesion and biofilm formation. These data suggest that the yeasts reduced the production of the extracellular matrix after radiotherapy, despite probably preserving metabolism process and the cell-wall structures.

Figure 13. Histological analysis of kidney sections from animals infected with *Candida tropicalis* ATCC 750 before and after irradiation, stained by hematoxylin and eosin.

Representative histology (from >50 examined in each situation): kidneys of not infected mice (A) and mice infected intravenously with 1×10^7 colony-forming units/ml nonirradiated (B); and after irradiation (C–F). In (A), we observe the normal kidney morphology, highlighting: glomeruli (g), Bowman's capsule (*) and tubules (t). In (B), observe hyperemic vessels (arrowhead) of the glomerulus and among the tubules, also reduction of Bowman's space. In the animals infected with irradiated yeasts, observe the occurrence of inflammatory infiltrate (i), groups of yeasts (y), tubules with acidophilic cells (arrow) and areas composed of cell debris (d). All mice were sacrificed on day 3 after infection, and sections were stained with hematoxylin and eosin. Magnification: 50 $\times$ and 100 $\times$.

We decided to perform assays with two types of cell lines to verify whether after irradiation the yeast adheres differently to healthy epithelium and cancerous epithelium, since the focus of the research is the effect of radiation on colonizing yeasts from patients with head and neck cancer under radiotherapy treatment. When adherence of *C. tropicalis* on normal BEC and TR146 cells was compared, apparently contrasting results were found: there was a significant increase in adhesion on BEC, but a decrease in adhesion to TR146 cells (Figures 9 & 10). BEC are fresh cells which seem to easily recognize the yeast, meanwhile cancer cells, which lose adhesion molecules on the cell surface [51], resulting a decrease in the receptors for adhesion of *C. tropicalis* to TR146 cells. Additionally, it is important to highlight the differences between the methods employed, as the protocols used for the yeast evaluation were different for each test: for assay with BEC it was possible to observe and count the yeasts and cells using light microscopy, while in the adhesion assay with TR146 cells, evaluation was performed by colony counting, making it impossible to know how many yeasts formed each colony.

The phagocytosis assay produced remarkable results, showing that the phagocytic index statistically decreased after irradiation (Figure 11). This result shows that irradiated yeasts became more resistant to killing, and may explain the high number of CFU recovered (Figure 12A) and visualized from animal kidneys infected with irradiated

	Gene symbol (Blast n)	Description (Blast2Go)	Length	e-value	Sim means
Before irradiation	CTRG_06049 <i>C. tropicalis</i> Mya3404	Adenosylhomocysteinase	394	4.71E-62	95.8%
	CTRG_00819 <i>C. tropicalis</i> Mya3404	Pgm2p phosphoglucomutase	909	6.78E-72	86.60%
	CTRG_05573 <i>C. tropicalis</i> Mya3404	Protein transport protein SEC61 alpha subunit	342	7.33E-38	92%
	CTRG_05084 <i>C. tropicalis</i> Mya3404	Dihydrolipoylysine-residue acetyltransferase component of pyruvate dehydrogenase complex, mitochondrial precursor	1020	5.46E-57	92.13%
	CTRG_04114 <i>C. tropicalis</i> Mya3404	Similar to potential NADH-dependent flavin oxidoreductase	287	7.38E-27	95.25%
After irradiation	CAGL0G01342g <i>C. glabrata</i> CBS138	Vacuolar segregation 7 (<i>C. Glabrata</i>)	1037	6.84E-47	93.33%
	LBRM_27_1880 <i>leishmania braziliensis</i> MHOM/BR/75/M2904	Tubulin binding cofactor c	974	7.35E-42	80.11%

Figure 14. Genes differentially expressed of *Candida tropicalis* ATCC 750 before and after irradiation recovered from animals infections assay.

Representational difference analysis techniques allow us to infer that five genes were not expressed, whereas two other genes were differentially expressed after irradiation.

yeast (Figure 12C & D), in contrast to the few yeasts found in kidneys of mice infected with nonirradiated yeasts (Figure 12B).

When analyzing histological images and the phagocytosis assay, it is possible to infer that the irradiated yeast became more aggressive and more efficient at causing the infectious process (Figure 13). A no less important finding was that visual clinical analysis of mice revealed an inflammatory process, only in the tail of the mice infected with irradiated yeast. This could reinforce the BEC adherence results, proving that the yeasts became more effective at adhering to cell surfaces.

The RDA technique was used in this study to verify the level of gene expression in the studied yeast. RDA was first developed to identify differences between two sequences of DNA in cancer cells [52], and was then adapted to detect differences in cDNA [53]. It has since been used on comparative gene expression in several areas of health [35,54].

To identify the different genes expressed after infection, the tester representing the cDNA of the yeasts after irradiation and the driver before irradiation were considered first. In a second assay, the tester and driver representing before and after irradiation were analyzed (Figure 14). One of the differentially expressed genes found in irradiated yeast was LBRM.27_1880 encoding Tubulin Binding Cofactor C. This protein is directly involved in the cellular cycle [55]. It is important to remember that the irradiated yeasts exhibited a higher growth rate, which corroborated with the results of the growth curve (Figure 5). Other genes related to the metabolism of the yeast were differentially

expressed. While further studies should be performed in order to elucidate the pathways that changed after irradiation, this was not the objective of the present study.

Conclusion

The present study confirmed some changes in yeasts by γ -radiation similar to a course of radiotherapy. Some of the alterations have already been observed by other authors, but our results added other important alterations which have not yet been reported. The results highlight that radiation treatment can cause changes in the virulence factors of *C. tropicalis*, such as increasing the growth rate, filamentation and adhesion to epithelial cells, and in addition can increase its ability to escape the phagocytosis process and consequently become more efficient at causing systemic infection in mice. These changes can directly affect the relationship between yeast and host, increasing the infection potential; and we strongly believe that these events occur in yeasts that colonize patients undergoing radiotherapy treatment, thereby justifying the increased risk of the death of cancer patients by candidemia. They may, therefore, be one reason for the increase in morbidity and mortality rates due to candidiasis by *C. tropicalis* that affect this patient group. We hope that these findings stimulate interest in the R&D of prophylactic methods to minimize the occurrence of yeast infections in cancer patients.

Future perspective

We observed that γ -radiation induced important changes in *C. tropicalis* increasing its virulence profile, which could directly affect the relationship between yeasts and hosts, and induce an increase in the risk of infection in patients undergoing radiotherapy treatment. More specific studies would be carried out to evaluate the same factors now studied, but on clinical isolates, moreover, it would be interesting to carry out this evaluation on fresh cancer cells, instead of cell lines. Also include other assays like if *C. tropicalis* melanin production is stimulated by radiation, since this molecule is related to resistance, including to phagocytosis and there are reports that this specie can produce melanin.

Summary points

- γ -irradiation induced important changes in morphology and functionality of *Candida tropicalis*.
- Some morphological changes induced by γ -irradiation were irreversible.
- *C. tropicalis* irradiated showed larger pseudohyphae after irradiation.
- After irradiation yeasts become more resistant to fluconazole.
- After irradiation *C. tropicalis* causes disseminated experimental infection more efficiently.
- γ -irradiation significantly increased the experimental murine systemic infection by *C. tropicalis*.

Acknowledgements

The authors would like to thank the Sant'ana Radiation Oncology Center of Maringá for the radiation procedures and physical calculations.

Financial & competing interests disclosure

The authors have received funding from Araucaria Foundation: Protocol 39310.413.29525.18042013 – Covenant number: 272/2014, National Council for Scientific and Technological Development (CNPq – Process number: 477918/2013-1) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), from Health Ministry of Education and Culture for scholarship (EM da Silva and TIE Svidzinski). The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Ethical conduct of research

The experimental protocol was approved by the Ethics Committee in Animal Experimentation of the State University of Maringá (CEEUA 110/2013).

References

Papers of special note have been highlighted as: • of interest; •• of considerable interest

1. Rim CH, Seong J. Application of radiotherapy for hepatocellular carcinoma in current clinical practice guidelines. *Radiat. Oncol. J.* 34(3), 160–167 (2016).
2. Jeong H, Bok S, Hong BJ, Choi HS, Ahn GO. Radiation-induced immune responses: mechanisms and therapeutic perspectives. *Blood Res.* 51(3), 157–163 (2016).
3. Wang Y, Guan H, Xie D-F *et al.* Proteomic analysis implicates dominant alterations of RNA metabolism and the proteasome pathway in the cellular response to carbon-ion irradiation. *PLoS ONE* 11(10), e0163896 (2016).
- **Excellent article to understand changes on cell metabolism in response to carbon-ion irradiation.**
4. Procop GW, Davis HA, Volz PA. Cobalt 60 radiation and growth of *Candida* species. *Mycoses* 31(9), 466–470 (1988).
5. Ahmadi A. Potential prevention: Aloe vera mouthwash may reduce radiation-induced oral mucositis in head and neck cancer patients. *Chin. J. Integr. Med.* 18(8), 635–640 (2012).
6. Sønstevedt T, Johannessen AC, Stühr L. A rat model of radiation injury in the mandibular area. *Radiat. Oncol.* 9(10), 129 (2015).
7. Karbach J, Walter C, Al-Nawas B. Evaluation of saliva flow rates, *Candida* colonization and susceptibility of *Candida* strains after head and neck radiation. *Clin. Oral Investig.* 16(4), 1305–1312 (2012).
8. Barlow ML, Cummings RJ, Pentland AP *et al.* Total body irradiation exacerbates dissemination of cutaneous *Candida albicans*. *Infection Radiat. Res.* 186(5), 436–446 (2016).
9. Ainaur MMAR, Mahesh P, Kumar KV, Deepak A, Mrinal U, Sharada P. An *in vitro* study of isolation of candidal strains in oral squamous cell carcinoma patients undergoing radiation therapy and quantitative analysis of the various strains using CHROMagar. *J. Oral Maxillofac. Pathol.* 21(1), 11–17 (2017).
10. Özçelik B, Citak S, Cesur S, Abbasoglu U, Içli F. *In vitro* susceptibility of *Candida* species isolated from cancer patients to some antifungal agents. *Drug Metabol. Drug Interact.* 20(1–2), 101–108 (2004).
11. Thaweboon S, Thameboon B, Srithavaj T, Choonharuangdej S. Oral colonization of *Candida* species in patients receiving radiotherapy in the head and neck area. *Quintessence Int.* 39(2), 52–57 (2008).
12. Mendling W, Haller I. The effect of therapeutic dose of gamma radiation on *Candida albicans* cells *in vitro*. *Geburtshilfe Frauenheilkd* 37(11), 947–951 (1977).
13. Bennett CB, Lewis LK, Karthikeyan G *et al.* Genes required for ionizing radiation resistance in yeast. *Nat. Genet.* 29(4), 426–434 (2001).
14. Mironenko NV, Alekhina IA, Zhdanova NN, Bular SA. Intraspecific variation in gamma-radiation resistance and genomic structure in the filamentous fungus *Alternaria alternata*: a case study of strains inhabiting Chernobyl reactor no. 4. *Ecotoxicol. Environ. Saf.* 45(2), 177–187 (2000).
15. Dadachova E, Bryan RA, Huang X *et al.* Ionizing radiation changes the electronic properties of melanin and enhances the growth of melanized fungi. *PLoS ONE* 2(5), e457 (2007).
- **Excellent article for understanding how ionizing radiation affects growth of melanized fungi.**
16. Casadevall A, Cordero RJ, Bryan R, Nosanchuk J, Dadachova E. Melanin, radiation, and energy transduction in fungi. *Microbiol. Spectr.* doi:10.1128/microbiolspec.FUNK-0037-2016 (2017) (Epub ahead of print).
17. Dambrosio D, Svidzinski TIE, Svidzinski AE, Dalalio MMO, Moliterno RA. Radiotherapy effect on frequency of *Candida* spp. and on virulence of *C. albicans* isolated from the oral cavity of head and neck cancer patients. *Revista de Ciênc Farm Básicas e Apl.* 30(2), 153–159 (2009).
18. Ramla S, Sharma V, Patel M. Influence of cancer treatment on the *Candida albicans* isolated from the oral cavities of cancer patients. *Support Care Cancer* 24(6), 2429–2436 (2016).
- **Excellent article for understanding how *Candida* suffers changes due to cancer treatment.**
19. Eliakim-Raz N, Babaoğlu R, Yahav D, Yanai S, Shaked H, Bishara J. Epidemiology, microbiology, clinical characteristics, and outcomes of candidemia in internal medicine wards – a retrospective study. *Int. J. Infect. Dis.* 52, 49–54 (2016).
- **Excellent article for understanding the clinical importance of the genus *Candida*.**
20. Nucci M, Queiroz-Telles F, Alvarado-Matute T *et al.* Epidemiology of candidemia in Latin America: a laboratory-based survey. *PLoS ONE* 8(3), e59373 (2013).
- **Excellent article for understanding *Candida* spp. epidemiology in Latin America.**
21. Doi AM, Pignatari ACC, Edmond MB *et al.* Epidemiology and microbiologic characterization of nosocomial candidemia from a Brazilian National Surveillance Program. *PLoS ONE* 11(1), e0146909 (2016).
22. Singh GK, Capoor MR, Nair D, Bhowmik KT. Spectrum of fungal infection in head and neck cancer patients on chemoradiotherapy. *J. Egypt Natl. Canc. Inst.* 29(1), 33–37 (2017).
23. Morales AT, França EJ, Furlaneto-Maia L, Quesada RM, Furlaneto MC. Phenotypic switching in *Candida tropicalis*: association with modification of putative virulence attributes and antifungal drug sensitivity. *Med. Mycol.* 52(1), 106–114 (2014).
- **Excellent article for understanding that the virulence factors are different for each phenotypic switch, even being the same yeast species.**

24. França EJ, Andrade CG, Furlaneto-Maia L *et al.* Ultrastructural architecture of colonies of different morphologies produced by phenotypic switching of a clinical strains of *Candida tropicalis* and biofilm formation by variant phenotypes. *Micron* 42(7), 726–732 (2011).
25. Klepser ME, Wolfe EJ, Jones RN, Nightingale CH, Pfaller MA. Antifungal pharmacodynamics characteristics of fluconazole and amphotericin B tested against *Candida albicans*. *Antimicrob. Agents Chemother.* 41(6), 1392–1395 (1997).
26. NCCLS. Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts: Approved Standard (2nd Edition). NCCLS Document M27-A2. PA, USA (2002). www.researchgate.net/file.PostFileLoader.html?id=55b324fe5e9d97d2638b45b8&assetKey=AS%3A273819233652736%401442295000257
27. Clinical and Laboratory Standards Institute (CLSI). Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeast (3rd Edition). Clinical and Laboratory Standards Institute, PA, USA (2008). https://clsi.org/media/1461/m27a3_sample.pdf
28. Porman AM, Hirakawa MP, Jones SK, Wang N, Bennett RJ. MTL-independent phenotypic switching in *Candida tropicalis* and a dual role for *Wor1* in regulating switching and filamentation. *PLoS Genet.* 9(3), e1003369 (2013).
- **Excellent article for understanding that filamentation is related to the colony morphology.**
29. Gibbons RJ, Etherden I. Comparative hydrophobicities of oral bacteria and their adherence to salivary pellicles. *Infect. Immun.* 41, 11190–11196 (1983).
30. Queiroz PA, Godoy JS, Mendonça P de S, Pedrosa RB, Svidzinski TI, Negri M. Adhesion and biofilm formation in artificial saliva and susceptibility of yeasts isolated from chronic kidney patients undergoing haemodialysis. *J. Med. Microbiol.* 64(9), 960–966 (2015).
31. Rupniak HT, Rowlatt C, Lane EB *et al.* Characteristics of four new human cell lines derived from squamous cell carcinomas of the head and neck. *J. Natl Cancer Inst.* 75(4), 621–635 (1985).
32. Murciano C, Moyes DL, Runglall M *et al.* Evaluation of the role of *Candida albicans* agglutinin-like sequence (Als) proteins in human oral epithelial cell interactions. *PLoS ONE* 7(3), e33362 (2012).
33. Ray A, Dittel BN. Isolation of mouse peritoneal cavity cells. *J. Vis. Exp.* 28(35), pii1488 (2010).
34. Xie J, Tao L, Nobile CJ *et al.* White-opaque switching in natural MTL α/α isolates of *Candida albicans*: evolutionary implications for roles in host adaptation, pathogenesis, and sex. *PLoS Biol.* 11(3), e1001525 (2013).
35. Siqueira VL, Cardoso RF, Caleffi-Ferracioli KR *et al.* Structural changes and differentially expressed genes in *Pseudomonas aeruginosa* exposed to meropenem–ciprofloxacin combination. *Antimicrob. Agents Chemother.* 58(7), 3957–3967 (2014).
36. Gress TM, Wallrapp C, Frohme M *et al.* Identification of genes with specific expression in pancreatic cancer by cDNA representational difference analysis. *Genes Chromosomes Cancer* 19(2), 97–103 (1997).
37. Molecular Biology Laboratory – UNB Department of Cell Biology. www.biomol.unb.br/phph/index.html
38. NCBI. <http://www.ncbi.nlm.nih.gov>
39. Butler G, Rasmussen MD, Lin MF *et al.* Evolution of pathogenicity and sexual reproduction in eight *Candida* genomes. *Nature* 459(7247), 657–662 (2009).
40. GeneBank Database. www.ncbi.nlm.nih.gov
41. UniProt. www.uniprot.org
42. InterPro. <http://www.ebi.ac.uk/interpro/>
43. Blast2GO Program Database. www.blast2go.com
44. Yoon SA, Vazques JA, Steffan PE, Sobel JD, Akins RA. High-frequency, *in vitro* reversible switching of *Candida lusitanae* clinical isolates from amphotericin B susceptibility to resistance. *Antimicrob. Agents Chemother.* 43(4), 836–845 (1999).
45. Deorukhkar SC, Saini S, Mathew S. Virulence factors contributing to pathogenicity of *Candida tropicalis* and its antifungal susceptibility profile. *Int. J. Microbiol.* 2014, 456878 (2014).
46. Morrow B, Anderson J, Wilson J, Soll DR. Bidirectional stimulation of the white-opaque transition of *Candida albicans* by ultraviolet irradiation. *J. Gen. Microbiol.* 135 (5), 1201–1208 (1989).
47. Farrag HA, A-Karam El-Din A, Mohamed El-Sayed ZG, Abdel-Latifissa S, Kamal MM. Microbial colonization of irradiated pathogenic yeast to catheter surfaces: relationship between adherence, cell surface hydrophobicity, biofilm formation and antifungal susceptibility. A scanning electron microscope analysis. *Int. J. Radiat. Biol.* 91(6), 519–527 (2015).
48. Silva-Dias A, Miranda IM, Branco J, Monteiro-Soares M, Pina-Vaz C, Rodrigues AG. Adhesion biofilm formation, cell surface hydrophobicity, and antifungal planktonic susceptibility: relationship among *Candida* spp. *Front. Microbiol.* 12(6), 205 (2015).
- **Excellent article to understand the difference in the virulence factors among the *Candida* species.**
49. Udayalaxmi, Jacob S, D'Souza D. Comparison between virulence factors of *Candida albicans* and non-*albicans* species of *Candida* isolated from genitourinary tract. *J. Clin. Diag. Res.* 8(11), DC15-7 (2014).
50. Tellapragada C, Eshwara VK, Johar R *et al.* Antifungal susceptibility patterns *in vitro* production of virulence factors, and evaluation of diagnostic modalities for the speciation of pathogenic *Candida* from blood stream infections and vulvovaginal candidiasis. *J. Patholog.* 2014, 142864 (2014).

51. Ertel A, Verghese A, Byers SW, Ochs M, Tozeren A. Pathway-specific differences between tumor cell lines and normal and tumor tissue cells. *Mol. Cancer* 5(1), 55 (2006).
52. Lisitsyn N, Lisitsin N, Wigler M. Cloning the differences between two complex genomes. *Science* 259(5097), 946–951 (1993).
53. Hubank M, Schatz DG. Identifying differences in mRNA expression by representational difference analysis of cDNA. *Nucleic Acids Res.* 22(25), 5640–5648 (1994).
54. Do Carmo Silva L, Tamayo Ossa DP, Castro SV *et al.* Transcriptome profile of the response of *Paracoccidioides* spp. to camphene thiosemicarbazide derivative. *PLoS ONE* 10(6), e0130703 (2015).
55. Hage-Sleiman R1, Herveau S, Matera EL, Laurier JF, Dumontet C. Tubulin binding cofactor C (TBCC) suppresses tumor growth and enhances chemosensitivity in human breast cancer cells. *BMC Cancer* 12(10), 135 (2010).

Carvacrol is highly disruptive against coagulase-negative staphylococci in *in vitro* biofilms

Vânia Gaio^{‡,1}, Cláudia Afonso Lima^{‡,1}, Fernando Oliveira¹, Ângela França¹ & Nuno Cerca^{*,1}

¹ CEB-LIBRO – Centre of Biological Engineering, Laboratory of Research in Biofilms Rosário Oliveira, University of Minho, Campus de Gualtar, 4710-057, Braga, Portugal

* Author for correspondence: Tel.: +351 2530 1968; Fax: +351 2550 4400; nunocerca@ceb.uminho.pt

[‡] Authors contributed equally

Aim: To evaluate the antimicrobial activity of carvacrol against stationary-phase planktonic and biofilm cells of coagulase-negative staphylococci and comparison to traditional antistaphylococcal antibiotics. **Materials & methods:** The antimicrobial effect of carvacrol and antibiotics against planktonic and biofilm cells were assessed through quantification of the number of culturable and/or viable cells. Confocal laser scanning microscopy was used to evaluate the effect of carvacrol on the biofilm structure. **Results:** A concentration of 4 mM of carvacrol demonstrated a potent antimicrobial effect, vastly superior than ciprofloxacin, rifampicin, gentamicin and vancomycin, both in planktonic and biofilm cells. **Conclusion:** Carvacrol is a potential antimicrobial agent, which applicability for the prevention and/or treatment of coagulase-negative staphylococci biofilm-associated infections is worthwhile investigating in more detail.

First draft submitted: 21 June 2017; Accepted for publication: 7 August 2017; Published online: 23 November 2017

Keywords: antimicrobial resistance • biofilm • nosocomial infections

The increasing number of morbid, chronically ill and immunocompromised or immunosuppressed patients, as well as the increasing use of medical devices, have led to the acknowledgement of a large variety of infections caused by coagulase-negative staphylococci (CoNS) [1]. *Staphylococcus epidermidis* is, by far, the most commonly found species in CoNS infections. However, other CoNS species such as *Staphylococcus equorum*, *Staphylococcus capitis*, *Staphylococcus haemolyticus*, *Staphylococcus hominis*, and *Staphylococcus warneri* have also been recognized as human pathogens, causing infections with serious health consequences [2–4]. Importantly, due to the emergence of methicillin resistance among CoNS species [1] and their remarkable ability to form biofilms on medical devices [5], the treatment of CoNS infections is usually very challenging. Moreover, CoNS biofilms are known to be more tolerant to antibiotics than their planktonic counterparts [6,7], often resulting in the progression of persistent and recalcitrant infections [8]. Over the last years, several alternatives to antibiotics have been considered for the treatment of these infections [9]. Worldwide, plant-originated compounds, such as essential oils, have been explored and emerged as noble alternative approaches to treat several infections and/or diseases [10–12]. Carvacrol, also known as 2-methyl-5-(1-methylethyl)-phenol, is one of the major components of essential oils of oregano species [13–15] and present a wide diversity of biological activities, such as anti-inflammatory [16], antioxidant [17], antitumor [18] and antimicrobial [19–21]. Moreover, a distinguished advantage of the treatment of infections with essential oils and/or their isolated compounds is the low rate of antimicrobial resistance development when compared with traditional treatment with antibiotics [22]. Carvacrol antimicrobial activity has been mainly attributed to its hydrophobic nature, despite its exact mechanism of action is still poorly understood [23–25]. So far, it is known that its free hydroxyl group interacts with the lipid bilayer of the cytoplasmic membrane causing the expansion and destabilization of the membrane structure, consequently increasing its fluidity and permeability [25–28]. It has been reported that carvacrol is able to interfere with planktonic growth and biofilm formation and/or maturation of clinically relevant pathogens, such as *S. aureus* and *S. epidermidis* [11,29–31]. Nevertheless, nothing is known regarding the susceptibility of other relevant CoNS species to carvacrol. Thus, the main goal of this study was to evaluate the antimicrobial

activity of this essential oil compound against several CoNS strains by evaluating the susceptibility of *S. equorum*, *S. capitis*, *S. haemolyticus*, *S. hominis*, *S. warneri* and also *S. epidermidis* planktonic and biofilm cells to carvacrol. Furthermore, the efficacy of this natural compound was compared with the efficacy of four of the most frequently used antibiotics to treat staphylococcal infections (ciprofloxacin, rifampicin, gentamicin and vancomycin).

Materials & methods

Bacteria & growth conditions

Six CoNS species, selected from a collection of previously characterized strains [32,33], were used in this study. CoNS cultures were obtained by inoculating one single colony into 1 ml of Tryptic Soy Broth (TSB; Liofilchem, Teramo, Italy), and incubating for 24 ($\pm$ 2) h at 37°C with shaking at 200 r.p.m. (20 mm orbit). A total of 24-h-old cultures were diluted 1:200 in 2 ml of fresh TSB, and incubated at 37°C with shaking at 200 r.p.m. (20 mm orbit) for 24 ($\pm$ 2) h, in order to obtain stationary phase planktonic cells. Biofilms were formed as described before [33], with minor modifications. Briefly, in a 96-well microtiter plate (Orange Scientific, Braine-l'Alleud, Belgium), 2 μ l of the 24-h-old cultures were inoculated into 200 μ l of TSB supplemented with 0.25% of glucose (Panreac Química, Barcelona, Spain) to promote biofilm formation. The plates were then incubated for 24 h at 37°C with shaking at 120 r.p.m. (10 mm orbit). Prior to any assay, spent medium was removed and biofilms carefully washed once with 200 μ l of 0.9% NaCl (VWR Chemicals, Leuven, Belgium).

Carvacrol & antibiotics minimum inhibitory concentration determination

The minimum inhibitory concentration (MIC) of carvacrol, ciprofloxacin and rifampicin, gentamicin and vancomycin (Sigma-Aldrich, MO, USA) was determined by using the broth microdilution method [34]. In brief, carvacrol (Sigma-Aldrich) was serially diluted in TSB (1:2) in order to obtain concentrations ranging from 4 to 0.03125 mM. Regarding the antibiotics, twofold dilutions in TSB were performed to obtain a wide range of concentrations, which varied for each antibiotic tested. A total of 200 μ l of each dilution were then placed in each well of a 96-well plate (Orange Scientific). Thereafter, each well was inoculated with 2 μ l of a suspension with approximately 1×10^8 colony-forming units (CFU)/ml of a planktonic culture. Moreover, 200 μ l of media with bacteria (without antibiotics) were used as a positive control for bacteria growth, and a negative control was performed with media without bacterial suspension. The plates were incubated at 37°C during 24 ($\pm$ 2) h without shaking. Bacterial growth inhibition in the presence of carvacrol or antibiotics was evaluated by direct visualization of medium turbidity, and MIC values were defined as the lowest concentrations of antimicrobials that inhibited visible growth of bacteria. This experiment was repeated three independent times, with technical duplicates.

Susceptibility of planktonic cells to carvacrol

In order to assess the effect of carvacrol on CoNS growth, 1×10^8 CFU/ml of stationary phase planktonic cells, grown as described above, were incubated with 4 mM of carvacrol ($2 \times$ MIC) for 5, 15, 45 and 90 min with shaking at 200 r.p.m. (20 mm orbit). A control was obtained before the addition of carvacrol, and the effect of carvacrol was assessed through dilution and plating of planktonic cells, as further explained. This experiment was performed at least three independent times, with technical duplicates. Additionally, viability of planktonic cells was assessed after 15 and 90 min of exposure to carvacrol by flow cytometry (EC800™, Sony Biotech Inc, CA, USA), using SYTO BC/propidium iodide staining [35]. A control with dead cells was obtained by incubating bacteria 15 min on ice in 1 ml of methanol (Thermo Fisher Scientific, Leicestershire, UK).

Biofilms susceptibility to carvacrol

After being washed once with 0.9% NaCl, 200 μ l of TSB with 4 mM of carvacrol were carefully added on top of the 24-h-old biofilms, and the 96-well plates incubated at 37°C, with shaking at 120 r.p.m. (10 mm orbit). After 5 and 90 min of incubation, the medium was removed, biofilms washed once and suspended in 200 μ l of 0.9% NaCl supplemented with 0.05% Tween-20. For each strain, three biofilms were pooled together. The suspension was then sonicated for 3 s at 40% amplitude (Cole Parmer Ultrasonic Processor, IL, USA) to disrupt aggregates without killing the cells [36], prior to dilution and plating (see further section). This assay was repeated at least three independent times.

Biofilms 3D structure analysis after treatment with carvacrol

To evaluate CoNS biofilms structure after interaction with carvacrol, biofilms were analyzed by confocal laser scanning microscopy (CLSM). Biofilms were prepared as above, with some modifications. Briefly, in a 24-well polystyrene plate (Thermo Fisher Scientific, NY, USA), 5 µl of a 24-h-old culture were inoculated into 1 ml of TSB supplemented with 0.25% glucose and incubated for 24 h (± 2), at 37°C with shaking at 120 r.p.m. (10 mm orbit). Thereafter, the biofilm bulk-fluid was discarded; biofilms were washed once with 0.9% NaCl supplemented with 0.05% Tween-20 and 200 µl of TSB with 4 mM of carvacrol were carefully placed on top of the biofilms. The plates were incubated at 37°C with shaking at 120 r.p.m. (10 mm orbit) during 90 min. After the incubation period, biofilms were washed with a 0.9% NaCl solution and the bottom of each well was removed using a previously heated metal heat cutter with the same diameter of the wells. These pieces were placed on the top of a glass slide and stained with SYTO BC (Molecular Probes, Thermo Fisher Scientific), following the manufacturer's instructions. Biofilms were analyzed using an OlympusTM FluoView FV1000 (Olympus, Lisbon, Portugal) confocal scanning laser microscope with a 60 × water-immersion objective (60×/1.2 W). Images were taken with 512 × 512 resolution.

Comparison between carvacrol & antibiotics antimicrobial activities

For planktonic assays, 1×10^8 CFU/ml of a stationary phase planktonic culture were incubated with 4.5 mg/l ciprofloxacin [37], 10 mg/l rifampicin [38], 10 mg/l gentamicin [39] or 40 mg/l vancomycin [38], which correspond to peak serum concentration (PSC) of each antibiotic, up to 90 min at 37°C with shaking at 200 r.p.m. (20 mm orbit). For biofilm assays, 24-h-old biofilms, grown as described above, were washed once with 0.9% NaCl supplemented with 0.05% Tween-20, and 200 µl of TSB with PSC of each antibiotic were carefully added on top of the biofilms. The microtiter plates were incubated at 37°C with shaking at 120 r.p.m. (10 mm orbit) up to 90 min. The growth of both planktonic and biofilm cells was assessed after the incubation period with antibiotics, as previously described for carvacrol assays. A control was obtained before adding antibiotics by plating planktonic or biofilm cells suspended in 0.9% NaCl and sonicated for 3 s and 40% amplitude. Carvacrol (4 mM) was tested at the same time to enable direct comparison with antibiotics efficacy. Sample collection and plating were performed as described above. This assay was repeated at least three independent times.

Dilution & plating of planktonic & biofilm cells for CFUs quantification

Dilution and plating was performed to allow the quantification of culturable cells by CFUs counting. For that purpose, at each time point, an aliquot of 20 µl was collected and transferred into 180 µl of 0.9% NaCl supplemented with 0.05% of Tween-20 (AppliChem, Darmstadt, Germany). Importantly, in biofilm assays the aliquot was obtained from a previously disrupted biofilm suspended in 0.9% NaCl, sonicated for 3 s at 40% amplitude. Thereafter, samples were serially diluted (tenfold dilutions) and at least three different dilutions were plated on Tryptic Soy Agar plates to allow counting of CFUs.

Statistical analysis

Statistical significance between control and carvacrol-, between control and antibiotic- and between carvacrol- and antibiotic-treated samples was determined with unpaired *t*-test (two-tailed) using GraphPad Prism version 6 (Trial version, CA, USA). Differences were considered significant when *p*-values were less than 0.05.

Results

Carvacrol & antibiotics MIC determination

The MIC of carvacrol and of the tested antibiotics that inhibited the growth of the six CoNS strains under study is shown in Table 1. With the exception of *S. epidermidis* SECOM020A.1 resistance to gentamicin, all strains were susceptible to the tested antibiotics, according to EUCAST breakpoints [40]. Furthermore, all CoNS strains were also susceptible to concentrations of carvacrol between 1 and 2 mM.

Carvacrol is highly effective against CoNS planktonic & biofilm cells

In order to further characterize carvacrol antimicrobial activity against CoNS, the susceptibility of stationary phase planktonic cells to carvacrol was evaluated overtime. As can be seen in Figure 1, a significant decrease in the $\log_{10}$ CFU/ml was detected after only 5 min of exposure to carvacrol in all CoNS strains tested: the reduction in culturability ranged from 0.96, for *S. capitis*, to 4.62 $\log_{10}$ CFU/ml for *S. epidermidis*. Furthermore, as soon as

Table 1. Minimum inhibitory concentration of carvacrol and antibiotics determined for the six coagulase-negative staphylococci species under study.

CoNS species/strains	MIC range				
	Carvacrol (mM)	Ciprofloxacin (mg/l)	Rifampicin (mg/l)	Gentamicin (mg/l)	Vancomycin (mg/l)
<i>S. epidermidis</i> 020A.1	2	0.5–1 (S)	0.015 (S)	2 (R)	2–4 (S)
<i>S. equorum</i> 060A	1–2	0.5 (S)	0.015 (S)	0.125–0.5 (S)	2 (S)
<i>S. capitis</i> 052A	2	0.25 (S)	0.015 (S)	0.5 (S)	1 (S)
<i>S. haemolyticus</i> 065A.1	1–2	0.5–1 (S)	0.015–0.03 (S)	0.25–0.5 (S)	2 (S)
<i>S. hominis</i> M11	1–2	1 (S)	0.015 (S)	1 (S)	2 (S)
<i>S. warneri</i> F16	2	0.25 (S)	0.03 (S)	0.5–1 (S)	2 (S)

CoNS: Coagulase-negative staphylococci; M: Minimum inhibitory concentration; MIC: Minimum inhibitory concentration; R: Resistant; S: Susceptible.

Figure 1. Number of culturable planktonic cells upon incubation with carvacrol. Stationary phase planktonic cells culturability was evaluated after treatment with 4 mM of carvacrol for 5, 15, 45 and 90 min. The statistical differences between control (0 min) and treated samples (5–90 min) were determined by unpaired t-test (two-tailed). The bars represent the mean plus or minus standard deviation of at least three independent assays. CFU: Colony-forming unit.

Table 2. Percentage of dead cells upon 15 and 90 min of exposure to carvacrol.

CoNS species/strains	% of dead cells (PI positive)	
	15 min	90 min
<i>S. epidermidis</i> 020A.1	98.76	99.61 ± 0.50
<i>S. equorum</i> 060A	99.13	99.53 ± 0.34
<i>S. capitis</i> 052A	98.69	98.58 ± 1.06
<i>S. haemolyticus</i> 065A.1	99.67	99.77 ± 0.06
<i>S. hominis</i> M11	99.33	97.82 ± 0.97
<i>S. warneri</i> F16	99.69	99.30 ± 0.96

The values represent the values of a single experiment for 15 min, and the mean plus or minus the standard deviation of three independent assays for 90 min of exposure. CoNS: Coagulase-negative staphylococci; PI: Propidium iodide.

15 min after treatment, no CFU were detected in any of the strains (Figure 1). To confirm that the absence of culturable cells was the result of cell death, flow cytometry analysis was performed using SYTO BC/propidium iodide (LIVE/DEAD) staining. As can be seen in Table 2, more than 97.8% of the cells quantified by flow cytometry were positive for propidium iodide, demonstrating that carvacrol did, in fact, cause cell death.

When analyzing the susceptibility of intact biofilms to carvacrol, a significant reduction in the number of *S. epidermidis* and *S. equorum* culturable cells was observed in the first 5 min of exposure, with a decrease of 0.7 and 1.5 log₁₀ CFU/ml, respectively. Regarding the other CoNS strains, significant reductions were only found after 90 min of exposure, ranging from 2.3, in the case of *S. warneri*, to slightly more than 4.0 log₁₀ CFU/ml in the case

Figure 2. Number of culturable biofilm cells upon incubation with carvacrol.

Biofilm cells were treated with 4 mM of carvacrol for 5 or 90 min. The statistical differences between control (0 min) and treated samples (5 and 90 min) were determined by unpaired t-test (two-tailed). The bars represent the mean plus or minus the standard deviation of at least three independent assays. CFU: Colony-forming unit.

Table 3. Minimum inhibitory concentration of carvacrol (mM) and number of culturable planktonic and biofilm cells (LO CFU/ml) upon 90 min of treatment with carvacrol, and prior to any treatment (CT) for additional 11 coagulase-negative staphylococci isolates.

CoNS species/strains	CAR MIC (mM)	Log CFU/ml			
		Planktonic cells		Biofilm cells	
		CT (0 min)	CAR (90 min)	CT (0 min)	CAR (90 min)
<i>S. epidermidis</i> RP62A (ATCC 35984)	2	8.55 ± 0.11	0	8.27 ± 0.27	3.65 ± 0.31
<i>S. epidermidis</i> 010B	2	8.15 ± 0.60	0	8.36 ± 0.27	3.47 ± 0.81
<i>S. equorum</i> 021B	2	8.38 ± 0.27	0	8.21 ± 0.03	3.31 ± 0.47
<i>S. capitis</i> 048A	2	8.53 ± 0.27	0	8.48 ± 0.21	3.59 ± 0.61
<i>S. capitis</i> 017A	2	8.51 ± 0.09	0	n.p.	n.p.
<i>S. haemolyticus</i> 047A	2	7.97 ± 0.52	0	8.23 ± 0.40	3.35 ± 0.56
<i>S. haemolyticus</i> 008A	2	7.99 ± 0.50	0	7.99 ± 0.83	3.47 ± 0.45
<i>S. hominis</i> 057B	2	7.92 ± 0.74	0	8.17 ± 0.29	3.60 ± 0.51
<i>S. hominis</i> 021A	2	7.96 ± 0.88	0	7.97 ± 0.33	3.48 ± 0.28
<i>S. warneri</i> F14	2	7.90 ± 0.57	0	8.39 ± 0.09	3.42 ± 0.40
<i>S. warneri</i> 055A	2	8.11 ± 0.40	0	n.p.	n.p.

CAR: Carvacrol; CFU: Colony-forming unit; CoNS: Coagulase-negative staphylococci; CT: Control; MIC: Minimum inhibitory concentration; n.p.: Nonperformed.

of *S. epidermidis* and *S. equorum* (Figure 2). The effect of carvacrol on planktonic and biofilm cultures of CoNS was further confirmed by repeating the antimicrobial killing assays using 11 more isolates of the tested species. As seen in Table 3, carvacrol was able to completely eradicate the planktonic cultures and significant killing most biofilm bacteria, in all tested isolates.

Finally, the effect of carvacrol on biofilm structure was also qualitatively analyzed by CLSM (Figure 3). Biofilms formed by *S. epidermidis*, *S. equorum*, *S. haemolyticus* and *S. hominis* exhibited a compact mushroom-like structure, while *S. warneri* and, particularly, *S. capitis* formed less dense biofilms. Despite, these structural differences, all biofilms treated with carvacrol showed a reduced maximum thickness, and the cell density at the lower layers was also reduced. Nevertheless, *S. epidermidis* and *S. haemolyticus* lower layers were apparently less affected by carvacrol.

Carvacrol is more effective than antibiotics against CoNS stationary phase planktonic & biofilm cells

Since, the MIC determination showed that all strains tested were susceptible to most antibiotics used, we used the PSC to evaluate antibiotics efficacy. Carvacrol had a significantly higher impact on the culturability of both

Figure 3. Representative confocal laser scanning microscopy images of biofilms after incubation with carvacrol.

Biofilms grown for 24 h were treated with 4 mM of carvacrol for 90 min. Biofilms were staining with SYTO9 (green). Z-stacks were taken at 5-μm intervals.

CAR: Carvacrol; CT: Control.

planktonic and biofilm cells than the antibiotics tested, after 90 min of incubation (Figure 4). Furthermore, an extra experiment was performed, by allowing incubation with antimicrobial agents for up to 360 min (Supplementary Figure 1). Not surprisingly, longer periods of time increased the activity of antibiotics in both planktonic and biofilm cultures. However, biofilm cultures incubated with carvacrol for 360 min also saw a significant increase in cell death. As such, despite the increase in antibiotic killing activity after 360 min, the activity of carvacrol remained substantially higher than any of the tested antibiotics.

Discussion

Carvacrol has been shown as a promising candidate to replace traditional antimicrobial chemotherapy which antimicrobial potential has already been assessed against several bacteria and fungi species, as *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Klebsiella pneumoniae*, *Escherichia coli*, *Candida albicans*, *Candida tropicalis*, *Torulopsis glabrata* [19], *Porphyromonas gingivalis*, *Fusobacterium nucleatum*, *Streptococcus mutans* [41] and several others *Streptococcus* isolates [42], among others [43–47]. Furthermore, previous studies revealed that carvacrol is capable of inhibiting the growth of *S. aureus* and *S. epidermidis* planktonic bacteria and even interfere with biofilm formation [29] and maturation [48], although, generally, at a lesser extent. However, and despite a few studies on *S. epidermidis* have been published, other relevant biofilm forming CoNS species have been entirely disregarded. Henceforth, due to their increasing clinical relevance, we aimed to investigate carvacrol antimicrobial activity against planktonic and biofilm cells of some of the most commonly isolated clinically relevant CoNS species [32,49–51].

Remarkably, after only 5 min of interaction with carvacrol a significant reduction, up to 4.6 log₁₀ CFU/ml, was observed in planktonic cells culturability. Furthermore, 15 min after adding carvacrol, no CFU were detected in any of the species tested.

The absence of CFU was directly correlated to cell death as at least 97.8% of the cells were positive for propidium iodide (Table 2). This was not surprising since previous studies associated carvacrol mechanism of action with cell-wall biosynthesis blockage [21,41,52]. Interestingly, contrary to what occurs in many cell wall-targeting antibiotics [6], carvacrol was very active against stationary phase planktonic cells. This prompted us to study the effect of carvacrol on biofilms. However, the effect of carvacrol on biofilms was less striking and it took more time, when compared with stationary phase planktonic cells. As both these cell populations are characterized by a lower metabolic activity than exponential planktonic cells, this further suggests that carvacrol activity is not dependent in cell metabolism.

Thus, the differences observed in carvacrol effect between planktonic and biofilm populations may be related to difficulties accessing the cells embedded in the biofilm matrix [53]. Our CLSM analysis partially confirmed this

Figure 4. Number of culturable planktonic and biofilm cells upon 90 min of treatment with carvacrol versus treatment with antibiotics with distinct mechanisms of action.

Stationary phase planktonic cells and 24-h biofilms were incubated with 4 mM of CAR, 4.5 mg/l of CIP, 10 mg/l of RIF, 10 mg/l of GEN or 40 mg/l of VAN, during 90 min. A CT was obtained by plating the bacterial suspension before adding the antimicrobial compounds. (A) *S. epidermidis* 020A.1; (B) *S. equorum* 060A; (C) *S. capitis* 052A; (D) *S. haemolyticus* 065A.1; (E) *S. hominis* M11; (F) *S. warneri* F16. The statistical differences were determined by unpaired *t*-test (two-tailed), between CT and treated samples (CAR, CIP, RIF, GEN and VAN) and between CAR- and antibiotic-treated samples (CIP, RIF, GEN and VAN). The bars represent the mean plus or minus standard deviation of at least three independent assays.

CAR: Carvacrol; CFU: Colony-forming unit; CIP: Ciprofloxacin; CT: Control; GEN: Gentamicin; RIF: Rifampicin; VAN: Vancomycin.

hypothesis, since species with higher biofilm formation ability were less affected in the lower layers of the biofilm structure.

To better assess the impact of carvacrol as an alternative to treat *S. epidermidis* biofilm infections, we compared the killing efficacies of carvacrol with four antibiotics used to treat staphylococcal infections with distinct mechanisms of action: nucleic acids (rifampicin and ciprofloxacin) or protein synthesis inhibitors (gentamicin) and cell wall targeting (vancomycin). Although, the differences between the decrease in the culturability of cells in contact with carvacrol or antibiotics were more pronounced after shorter incubation periods (90 min), the efficacy of carvacrol remained considerably higher than the one of other antimicrobials even after 360 min of incubation.

Conclusion

To our knowledge, this is the first study reporting the antimicrobial effect of carvacrol against a range of CoNS species, as well as to compare the efficacy of this natural compound with the antibacterial effect of traditionally used antibiotics. Regardless of the differences between *in vivo* and *in vitro* conditions, this study shows a high potential of carvacrol for the control of CoNS-originated infections, which gains particular importance given the low efficacy of several antimicrobial therapies focused in antibiotics. Remarkably, carvacrol action against biofilms was very promising in comparison to antibiotics. Further studies should include the use of *in vivo* models in order to evaluate the efficacy and relevance of carvacrol with the presence of host factors, since several compounds are known to lose their activity in the presence of host proteins.

Summary points

- Carvacrol displays a remarkable antibacterial activity against several *Staphylococcus* spp. bacteria, both against planktonic and biofilm cells.
- A concentration of 4 mM of carvacrol was able to completely eradicate an initial load of 10^8 colony-forming unit per ml of planktonic cells after 15 min of treatment.
- Biofilms were highly affected by carvacrol, with decreases in the culturability of about 3 log₁₀ colony-forming unit per ml after 90 min.
- The antimicrobial effect of carvacrol was significantly higher than the one observed with four traditional antistaphylococcal antibiotics.

Supplementary data

To view the supplementary data that accompany this paper please visit the journal website at: www.futuremedicine.com/doi/suppl/10.2217/fmb-2017-0122.

Financial & competing interests disclosure

This study was supported by the Portuguese Foundation for Science and Technology (FCT) under the scope of the strategic funding of UID/BIO/04469/2013 unit and COMPETE 2020 (POCI-01-0145-FEDER-006684) and the Project RECI/BBB-EBI/0179/2012 (FCOMP-01-0124-FEDER-027462). A França is supported by the FCT fellowship SFRH/BPD/99961/2014. N Cerca is an investigator FCT. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Ethical conduct of research

The authors state that they have obtained appropriate institutional review board approval or have followed the principles outlined in the Declaration of Helsinki for all human or animal experimental investigations. In addition, for investigations involving human subjects, informed consent has been obtained from the participants involved.

References

Papers of special note have been highlighted as: • of interest; •• of considerable interest

1. Becker K, Heilmann C, Peters G. Coagulase-negative staphylococci. *Clin. Microbiol. Rev.* 27(4), 870–926 (2014).
2. Spanu T, Sanguinetti M, Ciccaglione D *et al.* Use of the VITEK 2 system for rapid identification of clinical isolates of staphylococci from bloodstream infections. 41(9), 4259–4263 (2003).

3. Huang CR, Lu CH, Wu JJ *et al.* Coagulase-negative staphylococcal meningitis in adults: clinical characteristics and therapeutic outcomes. *Infection* 33(2), 56–60 (2005).
4. Shin JH, Kim SH, Jeong HS *et al.* Identification of coagulase-negative staphylococci isolated from continuous ambulatory peritoneal dialysis fluid using 16S ribosomal RNA, *tuf*, and *SodA* gene sequencing. *Perit. Dial. Int.* 31(3), 340–346 (2011).
5. Donlan RM, Costerton JW. Biofilms: survival mechanisms of clinically relevant microorganisms. *Clin. Microbiol.* 15(2), 167–193 (2002).
6. Cerca N, Martins S, Cerca F *et al.* Comparative assessment of antibiotic susceptibility of coagulase-negative staphylococci in biofilm versus planktonic culture as assessed by bacterial enumeration or rapid XTT colorimetry. *J. Antimicrob. Chemother.* 56(2), 331–336 (2005).
7. del Pozo JL, Patel R. The challenge of treating biofilm-associated bacterial infections. *Clin. Pharmacol. Ther.* 82(2), 204–209 (2007).
8. Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent infections. *Science* 284(5418), 1318–1322 (1999).
9. Singh G, Kapoor IPS, Pandey SK, Singh UK, Singh RK. Studies on essential oils: part 10; antibacterial activity of volatile oils of some spices. *Phyther. Res.* 16(7), 680–682 (2002).
10. Kalemka D, Kunicka A. Antibacterial and antifungal properties of essential oils. *Curr. Med. Chem.* 10(10), 813–829 (2003).
11. Nostro A, Roccaro AS, Bisignano G *et al.* Effects of oregano, carvacrol and thymol on *Staphylococcus aureus* and *Staphylococcus epidermidis* biofilms. *J. Med. Microbiol.* 56(4), 519–523 (2007).
- **In this article, Nostro *et al.* presents a study of the effects of distinct natural compounds on two *Staphylococcus* species, in which several studies were based to assess the potential antimicrobial effect of carvacrol and other essential oils.**
12. Reichling J, Schnitzler P, Suschke U, Saller R. Essential oils of aromatic plants with antibacterial, antifungal, antiviral, and cytotoxic properties – an overview. *Forsch. Komplementarmed.* 16(2), 79–90 (2009).
13. Hüsnü K, Başer C, Demirci F. Chemistry of essential oils. In: *Flavours and Fragrances Chemistry, Bioprocessing and Sustainability*. Berger RG (Ed.). Springer-Verlag Berlin Heidelberg, 43–86 (2007).
14. Kirimer N, Baser KHC, Tumen G. Carvacrol-rich plants in Turkey. *Chem. Nat. Compd.* 31(1), 37–41 (1995).
15. Solórzano-Santos F, Miranda-Novales MG. Essential oils from aromatic herbs as antimicrobial agents. *Curr. Opin. Biotechnol.* 23(2), 136–141 (2012).
16. Hajhashemi V, Ghannadi A, Karim S. Antinociceptive and anti-inflammatory effects of *Satureja hortensis* L. extracts and essential oil. 82, 83–87 (2002).
17. Prieto JM, Iacopini P, Cioni P, Chericoni S. *In vitro* activity of the essential oils of *Origanum vulgare*, *Satureja montana* and their main constituents in peroxynitrite-induced oxidative processes. *Food Chem.* 104(3), 889–895 (2007).
18. Karkabounas S, Kostoula OK, Daskalou T *et al.* Anticarcinogenic and antiplatelet effects of carvacrol. *Exp. Oncol.* 28(2), 121–125 (2006).
19. Aligiannis N, Kalpoutzakis E, Mitaku S, Chinou IB. Composition and antimicrobial activity of the essential oils of two *Origanum* species. *J. Agric. Food Chem.* 49(9), 4168–4170 (2001).
20. Svoboda K, Hampson J. Bioactivity of essential oils of selected temperate aromatic plants: antibacterial, antioxidant, antiinflammatory and other related pharmacological activities. *Spec. Chem. 21st Century* 1–17 (1999).
21. Machado D, Gaspar C, Palmeira-de-Oliveira A *et al.* *Thymbra capitata* essential oil as potential therapeutic agent against *Gardnerella vaginalis* biofilm-related infections. *Future Microbiol.* 12(5), 407–416 (2017).
- **This manuscript refers to *in vitro* assays performed with *Thymbra capitata* essential oil and some of its major constituents, highlighting the therapeutic potential of carvacrol to treat biofilm-related infections.**
22. Hammer KA, Carson CF, Rileya TV. Effects of *Melaleuca alternifolia* (tea tree) essential oil and the major monoterpene component terpinen-4-ol on the development of single- and multistep antibiotic resistance and antimicrobial susceptibility. *Antimicrob. Agents Chemother.* 56(2), 909–915 (2012).
23. de Sousa JP, Torres R de A, de Azerêdo GA, Figueiredo RCBQ, Vasconcelos MA, de Souza EL. Carvacrol and 1,8-cineole alone or in combination at sublethal concentrations induce changes in the cell morphology and membrane permeability of *Pseudomonas fluorescens* in a vegetable-based broth. *Int. J. Food Microbiol.* 158(1), 9–13 (2012).
24. Nostro A, Marino A, Blanco AR *et al.* *In vitro* activity of carvacrol against staphylococcal preformed biofilm by liquid and vapour contact. *J. Med. Microbiol.* 58(6), 791–797 (2009).
25. Lambert RJW, Skandamis PN, Coote PJ, Nychas GJE. A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *J. Appl. Microbiol.* 91(3), 453–462 (2001).
26. Ben Arfa A, Combes S, Preziosi-Belloy L, Gontard N, Chalier P. Antimicrobial activity of carvacrol related to its chemical structure. *Lett. Appl. Microbiol.* 43(2), 149–154 (2006).
- **In this article, Arfa *et al.* studied the antimicrobial activity of carvacrol and the relation to its chemical structure, thus trying to clarify the mechanism of action of this essential oil.**

27. Cristani M, D'Arrigo M, Mandalari G *et al.* Interaction of four monoterpenes contained in essential oils with model membranes: implications for their antibacterial activity. *J. Agric. Food Chem.* 55(15), 6300–6308 (2007).
28. Helander IM, Alakomi H-L, Latva-Kala K *et al.* Characterization of the action of selected essential oil components on Gram-negative bacteria. *J. Agric. Food Chem.* 46(9), 3590–3595 (1998).
29. Knowles J, Roller S. Efficacy of chitosan, carvacrol, and a hydrogen peroxide-based biocide against foodborne microorganisms in suspension and adhered to stainless steel. *J. Food Prot.* 64(10), 1542–1548 (2001).
30. Knowles JR, Roller S. Antimicrobial action of carvacrol at different stages of dual-species biofilm development by *Staphylococcus aureus* and *Salmonella enterica* serovar Typhimurium. *Society* 71(2), 797–803 (2005).
31. Nostro A, Blanco AR, Cannatelli MA *et al.* Susceptibility of methicillin-resistant staphylococci to oregano essential oil, carvacrol and thymol. *FEMS Microbiol. Lett.* 230(2), 191–195 (2004).
32. Oliveira F, Cerca N. Antibiotic resistance and biofilm formation ability among coagulase-negative staphylococci in healthy individuals from Portugal. *J. Antibiot. (Tokyo)* 66(12), 739–741 (2013).
33. Oliveira F, Lima CA, Bras S, França A, Cerca N. Evidence for inter- and intraspecies biofilm formation variability among a small group of coagulase-negative staphylococci. *FEMS Microbiol. Lett.* 362(20), 1–20 (2015).
34. National Committee for Clinical Laboratory Standards. Methods for dilution: antimicrobial susceptibility tests for bacteria that grow aerobically; Approved Standard (9th Edition).
35. Ben-amor K, Heilig H, Smidt H *et al.* Genetic diversity of viable, injured, and dead fecal bacteria assessed by fluorescence-activated cell sorting and 16S rRNA gene analysis. *Society* 71(8), 4679–4689 (2005).
36. Freitas AI, Vasconcelos C, Vilanova M, Cerca N. Optimization of an automatic counting system for the quantification of *Staphylococcus epidermidis* cells in biofilms. *J. Basic Microbiol.* 54(7), 750–757 (2014).
37. Bayer HealthCare Pharmaceuticals Inc. CIPRO[®] (ciprofloxacin hydrochloride) Tablets, 1–31 (2004).
38. National Committee for Clinical Laboratory Standards. Methods for dilution: antimicrobial susceptibility tests for bacteria that grow aerobically (5th Edition). Approved Standard M7-A5. NCCLS, Wayne, PA, USA.
39. Demczar DJ, Nafziger AN, Bertino JS. Pharmacokinetics of gentamicin at traditional versus high doses: implications for once-daily aminoglycoside dosing. *Antimicrob. Agents Chemother.* 41(5), 1115–1119 (1997).
40. European Committee on Antimicrobial Susceptibility Testing. EUCAST: Definitions of clinical breakpoints and epidemiological cut-off values (2017). www.eucast.org/clinical_breakpoints/
41. Ciandrini E, Campana R, Federici S *et al.* *In vitro* activity of carvacrol against titanium-adherent oral biofilms and planktonic cultures. *Clin. Oral Investig.* 18(8), 2001–2013 (2014).
- **In this article, the antibacterial effect of carvacrol was tested against biofilms, stating its potential to be used in biofilm-related infections, as addressed in our manuscript.**
42. Magi G, Marini E, Facinelli B. Antimicrobial activity of essential oils and carvacrol, and synergy of carvacrol and erythromycin, against clinical, erythromycin-resistant Group A Streptococci. *Front. Microbiol.* 6, 1–7 (2015).
43. Ahmad A, Khan A, Akhtar F *et al.* Fungicidal activity of thymol and carvacrol by disrupting ergosterol biosynthesis and membrane integrity against *Candida*. *Eur. J. Clin. Microbiol. Infect. Dis.* 30(1), 41–50 (2011).
44. Chavan PS, Tupe SG. Antifungal activity and mechanism of action of carvacrol and thymol against vineyard and wine spoilage yeasts. *Food Control.* 46, 115–120 (2014).
45. Didry N, Dubreuil L, Pinkas M. Activity of thymol, carvacrol, cinnamaldehyde and eugenol on oral bacteria. *Pharm. Acta Helv.* 69(1), 25–28 (1994).
46. Ultee A, Slump RA, Steging G, Smid EJ. Antimicrobial activity of carvacrol toward *Bacillus cereus* on rice. *J. Food Prot.* 63(5), 620–624 (2000).
47. Ultee A, Bennik MHJ, Moezelaar R. The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Appl. Environ. Microbiol.* 68(4), 1561–1568 (2002).
48. Burt SA, Ojo-Fakunle VTA, Woertman J, Veldhuizen EJA. The natural antimicrobial carvacrol inhibits quorum sensing in *Chromobacterium violaceum* and reduces bacterial biofilm formation at sub-lethal concentrations. *PLoS ONE* 9(4), 1–6 (2014).
49. Otto M. *Staphylococcus epidermidis* – the ‘accidental’ pathogen. *Nat. Rev. Microbiol.* 7(8), 555–567 (2009).
50. Akinkunmi EO, Lamikanra A. Species distribution and antibiotic resistance in coagulase-negative staphylococci colonizing the gastrointestinal tract of children in Ile-Ife, Nigeria. *Trop. J. Pharm. Res.* 9(1), 35–43 (2010).
51. Delmas J, Chacornac JP, Robin F, Giammarinaro P, Talon R, Bonnet R. Evaluation of the VITEK 2 system with a variety of *Staphylococcus* species. *J. Clin. Microbiol.* 46(1), 311–313 (2008).
52. Salgueiro LR, Pinto E, Goncalves MJ *et al.* Chemical composition and antifungal activity of the essential oil of *Thymbra capitata*. *Planta Med.* 70(6), 572–575 (2004).
53. Mah TFC, O'Toole GA. Mechanisms of biofilm resistance to antimicrobial agents. *Trends Microbiol.* 9(1), 34–39 (2001).

Histone acetyltransferase encoded by *NGG1* is required for morphological conversion and virulence of *Candida albicans*

De-Dong Li^{1,2,3}, Beth Burgwyn Fuchs², Yan Wang³, Xiao-Wen Huang^{2,4}, Dan-Dan Hu³, Yan Sun¹, Dong Chai¹, Yuan-Ying Jiang^{*,3} & Eleftherios Mylonakis^{**,2}

¹Clinical Pharmacy Center, Department of Pharmacy, Chinese PLA General Hospital, Beijing 100853, China

²Division of Infectious Disease, Rhode Island Hospital, Alpert Medical School & Brown University, RI 02903, USA

³New Drug Research & Development Center, School of Pharmacy, Second Military Medical University, Shanghai 200433, China

⁴Department of Dermatology, Nanfang Hospital, Southern Medical University, Guangzhou 510515, Guangdong, China

* Author for correspondence: 13761571578@163.com

** Author for correspondence: emylonakis@lifespan.org

Aim: To assess the function of *Ngg1* in *Candida albicans* and reveal the role of *NGG1* in the morphological conversion and virulence of *C. albicans*. **Materials & methods:** *C. albicans* *NGG1* gene was deleted in the wild-type strain SC5314 and the function of *Ngg1* was assessed by western blot analysis. The phenotypes and the virulence of the *ngg1* mutants were examined. Microarray analysis was performed to explore the mechanism. **Results:** The *ngg1* mutants attenuated acetylated histone H3, obviously reduced filamentous growth and showed significantly diminished pathogenicity in all the infection models. **Conclusion:** This study suggested the histone acetyltransferase activity of *C. albicans* *Ngg1* and revealed the important role of *NGG1* in morphological conversion and virulence of *C. albicans*.

First draft submitted: 10 May 2017; Accepted for publication: 27 July 2017; Published online: 7 November 2017

Keywords: *Caenorhabditis elegans* • *Candida albicans* • *Galleria mellonella* • histone acetyltransferase • morphological conversion • *NGG1* • virulence

Candida albicans, the most common fungal pathogen, causes diseases varying from superficial mucosal infections to lethal systemic disorders, especially in immunocompromised individuals [1]. Bloodstream infections caused by *C. albicans* are associated with an estimated 40% mortality [2]. The virulence of *C. albicans* is mainly attributed to its ability to undergo reversible morphological conversion utilizing hyphae and biofilm to pathogenize a host [3,4]. The study on virulence factors of *C. albicans*, especially morphological factors, is of great significance for the discovery of antifungal targets and the treatment of fungal infections.

In *C. albicans*, morphological conversion is tightly correlated with the expression of hyphal-specific genes which are mainly controlled by transcriptional factors, such as Efg1, Cph1, Flo8 and Tec1 [3]. For example, Efg1 is thought to be the major regulator of hyphal formation, which is required for hyphal formation in response to serum, CO₂, neutral pH and *N*-acetyl-D-glucosamine (GlcNAc) in liquid media [5–7]. Cph1 and its upstream activating pathway are required for hyphal formation on solid Spider media [8,9].

In addition, recent studies demonstrated that the chromatin modifying histone acetyltransferase (HAT) also contributes to the morphological conversion and virulence of *C. albicans*. The HAT complex NuA4, which primarily acetylates H4 and H2A in *C. albicans*, was responsible for nucleosomal H4 acetylation in the promoters of hyphal-specific genes during hyphal induction [10]. Gcn5, a catalytic subunit of many HAT complexes, contributes to the invasive and filamentous growth of *C. albicans* [11]. Moreover, Rtt109, another HAT which acetylates histone H3 lysine 56 in *C. albicans*, is required for virulence of *C. albicans* and affects the morphology of clones [12].

HAT genes are not specific to yeast and they are evolutionarily conserved and part of the human genome. In human cells, transcriptional adapter 3 (Tada3) plays a key role in linking HAT-containing complexes to p53; thereby regulating p53 acetylation, stability and transcriptional activation following DNA damage [13]. The fungal ortholog Ada3 in *Saccharomyces cerevisiae* is a subunit of HAT complexes ADA, SAGA and SLIK, and plays an important role in chromatin modification and transcription activation [14–16]. There are homologs of Ada3 in *Candida* spp. including *C. albicans*, *C. glabrata*, *C. dubliniensis* and *C. parapsilosis* (available at *C. albicans* genome database [17]). Despite extensive analysis of human Tada3 and *S. cerevisiae* Ada3, the role of their homologs in *Candida* spp. is not well understood. In *C. albicans*, the homolog of Ada3 is known as Ngg1. In this study, we confirm the HAT activity of Ngg1 in *C. albicans* and investigated the role of *NGG1* in the morphological conversion and pathogenicity of *C. albicans*.

Materials & methods

Strains & growth conditions

C. albicans strains used in this study are listed in Table 1. Strains were routinely grown in YPD (1% yeast extract, 2% peptone and 2% dextrose) liquid media at 30°C in a shaking incubator. *C. elegans glp-4; sek-1* strains were propagated on nematode growth media on lawns of *Escherichia coli* OP50 by using standard methods [18]. To evaluate the growth rate of the *C. albicans* strains, the organisms were grown overnight at 30°C in liquid YPD media and then subcultured by inoculation into fresh YPD media to 1×10^6 cells/ml. The cells were cultured at 30°C under constant shaking (200 rpm) and counted at different time points after culture.

Generating homozygous mutant strain

The *ngg1*Δ/Δ mutant was generated using the pop-out flipper cassette from pSFS2 containing the dominant nourseothricin resistance marker CaSAT1 as described previously [19]. Briefly, a 0.54 kb ApaI-XhoI fragment containing the 5' end of *NGG1* gene was amplified with primers NGG1-1 and NGG1-2 and cloned into the pSFS2 vector upstream of the *MAL2p-CaFLP-CaSAT1* cassette, yielding plasmid pdl701. In the same way, a 0.61 kb SacII-SacI fragment containing the 3' end of the *NGG1* gene was amplified with primers NGG1-3 and NGG1-4 and cloned into the pdl701 vector downstream of the *MAL2p-CaFLP-CaSAT1* cassette, yielding the final plasmid pdl702. The pdl702 plasmid was digested with ApaI and SacI and the 5.33 kb fragment was used for transformation of *C. albicans* SC5314 using the lithium acetate method [20]. Nourseothricin resistant transformants were identified via PCR using primers NGG1-7, NGG1-8, NGG1-9 and NGG1-10 (Supplementary Figure 1). After deleting the nourseothricin resistance marker (*caSAT1*), the same procedure was repeated to delete the remaining functional allele of *NGG1*. A 0.59 kb fragment containing internal sequence of *NGG1* was amplified

Table 1. *Candida albicans* strains used in this study.

Strain	Relevant genotype [†]	Source or references
SC5314	Wild-type	
<i>ngg1</i> -A	<i>ngg1::FRT/ngg1::FRT</i>	This study
<i>ngg1</i> -B	<i>ngg1::FRT/ngg1::FRT</i>	This study

[†]Apart from the indicated features, all strains have the genotype of their parents.

with primers NGG1-11 and NGG1-12 to identify the deletion of *NGG1*. Primers used to identify all strains are listed in Supplementary Table 1.

Western blot analysis

C. albicans cells were grown in 20 ml of YPD liquid media at 30°C. Exponentially growing *C. albicans* cells were washed with PBS, resuspended in 0.5 ml of lysis buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1% Triton X-100 and 1 × proteinase inhibitor mix) and then transferred to Lysis Matrix C tube. Cells were lysed at 4°C using a Fast-Prep system (FastPrep-24™ 5G; MP Biomedical, CA, USA). Cell lysates were centrifuged at 13,000 r.p.m., 1 min at 4°C and repeated once. The concentration of total protein was measured using bicinchoninic acid (BCA) method. Proteins were boiled for 5 min in loading buffer. A total of 16 µg proteins were separated by 15% SDS-PAGE gel and transferred to polyvinylidene difluoride membranes. After blocking with 5% non-fat milk in PBS-T, membranes were probed with rabbit anti-H3 antibody (1:1000; Abcam, MA, USA) or rabbit anti-H3Ac antibody (1:2000; Millipore, MA, USA) overnight at 4°C. After washing for three-times with PBS-T, membranes were incubated with HRP-conjugated goat anti-rabbit IgG (H + L) antibody (1:1000; Applygen, Beijing, China) for 1 h at room temperature. After further washing with PBS-T, bounds were detected by ECL detecting system (Millipore).

In vitro biofilm formation assay

Biofilm formation assay was carried out as described by Nobile *et al.* [21] with slight modifications. Briefly, 2 ml *C. albicans* suspensions (1.0×10^6 cells/ml in Spider media) were inoculated into a 12-well plate with a prepared silicone square (1.5 × 1.5 cm cut from Cardiovascular Instrument silicone sheets [PR72034-06N, Bectec Medical, Inc, CA, USA]) and incubated statically at 37°C. After allowing 90 min for adhesion, the squares were washed with PBS slightly and moved to a fresh 12-well plate containing 2 ml of fresh Spider media. The plates were incubated at 37°C for an additional 60 h at 75 r.p.m. agitation, to allow for biofilm formation. Biofilms were removed from their respective silicone squares, dried overnight, and weighed the following day. The average biomass for each strain was calculated from four independent samples.

Spot assays

To assess the adaptability of the *ngg1* mutants to various stresses, the *C. albicans* wild-type strain SC5314 and the *ngg1Δ/Δ* mutants were all grown to mid-log phase. Strains were washed, resuspended in PBS to 1×10^6 cells/ml and spotted as tenfold serial dilution onto YPD plates with or without chemical agents. Growth differences were detected after a 48-h incubation period.

Virulence assays

The *C. elegans* infection model was carried out using previously described protocols [18]. Briefly, 100 µl of overnight grown *C. albicans* was spread into a square lawn on a BHI plate containing kanamycin (45 µg/ml), followed by incubation for approximately 20 h at 30°C. A total of 400–500 synchronized adult *C. elegans glp-4; sek-1* nematodes were added to the center of the *C. albicans* lawns. The plates were incubated at 25°C for 4 h. Worms were then carefully collected and washed with sterile M9. A total of 60–70 worms were then pipetted into a single well of a 6-well tissue culture plate containing 2 ml of liquid media (80% M9, 20% BHI) and kanamycin (45 µg/ml). Dead worms were scored daily and removed from the assay.

The *Galleria mellonella* infection model was carried out using previously described protocols [22]. Briefly, *G. mellonella* larvae (Vanderhorst Wholesale, Inc, OH, USA) at their final instar stage were inoculated with 1×10^6 CFUs of *C. albicans* suspended in PBS. Each infection group contained 16 randomly chosen larvae of the appropriate weight (300 ± 25 mg). The inoculum was injected in a 10 µl volume directly to the last, left proleg using a Hamil-

ton syringe [22]. After inoculation, larvae were incubated at 37°C. A mock inoculation with PBS was performed in each experiment to monitor killing due to physical injury or infection by pathogenic contaminants. The number of dead larvae was scored daily.

Mice were infected with *C. albicans* according to an established protocol [23]. CD1 female mice (6-week old, 18–22 g) were infected with 1.5×10^6 CFUs of *C. albicans* suspended in PBS via a tail vein injection in a 100 µl volume. A total of 15 mice were inoculated per strain tested. After 48 h of infection, kidneys were harvested for fungal burden evaluation and pathological analysis ($n = 5$). The other ten mice per group were observed daily for survival assay.

Gene expression analysis

Exponentially growing *C. albicans* cells were harvested, washed twice with PBS and resuspended in Spider media to 5×10^6 cells/ml. Following incubating statically at 37°C for 1 h, the cells were collected and stored in liquid nitrogen. *Candida* total RNA preparation and *Candida* genome microarray (CapitalBio Corp, Beijing, China) were performed as described previously [24]. For biological replicates, triplicate independent experiments were carried out and only genes showing a consistent pattern were considered. The genes were named according to the *C. albicans* genome database and divided according to their biological functions.

RT-qPCR was performed to confirm the microarray results. cDNA was synthesized from total RNA using the cDNA synthesis kit (TaKaRa, Dalian, China). RT-qPCR reactions were performed with SYBR Green I (TaKaRa) using LightCycler system (Roche Molecular Biochemical, Basel, Switzerland). Primers used in RT-qPCR are shown in Supplementary Table 2. The change of fluorescence in each cycle was monitored by the system and the threshold cycle (C_T) was measured. 18S rRNA was used as an internal control. Gene expression level of the *ngg1* mutant relative to that of the wild-type strain was calculated using the formula $2^{-\Delta\Delta C_T}$, where ΔC_T was the C_T value of target genes minus that of 18S and $\Delta\Delta C_T$ was the mean ΔC_T values of the *ngg1* mutant minus that of the wild-type strain.

Statistical analyses

Survival curves were generated by using the Kaplan–Meier method and differences were determined by using the log-rank test (STATA software). Student's *t*-test was used to compare biofilm formation. A *p*-value of <0.05 was considered statistically significant.

Results

HAT activity of Ngg1 in *C. albicans*

Ada3 is an important HAT that is well studied in mammals and *S. cerevisiae* [13–14,25–26]. To explore whether there are homologs of Ada3 in *C. albicans*, we searched the *C. albicans* genome database and found that protein Ngg1 (orf19.3023) shared high similarity with *S. cerevisiae* Ada3. As shown in Figure 1A, all of human Tada3, *S. cerevisiae* Ada3 and *C. albicans* Ngg1 contain an Ada3 domain, which is an essential component of the Ada transcriptional coactivator complex and is evolutionarily conserved among eukaryotes [27]. Moreover, Protein BLAST was performed to compare the similarity of these three amino acid sequences. The analysis showed that *C. albicans* Ngg1 and *S. cerevisiae* Ada3 have 51% amino acid similarity with an *E*-value of 9×10^{-86} , while the *C. albicans* Ngg1 and human Tada3 have 34% amino acid similarity with an *E*-value of 9×10^{-12} (Supplementary Figure 2). These results indicated that compared with the human protein Tada3, *C. albicans* Ngg1 shows much higher conservation with *S. cerevisiae* Ada3.

To explore the function of *C. albicans* Ngg1, we constructed an *ngg1*Δ/Δ mutant using the SAT1 cassette strategy [19]. But unfortunately, though different transformation methods (including lithium acetate transformation and electroporation [20,28–30]) were attempted for more than three-times each, we could not generate a reconstituted strain. We suspect that the deletion of *NGG1* affects the transformation of the strain [31] as no colony could grow after the transformation (data not shown). Because of this limitation and in order to control unwanted genetic rearrangements that might happen during mutant construction in *C. albicans* [32,33], two independent *ngg1*Δ/Δ isolates were derived for evaluation [34] (Table 1). As shown in Figure 1B, compared with the wild-type strain SC5314, deletion of *NGG1* drastically decreased the acetylation of histone H3 in both *ngg1*-A and *ngg1*-B. This finding suggests that *C. albicans* Ngg1 is similar to *S. cerevisiae* Ada3 as a HAT and is required for histone H3 acetylation in *C. albicans*.

Figure 1. HAT activity of Ngg1 in *Candida albicans*. (A) Schematic depiction of Ada3 domain in *C. albicans* Ngg1, *Saccharomyces cerevisiae* Ada3 and human Tada3. (B) Western blot analysis of the acetylation levels of histone H3 in the wild-type strain SC5314, *ngg1-A* and *ngg1-B*.

NGG1 plays an important role in morphological conversion & growth of *C. albicans*

C. albicans can grow in multiple morphological states and individual cells appear as budded or filamentous cells [3]. To analyze the role of *NGG1* in the morphological conversion of *C. albicans*, we used various hyphae-inducing media, including Lee's, Spider, serum-containing and yeast-extract peptone sucrose (YPS) embedded media, to examine the phenotype of the *ngg1* mutant strains. On solid Lee's and Spider media, the wild-type strain produced wrinkled colonies with radial hyphae around, while the *ngg1* mutants formed defective smooth colonies (Figure 2A). The *ngg1* mutants demonstrated impaired hyphal development in liquid hyphae-inducing media as well. As shown in Figure 2B, the *ngg1* mutants developed short filaments or budded cells; while SC5314 produced florid filaments after 3.5-h incubation at 37°C. Prolonging the incubation time to 8 h could not improve the hyphal development of the mutants (Figure 2C). The *ngg1* mutants were also defective in invasive growth when embedded in YPS media. Under the embedded condition, the wild-type colony invaded well with long heterogeneous filaments, while the *ngg1* mutants showed difficulty to invade the matrix exhibiting smooth clones in the agar (Figure 2A). These observations indicated that *NGG1* is required for the filamentous growth of *C. albicans* under both aerobic and microaerophilic conditions.

Since the *S. cerevisiae ada3*-null mutant demonstrates decreased utilization of carbon sources and slower growth [35], we then examined the proliferation ability of the *ngg1* mutants in YPD liquid media. As shown in Figure 2D and E, *C. albicans ngg1* mutants grew slowly and demonstrated a moderately longer doubling time (T_d) compared with the wild-type strain (92 and 91 min for the mutants vs 71 min for the wild-type at 30°C; 66 and 65 min for the mutants vs 52 min for the wild-type at 37°C), which implied the energy utilization disorder of the *ngg1* mutants. Hyphal formation and energy utilization are critical for the growth of *C. albicans* biofilms [36,37], so we further examined the biofilm formation of the *ngg1* mutants. As expected, the *ngg1* mutants showed significant reduced biofilm biomass, compared with SC5314 according to biofilm dry mass determination ($p < 0.01$, Figure 2F).

NGG1 is required for stress responses of *C. albicans*

To evaluate the ability of *ngg1* mutants to adapt to various environmental stresses, the strains were exposed to oxidative and cell wall stressors. As shown in Figure 3, the *ngg1* mutants were hypersensitive to various oxidative stresses. More specifically, under 5 mM hydrogen peroxide (H_2O_2), the *ngg1* mutants showed slight growth defects. However, exposure to 0.1 mM menadione and 1 mM diamide caused apparent growth inhibition of the mutants.

Figure 2. *NGG1* is needed for morphological conversion and growth of *Candida albicans*. Results for the wild-type strain SC5314, *ngg1*-A and *ngg1*-B are shown. **(A)** Filamentation defect of the *ngg1* mutants on solid hyphae-inducing media. Time of incubation and temperature were as follows: YPD, 4 days at 30°C; Lee's, 5 days at 30°C; Spider, 5 days at 37°C; YPS (embedded growth), 3 days at 25°C. **(B & C)** Filamentation defect of the *ngg1* mutants in liquid hyphae-inducing media. Growth was monitored after 3.5 h (B) or 8 h (C) incubation at 37°C. **(D & E)** Growth curve of log-phase cultures of the indicated strains at 30 or 37°C. Doubling time was derived by linear regression on the log₁₀ scale. **(F)** Biofilm dry mass determination of *C. albicans* strains. Strains were cultured in Spider media. After 60-h incubation, the biofilm biomass was dried and weighed. The data are shown as mean ± standard deviation from four independent experiments. Statistical analysis was determined by Student's *t*-test.

** Indicates *p* < 0.01, compared with SC5314.

CFU: Colony-forming unit; FBS: Fetal bovine serum; YPD: Yeast-extract peptone dextrose; YPS: Yeast-extract peptone sucrose.

Figure 3. Deletion of *NGG1* gene compromises adaptability to various stresses. *Candida albicans* wild-type strain SC5314 and two *ngg1*Δ/Δ mutants *ngg1*-A, *ngg1*-B were all grown to mid-log phase and then spotted as serial dilutions onto YPD plates with or without chemical agents as indicated. Growth differences were detected after a 48-h incubation period. SDS: Sodium dodecyl sulphate; YPD: Yeast-extract peptone dextrose.

Meanwhile, the sensitivity of *ngg1* mutants to cell wall stresses including sodium dodecyl sulphate and caffeine were dramatically increased, which strongly suggested a cell wall defect.

We further investigated the impact of *NGG1* deletion on drug susceptibility to different antifungal agents. Like other cell wall stresses, the *ngg1* mutants were sensitive to caspofungin, with 0.25 µg/ml caspofungin completely preventing growth of the mutants (Figure 3). Genotoxic drug 5-fluorocytosine was used to detect the genomic stability of the mutants. The *ngg1* mutants were hypersensitive to 5-fluorocytosine, which implied the impaired DNA damage repair capability (Figure 3). Moreover, the *ngg1* mutants showed hypersensitivity to azole antifungal agents, demonstrated through growth inhibition at 4 µg/ml fluconazole and 4 µg/ml ketoconazole. However, the deletion of *NGG1* did not change the sensitivity of the strains to another membrane targeting antifungal agent amphotericin B (Figure 3).

Pathogenicity of *C. albicans* is diminished in the absence of *NGG1*

Morphological conversion and biofilm formation are critical virulence determinants of *C. albicans* [38]. Considering the defects of the *ngg1* mutants in morphological conversion and biofilm formation, the virulence of *C. albicans* was evaluated using different infection models. First, two invertebrate infection models including *Caenorhabditis elegans* and *G. mellonella* were used. As shown in Figure 4, the *ngg1* mutants were extremely deficient in pathogenesis, compared with the wild-type strain in both *C. elegans* and *G. mellonella* models. More specifically, after 4 days of infection with *C. albicans*, over 80% *C. elegans* worms in the SC5314 group died, while only 25% worms died in the *ngg1* mutant groups ($p < 0.001$, Figure 4A). Consistent with this result, 1×10^6 CFUs/larva of the wild-type strain SC5314 caused all the *G. mellonella* larvae to die within 3 days. In contrast, over 60% *G. mellonella* larvae were still alive 3 days after infection with the *ngg1* mutants ($p < 0.001$, Figure 4B).

Further, a murine systemic candidiasis model was used to confirm the results. We infected CD1 female mice with 1.5×10^6 CFUs of the wild-type strain or the *ngg1* mutants by tail vein injection. The mortality of the mice was monitored daily and the results showed that the *ngg1* mutants were avirulent in mice. More specifically, as shown in Figure 5A, wild-type strain caused all the mice to die within 9 days while all the mice survived for more than 21 days after infection with the *ngg1* mutant strains. To determine whether these phenotypes correlated with proliferation of the pathogen in the animal, we retrieved kidneys from infected mice and assessed the fungal burden at day 2 postinfection. Fungi were recovered from the evaluated kidneys of wild-type infected mice, and the fungal

Figure 4. NGG1 is essential for virulence of *Candida albicans* in the *Caenorhabditis elegans* and *Galleria mellonella* infection model. (A) Survival of *C. elegans* over 6 days ($n = 60\text{--}70$ worms per strain of *C. albicans*). Compared with infection with *C. albicans* SC5314, the *ngg1* mutants are less virulent to *C. elegans* ($p < 0.001$). (B) Survival curves of *G. mellonella* infected with different *C. albicans* mutants ($n = 16$ larvae per strain of *C. albicans*). The *ngg1* mutants are less virulent to *G. mellonella* compared with the groups infected with the wild-type strain SC5314 with $p < 0.001$. PBS: Phosphate-buffered saline.

burden was approximately 5×10^5 CFUs/g. In contrast, mice infected with the *ngg1* mutants carried negligible amounts of yeast, which were around 10 CFUs/g in kidneys ($p < 0.001$, Figure 5B). In addition, pathological analysis by Periodic Acid Schiff staining was also performed to evaluate the fungal burden in kidneys. As shown in Figure 5C, compared with the dense hyphae in SC5314 group, there was almost no fungal cell observed in PBS group or *ngg1*Δ/Δ infected groups. Collectively, these results demonstrated that *NGG1* is required for the virulence of *C. albicans*, especially to the mammals.

ngg1 mutant displays an altered expression of genes related to energy utilization & mitosis

To gain further insight into the altered phenotypes of the *ngg1* mutant cells, we performed microarray analysis of gene expression. We compared the expression profiles of the wild-type strain and *ngg1-A* during filament formation in Spider media. Three independent biological experiments were carried out and genes showed a greater than twofold change with $q\text{-value} \leq 0.05$ were selected and analyzed. Overall, 772 genes were altered at the transcription level, in which 467 were up-regulated genes and 305 were downregulated (Supplementary Table 3). We performed GO-term analysis to quantify the biological process-related genes. The upregulated genes were mainly involved in the biosynthesis, transportation and metabolism of organic acids, carboxylic acids, fatty acids and amino acids, as well as intracellular oxidation–reduction reaction (Figure 6A). Meanwhile, many of the genes most highly down regulated after *NGG1* deletion were related to the regulation of cell cycle and cellular component organization, the segregation and organization of chromosomes and the regulation of phosphorylation (Figure 6B). Many of the GO-terms were overlapped and the biological processes can be generally categorized as energy utilization and mitosis related.

RT-qPCR was performed to validate genes that displayed transcriptional differences by microarray. These genes included metabolic genes (*LEU42*, *HIS3*, *HXT5*, *HGT12*, *ICL1*, *MSL1* and *PXP2*), oxidation–reduction gene (*SOD1*), mitotic genes (*CLB2*, *SMC2*, *MAD2* and *MTW1*) and genes regulation of cellular component organization (*PEA2*, *EAF3*, *HRR25* and *MUM2*). Similar changes in gene expression were detected by RT-qPCR (Supplementary Figure 3).

Figure 5. The *ngg1* mutants are avirulent in the murine candidiasis model. Mice were inoculated via the tail vein with 1.5×10^6 CFUs of the indicated strains of *Candida albicans*. (A) Survival over 21 days. (B) Fungal burden of the kidneys of mice 2 days after inoculation with the indicated strains. (C) Periodic Acid Schiff stained thin sections of kidneys 2 days after inoculation with different strains. C1: PBS, C2: SC5314, C3: *ngg1-A*, C4: *ngg1-B*. Pathology images were observed with an optical microscope at 400-fold magnification. CFU: Colony-forming unit.

Discussion

In this study, we explored the function of *NGG1* and its role in filamentous growth and virulence of *C. albicans*. *C. albicans* *NGG1* is a 1884 bp DNA sequence located on chromosome 1. The deletion of *NGG1* decreased the acetylation of histone H3, which suggested the HAT function of *C. albicans* Ngg1. Morphological conversion and growth defects of the *ngg1* mutants are important findings of this study. Moreover, the *ngg1* mutants showed dramatically diminished virulence in invertebrates and mice. Microarray analysis revealed the underlying gene expression changes for the serious phenotype variation of the *ngg1* mutants, in which energy utilization and mitosis related genes significantly changed expression.

Taken together the morphological conversion and growth defects of the *ngg1* mutants and the results of the microarray analysis, we postulate that the effects of *NGG1* on the hyphal growth and proliferation of *C. albicans* are mainly caused by mitosis arrest and energy utilization defects. Notably, the knockout of *NGG1* appears to cause altered expression of genes related to the regulation of cell cycle and cellular component organization, the segregation and organization of chromosomes and the regulation of phosphorylation, which means the mitosis of the strain is affected. Interestingly, there is global link between histone acetylation and mitosis, and histone acetylation plays a critical role in marking and activating genes during the mitotic stage of the cell cycle [39].

Also, the microarray results showed that genes involved in the biosynthesis, transportation and metabolism of organic acids, carboxylic acids, fatty acids and amino acids were upregulated in the *ngg1* mutant. Particularly, glyoxylate cycle genes, including *ICL1* (isocitrate lyase) and *MLS1* (malate synthase) were upregulated, which

Figure 6. Significant biological process terms for the differentially expressed genes caused by *NGG1* deletion. GO-term analysis shows the significant biological process terms for the (A) upregulated and (B) downregulated genes caused by *NGG1* deletion in Spider media. Enriched gene ontology terms were calculated from all genes with q-value (%) ≤ 5 and fold change of $\log 2 \leq -1.0$ or ≥ 1.0 .

indicates some degree of energy utilization disorder of the *ngg1* mutant. Meanwhile, the up-regulation of *PXP2* (putative acyl-CoA oxidase; enzyme of fatty acid β -oxidation) further indicates the impaired glucose metabolism of the mutant and implies that, in the mutant, the energy supply is more dependent on the oxidation of fatty acids. Consistent with our results, similar gene expression changes are also found after the deletion of *RTT109*, that is another HAT coding gene [12]. The mechanism by which *NGG1* affects the energy utilization of *C. albicans* is unknown, but researches on *S. cerevisiae* *ADA3* revealed that *ADA3* plays an important role in glucose repression of *S. cerevisiae*, which can give us some clues [25,40].

Energy utilization disorder is a key point affecting the morphological conversion of the strains. During the hyphal growth and biofilm formation of *C. albicans*, much more energy was needed to form stable cell wall and extracellular matrix [3,41–42]. Consistent with this hypothesis, the *ngg1* mutants were sensitive to sodium dodecyl sulphate, caffeine and caspofungin, which implied the damaged cell wall of the mutants [43,44]. Meanwhile, the microarray results showed that the cell wall formation related genes, including many glycosylphosphatidylinositol (GPI) anchored protein coding genes such as *ALS* family, *PGA* family and *IFF* family genes [45] were up-regulated (Supplementary Table 3), which manifested the feedback regulation caused by the damaged cell wall of the mutant.

Apart from the slow metabolism, the intrinsic defect of the *ngg1* mutants also contributes to the reduced filamentation of the strains. After several days on solid Lee's and Spider media, the clone size of the mutants was similar with that of the wild-type strain (Figure 2A), which indicates that the growth of the mutants is not significantly affected by their slow metabolism. However, the wild-type strain produced wrinkled colonies with radial hyphae around, while the *ngg1* mutants just formed smooth colonies (Figure 2A). Moreover, when prolonged the observation time of the strains in liquid-Spider media at 37°C from 3.5 to 8 h, the mutants still developed short filaments or budded cells, while growth curves demonstrated remarkable proliferation of the mutants after 8-h growth at 37°C (Figure 2D). Overall, we believe that both the slow metabolism and the intrinsic defect of the *ngg1* mutants caused the reduced filamentation of the strains.

Importantly, the *ngg1* mutants showed significantly diminished pathogenicity in different infection models. Specifically, 1.5×10^6 CFUs of the *ngg1* mutants were avirulent in mice, which demonstrated the dramatic effects of *NGG1* on the virulence of *C. albicans*. Many studies have shown that the hyphal growth and biofilm formation are critical to the pathogenicity of *C. albicans* thus defects in morphological conversion lead to diminished pathogenicity [3,38,46]. In addition, the possible role of *NGG1* in oxidative stress responses and DNA damage repair is another factor that affects the pathogenicity of *C. albicans*. In this study, we found that the *ngg1* mutants were hypersensitive to various oxidative stresses and the expression of oxidation–reduction-related genes including *SOD1* were upregulated in the *ngg1* mutant, which indicated the disorder of the oxidation–reduction state of the mutant. Oxidative stress responses and DNA damage repair machinery are critical for *C. albicans* to escape from the host innate immunity, as macrophages and neutrophils can produce reactive oxygen species to cause fatal oxidative damage and DNA damage to the strains [47,48]. Moreover, *C. albicans* *ngg1* mutants were hypersensitive to the genotoxic drug 5-fluorocytosine, which implied the decreased ability of DNA damage repair of the strains.

The hypersensitivity of the *ngg1* mutants to azoles, including fluconazole and ketoconazole, is an interesting finding. As ergosterol biosynthesis plays a significant role in the sensitivity of *C. albicans* to azoles [49,50], we analyzed the expression of ergosterol biosynthesis genes from the microarray data. However, no significant change was observed in the expression of all of the ergosterol biosynthesis genes that play important roles in the sensitivity of *Candida* to antifungal agents (including *ERG1*, *ERG3*, *ERG7*, *ERG9*, *ERG11* and *ERG25*) [51]. Additional studies, including direct evaluation of ergosterol content may help elucidate the relationship between the hypersensitivity of the mutants to azoles and their biosynthesis of ergosterol.

Recent studies suggest that histone modification inhibitory agents could provide a promising approach for the treatment of fungal infections. For example, genetic or chemical inhibition of the HAT Rtt109, which target H3K56, could reduce *C. albicans* virulence in mice [52]. Meanwhile, the inhibition of histone deacetylase activity of Hos2 by trichostatin A leads to reduced virulence of *C. albicans* [53]. Moreover, a specific Hos2 inhibitor termed MGCD290 has entered clinical trials as potential antifungal drug [54]. Interestingly, Hos2 shows high conservation between *C. albicans* and humans, with Protein BLAST *E*-value of 7×10^{-174} [55]. As noted above, the *C. albicans* Ngg1 and human Tada3 have lower similarity with an *E*-value of 9×10^{-12} . Therefore, we favor the hypothesis that Ngg1 can be considered as a potential antifungal target.

A limitation of this study is that we were unable to construct the reconstituted strain, in spite of multiple efforts over a 12-month period. During the transformation, we found that the mutant cells were very easy to die (data not shown). Of note is that, the deletion of HATs affects the transformation of yeast [31]. Moreover, the *S. cerevisiae*

ada3 mutant showed abnormal vitality and cannot even grow at 37°C [26]. Considering the serious phenotype variation of the *ngg1* mutants, we suspect that cells were unable to survive through transformation.

Conclusion & future perspective

In summary, this study verified the HAT activity of *C. albicans* Ngg1 and revealed its vital role in morphological conversion and virulence. The deletion of *NGG1* caused extensive gene expression changes of *C. albicans*, in which energy utilization defects and mitosis arrest are the key factors affecting the morphological conversion and pathogenicity. Considering the low similarity between *C. albicans* Ngg1 and human Tada3, these results are very encouraging and suggest that Ngg1 could be a site for targeting antifungals.

Summary points

Background

- Despite extensive analysis of human Tada3 and *Saccharomyces cerevisiae* Ada3, their homolog in *Candida albicans*, Ngg1, is not well understood.

Results

- Deletion of *NGG1* drastically decreased the acetylation of histone H3 in *C. albicans*, which verified the HAT activity of *C. albicans* Ngg1.
- Two different *ngg1* mutants showed significant defects in morphological conversion and growth.
- Pathogenicity of *C. albicans* is diminished in the absence of *NGG1*.
- The *ngg1* mutants displayed an altered expression of genes related to energy utilization and mitosis.

Conclusion

- *NGG1* is an important HAT in *C. albicans* and plays an important role in morphological conversion and virulence of *C. albicans*.

Supplementary data

To view the supplementary data that accompany this paper please visit the journal website at: www.futuremedicine.com/doi/suppl/10.2217/fmb-2017-0084.

Acknowledgements

The authors thank J Morschhauser (Institut für Molekulare Infektionsbiologie, University of Würzburg, Germany) for providing the plasmid pSFS2.

Financial and competing interests disclosure

This work was funded by the National Natural Science Foundation of China 81501730 to DD Li, 81573724 to Y Sun. All of the transcription data were deposited at the NCBI Gene Expression Omnibus (GEO) database with accession number GSE93200. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Ethical conduct of research

All murine protocols were approved by the Rhode Island Hospital Institutional Animal Care and Use Committee.

References

Papers of special note have been highlighted as: • of interest; •• of considerable interest

1. Kim J, Sudbery P. *Candida albicans*, a major human fungal pathogen. *J. Microbiol.* 49(2), 171–177 (2011).
2. Wisplinghoff H, Ebbers J, Geurtz L *et al.* Nosocomial bloodstream infections due to *Candida* spp. in the USA: species distribution, clinical features and antifungal susceptibilities. *Int. J. Antimicrob. Agents* 43(1), 78–81 (2014).
3. Sudbery PE. Growth of *Candida albicans* hyphae. *Nat. Rev. Microbiol.* 9(10), 737–748 (2011).
4. Odds FC. Morphogenesis in *Candida albicans*. *Crit. Rev. Microbiol.* 12(1), 45–93 (1985).
5. Stoldt VR, Sonneborn A, Leuker CE, Ernst JF. Efg1p, an essential regulator of morphogenesis of the human pathogen *Candida albicans*, is a member of a conserved class of bHLH proteins regulating morphogenetic processes in fungi. *EMBO J.* 16(8), 1982–1991 (1997).

6. Lo HJ, Kohler JR, Didomenico B, Loebenberg D, Cacciapuoli A, Fink GR. Nonfilamentous *C. albicans* mutants are avirulent. *Cell* 90(5), 939–949 (1997).
7. Braun BR, Johnson AD. TUP1, CPH1 and EFG1 make independent contributions to filamentation in *Candida albicans*. *Genetics* 155(1), 57–67 (2000).
8. Liu H, Kohler J, Fink GR. Suppression of hyphal formation in *Candida albicans* by mutation of a STE12 homolog. *Science* 266(5191), 1723–1726 (1994).
9. Leberer E, Marcus D, Broadbent ID *et al.* Signal transduction through homologs of the Ste20p and Ste7p protein kinases can trigger hyphal formation in the pathogenic fungus *Candida albicans*. *Proc. Natl Acad. Sci. USA* 93(23), 13217–13222 (1996).
10. Lu Y, Su C, Mao X, Raniga PP, Liu H, Chen J. Efg1-mediated recruitment of NuA4 to promoters is required for hypha-specific Swi/Snf binding and activation in *Candida albicans*. *Mol. Biol. Cell* 19(10), 4260–4272 (2008).
11. Chang P, Fan X, Chen J. Function and subcellular localization of Gcn5, a histone acetyltransferase in *Candida albicans*. *Fungal Genet. Biol.* 81, 132–141 (2015).
12. Lopes Da Rosa J, Boyartchuk VL, Zhu LJ, Kaufman PD. Histone acetyltransferase Rtt109 is required for *Candida albicans* pathogenesis. *Proc. Natl Acad. Sci. USA* 107(4), 1594–1599 (2010).
- **Important research about the fungal-specific histone acetyltransferase Rtt109 in *Candida albicans*. Indicated that Rtt109 is particularly important for fungal pathogenicity and is suitable to be a unique target for therapeutic antifungal compounds.**
13. Nag A, Germaniuk-Kurowska A, Dimri M *et al.* An essential role of human Ada3 in p53 acetylation. *J. Biol. Chem.* 282(12), 8812–8820 (2007).
14. Balasubramanian R, Pray-Grant MG, Selleck W, Grant PA, Tan S. Role of the Ada2 and Ada3 transcriptional coactivators in histone acetylation. *J. Biol. Chem.* 277(10), 7989–7995 (2002).
15. Pray-Grant MG, Schieltz D, McMahon SJ *et al.* The novel SLIK histone acetyltransferase complex functions in the yeast retrograde response pathway. *Mol. Cell. Biol.* 22(24), 8774–8786 (2002).
16. Sterner DE, Berger SL. Acetylation of histones and transcription-related factors. *Microbiol. Mol. Biol. Rev.* 64(2), 435–459 (2000).
17. Candida Genome Database. <http://www.candidagenome.org/>
18. Pukkila-Worley R, Peleg AY, Tampakakis E, Mylonakis E. *Candida albicans* hyphal formation and virulence assessed using a *Caenorhabditis elegans* infection model. *Eukaryot. Cell* 8(11), 1750–1758 (2009).
19. Reuss O, Vik A, Kolter R, Morschhauser J. The SAT1 flipper, an optimized tool for gene disruption in *Candida albicans*. *Gene* 341, 119–127 (2004).
- **Provides detailed information on the gene knockout method which is widely used in *C. albicans*.**
20. Walther A, Wendland J. An improved transformation protocol for the human fungal pathogen *Candida albicans*. *Curr. Genet.* 42(6), 339–343 (2003).
21. Nobile CJ, Andes DR, Nett JE *et al.* Critical role of Bcr1-dependent adhesins in *C. albicans* biofilm formation *in vitro* and *in vivo*. *PLoS Pathog.* 2(7), e63 (2006).
22. Fuchs BB, O'Brien E, Khoury JB, Mylonakis E. Methods for using *Galleria mellonella* as a model host to study fungal pathogenesis. *Virulence* 1(6), 475–482 (2010).
23. Breger J, Fuchs BB, Aperis G, Moy TI, Ausubel FM, Mylonakis E. Antifungal chemical compounds identified using a *C. elegans* pathogenicity assay. *PLoS Pathog.* 3(2), e18 (2007).
24. Liang RM, Yong XL, Jiang YP *et al.* 2-amino-nonyl-6-methoxyl-tetralin muriate activity against *Candida albicans* augments endogenous reactive oxygen species production – a microarray analysis study. *FEBS J.* 278(7), 1075–1085 (2011).
25. Brandl CJ, Martens JA, Margalio A *et al.* Structure/functional properties of the yeast dual regulator protein NGG1 that are required for glucose repression. *J. Biol. Chem.* 271(16), 9298–9306 (1996).
26. Pina B, Berger S, Marcus GA, Silverman N, Agapite J, Guarente L. ADA3: a gene, identified by resistance to GAL4-VP16, with properties similar to and different from those of ADA2. *Mol. Cell Biol.* 13(10), 5981–5989 (1993).
- **First article focused on ADA3 in *Saccharomyces cerevisiae*, found that the *ada3* mutant cannot grow at 37°C on any media.**
27. Marchler-Bauer A, Bo Y, Han L *et al.* CDD/SPARCLE: functional classification of proteins via subfamily domain architectures. *Nucleic Acids Res.* 45(D1), D200–D203 (2017).
28. Kawai S, Hashimoto W, Murata K. Transformation of *Saccharomyces cerevisiae* and other fungi: methods and possible underlying mechanism. *Bioeng. Bugs* 1(6), 395–403 (2010).
- **Provides good overview of the yeast transformation methods and discusses some common modifications to improve transformation efficiency.**
29. de Backer MD, Maes D, Vandoninck S, Logghe M, Contreras R, Luyten WH. Transformation of *Candida albicans* by electroporation. *Yeast* 15(15), 1609–1618 (1999).
30. Thompson JR, Register E, Curotto J, Kurtz M, Kelly R. An improved protocol for the preparation of yeast cells for transformation by electroporation. *Yeast* 14(6), 565–571 (1998).

31. Soltani J, van Heusden GP, Hooykaas PJ. Deletion of host histone acetyltransferases and deacetylases strongly affects *Agrobacterium*-mediated transformation of *Saccharomyces cerevisiae*. *FEMS Microbiol. Lett.* 298(2), 228–233 (2009).
32. Arbour M, Epp E, Hogues H *et al.* Widespread occurrence of chromosomal aneuploidy following the routine production of *Candida albicans* mutants. *FEMS Yeast Res.* 9(7), 1070–1077 (2009).
33. Bouchonville K, Forche A, Tang KE, Selmecki A, Berman J. Aneuploid chromosomes are highly unstable during DNA transformation of *Candida albicans*. *Eukaryot. Cell* 8(10), 1554–1566 (2009).
34. Noble SM, French S, Kohn LA, Chen V, Johnson AD. Systematic screens of a *Candida albicans* homozygous deletion library decouple morphogenetic switching and pathogenicity. *Nat. Genet.* 42(7), 590–598 (2010).
35. Dudley AM, Janse DM, Tanay A, Shamir R, Church GM. A global view of pleiotropy and phenotypically derived gene function in yeast. *Mol. Syst. Biol.* 1, 2005.0001 (2005).
36. Yeater KM, Chandra J, Cheng G *et al.* Temporal analysis of *Candida albicans* gene expression during biofilm development. *Microbiology* 153(Pt 8), 2373–2385 (2007).
37. Desai JV, Mitchell AP. *Candida albicans* biofilm development and its genetic control. *Microbiol. Spectr.* 3(3), doi:10.1128/microbiolspec.MB-0005-2014 (2015) (Epub ahead of print).
38. Hirota K, Yumoto H, Sapaar B, Matsuo T, Ichikawa T, Miyake Y. Pathogenic factors in *Candida* biofilm-related infectious diseases. *J. Appl. Microbiol.* 122(2), 321–330 (2017).
39. Valls E, Sanchez-Molina S, Martinez-Balbas MA. Role of histone modifications in marking and activating genes through mitosis. *J. Biol. Chem.* 280(52), 42592–42600 (2005).
40. Brandl CJ, Furlanetto AM, Martens JA, Hamilton KS. Characterization of NGG1, a novel yeast gene required for glucose repression of GAL4p-regulated transcription. *EMBO J.* 12(13), 5255–5265 (1993).
41. Land GA, McDonald WC, Stjernholm RL, Friedman TL. Factors affecting filamentation in *Candida albicans*: relationship of the uptake and distribution of proline to morphogenesis. *Infect. Immun.* 11(5), 1014–1023 (1975).
42. Li DD, Wang Y, Dai BD *et al.* ECM17-dependent methionine/cysteine biosynthesis contributes to biofilm formation in *Candida albicans*. *Fungal Genet. Biol.* 51, 50–59 (2013).
43. Kofla G, Ruhnke M. Pharmacology and metabolism of anidulafungin, caspofungin and micafungin in the treatment of invasive candidosis: review of the literature. *Eur. J. Med. Res.* 16(4), 159–166 (2011).
44. Plaine A, Walker L, Da Costa G *et al.* Functional analysis of *Candida albicans* GPI-anchored proteins: roles in cell wall integrity and caspofungin sensitivity. *Fungal Genet. Biol.* 45(10), 1404–1414 (2008).
45. Richard ML, Plaine A. Comprehensive analysis of glycosylphosphatidylinositol-anchored proteins in *Candida albicans*. *Eukaryot. Cell* 6(2), 119–133 (2007).
46. Tsui C, Kong EF, Jabra-Rizk MA. Pathogenesis of *Candida albicans* biofilm. *Pathog. Dis.* 74(4), ftw018 (2016).
47. Dantas Ada S, Day A, Ikeh M, Kos I, Achan B, Quinn J. Oxidative stress responses in the human fungal pathogen, *Candida albicans*. *Biomolecules* 5(1), 142–165 (2015).
48. Lorenz MC, Bender JA, Fink GR. Transcriptional response of *Candida albicans* upon internalization by macrophages. *Eukaryot. Cell* 3(5), 1076–1087 (2004).
49. Morschhauser J. The development of fluconazole resistance in *Candida albicans* – an example of microevolution of a fungal pathogen. *J. Microbiol.* 54(3), 192–201 (2016).
50. Akins RA. An update on antifungal targets and mechanisms of resistance in *Candida albicans*. *Med. Mycol.* 43(4), 285–318 (2005).
51. Borecka-Melkusova S, Moran GP, Sullivan DJ, Kucharikova S, Chorvat D Jr, Bujdakova H. The expression of genes involved in the ergosterol biosynthesis pathway in *Candida albicans* and *Candida dubliniensis* biofilms exposed to fluconazole. *Mycoses* 52(2), 118–128 (2009).
52. Wurtele H, Tsao S, Lepine G *et al.* Modulation of histone H3 lysine 56 acetylation as an antifungal therapeutic strategy. *Nat. Med.* 16(7), 774–780 (2010).
53. Hnisz D, Majer O, Frohner IE, Komnenovic V, Kuchler K. The Set3/Hos2 histone deacetylase complex attenuates cAMP/PKA signaling to regulate morphogenesis and virulence of *Candida albicans*. *PLoS Pathog.* 6(5), e1000889 (2010).
54. Pfaller MA, Messer SA, Georgopapadakou N, Martell LA, Besterman JM, Diekema DJ. Activity of MGCD290, a Hos2 histone deacetylase inhibitor, in combination with azole antifungals against opportunistic fungal pathogens. *J. Clin. Microbiol.* 47(12), 3797–3804 (2009).
55. Tscherner M, Stappler E, Hnisz D, Kuchler K. The histone acetyltransferase Hat1 facilitates DNA damage repair and morphogenesis in *Candida albicans*. *Mol. Microbiol.* 86(5), 1197–1214 (2012).

Comparative genomics of type 1 IncC plasmids from China

Lizhi Ma^{‡,1}, Zhe Yin^{‡,2}, Defu Zhang^{2,4}, Zhe Zhan², Qian Wang¹, Xiongbo Duan³, Huixia Gao³, Quanhui Liang², Yuzong Zhao^{2,4}, Jiao Feng², Yachao Zhao², Yigang Tong², Erhei Dai^{*,3} & Dongsheng Zhou^{**,2}

¹Department of Emergency Medicine, General Hospital of Chinese People's Armed Police Forces, Beijing 100039, China

²State Key Laboratory of Pathogen & Biosecurity, Beijing Institute of Microbiology & Epidemiology, Beijing 100071, China

³Department of Laboratory Medicine, The Fifth Hospital of Shijiazhuang, Hebei Medical University, Shijiazhuang, Hebei 050021, China

⁴College of Food Science & Project Engineering, Bohai University, Jinzhou 121013, China

* Author for correspondence: daieh2008@126.com

** Author for correspondence: dongshengzhou1977@gmail.com

‡ Authors contributed equally

Aim: This study dealt with genomic characterization of type 1 IncC resistance plasmids, capable of spreading across taxonomic borders, from China. **Materials & methods:** p112298-tetA was sequenced and compared with type 1 IncC reference plasmid pR148 and two available sequenced type 1 IncC plasmids pH536-NDM and pVAS3-1 from China. **Results:** These plasmids contained one or more exogenous resistance islands, which included the ARI-A islands, the ARI-B islands, the *ISEcp1-bla_{CMY}* units and the *bla_{KPC-2}* region and were inserted at various sites in the IncC backbone and thus represented three distinct lineages. **Conclusion:** Complex rearrangement and homologous recombination events have occurred during evolution of p112298-tetA, making it significantly differ modularly from the other three plasmids with respect to both plasmid backbone and exogenous resistance regions.

First draft submitted: 25 April 2017; Accepted for publication: 28 July 2017; Published online: 15 November 2017

Keywords: *Citrobacter freundii* • multidrug resistance • p112298-tetA • type 1 IncC plasmid

IncA and IncC plasmids with very similar exclusion and phage sensitivity were revealed and combined into the IncA/C group more than 40 years ago. IncA/C was recently divided into two types, IncA/C₁ (reference plasmid RA1 [1]) and IncA/C₂ [2], which corresponded to IncA and IncC and have very separate evolutionary stories with significant nucleotide divergence between their backbones [3]. IncA and IncC plasmids are actually compatible [4] and thus, IncA/C is an incorrect designation and should be separated again into IncA and IncC [5].

IncC plasmids have been identified worldwide from environmental, animal and clinical isolates of various species of Enterobacteriaceae [3,6], *Vibrio* [7–9], and *Aeromonas* [10,11] and also *Photobacterium damsela* [12], indicating their ability to spread across taxonomic borders. Plasmid backbone comparison have further divided IncC into two subtypes, type 1 (reference plasmid pR148 [10]) and type 2 (reference plasmid pR55 [13]), and these two lineages have diverged a long time ago and can be distinguished by two genes of substitution (*orf1832* in type 1 or *orf1847* in type 2 and *rhs1* in type 1 or *rhs2* in type 2) and two small intergenic insertions (i1 and i2, 428 and 462 bp in length, respectively, found in type 2 rather than type 1) [14].

Three kinds of exogenous resistance island, including an *ISEcp1-bla_{CMY}* transposition unit, ARI-A and ARI-B, are often found to integrate into the IncC backbones, making them to be important vehicles of spread of multidrug resistance [3,6]. Most of type 1 IncC plasmids possess the *ISEcp1-bla_{CMY}* unit and ARI-A and meanwhile some of them carry ARI-B [3]. ARI-B (but not the *ISEcp1-bla_{CMY}* unit and ARI-A) is found in type 2 IncC plasmids but it is not always present and type 2 IncC plasmids often carry additional type(s) of resistance island besides ARI-B [3]. Most of the available evidences point to a single acquisition of ARI-A in an ancestral strain, followed by evolution *in situ* in the type 1 lineage, while ARI-B may have been acquired independently by type 1 and type 2 lineages.

Table 1. Type 1 and type 2 IncC plasmids from China.

Plasmid	Host bacterium	Accession number	IncC subtype	<i>orf1832/orf1847</i>	<i>rhs1/rhs2</i>	i1	i2	Location
pR148	<i>Aeromonas hydrophila</i>	JX141473	Type 1 reference	<i>orf1832</i>	<i>rhs1</i>	-	-	Thailand
pHS36-NDM	<i>Salmonella enterica</i> serovar Stanley	KU726616	Type 1	<i>orf1832</i>	Δ <i>rhs1</i>	-	-	China
pVAS3-1	<i>Vibrio alginolyticus</i>	KU160531	Type 1	<i>orf1832</i>	<i>rhs1</i>	-	-	China
p112298-tetA	<i>Citrobacter freundii</i>	KY986974	Type 1	<i>orf1832</i>	<i>rhs1</i>	-	+	China
pR55	<i>Klebsiella pneumoniae</i>	JQ010984	Type 2 reference	<i>orf1847</i>	<i>rhs2</i>	+	+	France
pVC1447	<i>Vibrio cholerae</i>	KM083064	Type 2	<i>orf1847</i>	<i>rhs2</i>	+	+	China
pVC1307	<i>Vibrio cholerae</i>	KJ817377	Type 2	<i>orf1847</i>	<i>rhs2</i>	+	+	China
pKP36-2	<i>Klebsiella pneumoniae</i>	CP017387	Type 2	<i>orf1847</i>	Δ <i>rhs2</i>	+	+	China
pSCEC2	<i>Escherichia coli</i>	KF152885	Type 2	<i>orf1847</i>	<i>rhs2</i>	-	+	China
pKPHS3	<i>Klebsiella pneumoniae</i>	CP003225	Type 2	-	<i>rhs2</i>	+	-	China
pNDM1.SZ1	<i>Enterobacter cloacae</i>	KU302801	NA	-	-	-	-	China
pTR2	<i>Klebsiella pneumoniae</i>	KJ187752	NA	-	-	-	-	China

Type 1: presence of *orf1832* and *rhs1*; Type 2: presence of *orf1847*, *rhs2*, i1 and i2.
pR148, HS36-NDM, pVAS3-1 and p112298-tetA were included in the comparative genomics analysis in this work.
+: Presence; -: Absence; NA: Not applicable; pKP36-2: Plasmid 2 of *K. pneumoniae* KP36.

We recently reported two fully sequenced multidrug resistant (MDR) plasmids, p112298-KPC (could not be assigned into any of the known incompatibility groups) [15] and p112298-catA (belonging to IncHI5) [16], from a clinical *Citrobacter freundii* isolate. This follow-up study determined the complete sequence of the third MDR plasmid p112298-tetA (belonging to type 1 IncC) from this isolate. Detailed genomic comparison of the three available sequenced type 1 IncC plasmids from China, including p112298-tetA and the reference plasmid pR148 [10] indicated significant modular differences in both plasmid backbones and accessory regions and thus represented distinct lineages. Data presented here provide a deeper insight into diversification and evolutionary history of type 1 IncC plasmids.

Materials & methods

C. freundii 112298 was isolated in 2013 from a midstream urine specimen from a patient with septic shock in a Chinese teaching hospital [15]. Plasmids were transferred in an attempt from the 112298 isolate into *Escherichia coli* TOP10 via electroporation, followed by selection of a *tetA*(A)- and *bla*_{KPC}-positive electroporant [15]. Activity of Ambler class A/B/D carbapenemases in bacterial cell extracts was determined by a modified CarbaNP test [15]. Bacterial antimicrobial susceptibility was tested by the broth dilution method and interpreted as per CLSI guidelines [17].

To determine the complete sequences of the plasmids in a wild-type strain, genomic DNA of the 112,298 isolate was sequenced from a mate pair library with average insert size of 5000 bp, using a MiSeq sequencer (Illumina, CA, USA). In order to get complete plasmid sequences, DNA contigs were assembled using Newbler 2.6 [18] and gaps between contigs were filled using a combination of PCR and Sanger sequencing using an ABI 3730 Sequencer. Open reading frames and pseudogenes were predicted using RAST 2.0 [19] combined with BLASTP/BLASTN [20] searches against the UniProtKB/Swiss-Prot [21] and RefSeq [22] databases. Annotation of resistance genes, mobile elements and other features was carried out using CARD [23], ResFinder [24], ISfinder [25] and INTEGRALL [26]. Multiple and pairwise sequence comparisons were performed using MUSCLE 3.8.31 [27] and BLASTN, respectively. Gene organization diagrams were drawn in Inkscape 0.48.1.

The complete sequence of p112298-tetA was submitted to GenBank under accession number KY986974.

Results

Overview of sequenced type 1 IncC plasmids from China

The complete nucleotide sequence of p112298-tetA determined herein was 190,310 bp in length with a mean G + C content of 52.5% and a total of 235 predicted open reading frames (ORFs; Supplementary Figure 1). All the ten fully sequenced IncC plasmids from China, including p112298-tetA, available in the public sequence databases were collected (Table 1). Based on presence or absence of *orf1832/orf1847*, *rhs1/rhs2*, i1 and i2, three of these plasmids including p112298-tetA were assigned into type 1 and five were recognized as type 2, while the

Table 2. Major features of plasmids analyzed.

Category	Plasmids			
	pR148	pHS36-NDM	pVAS3-1	p112298-tetA
Host bacterium	<i>Aeromonas hydrophila</i>	<i>Salmonella enterica</i> serovar Stanley	<i>Vibrio alginolyticus</i>	<i>Citrobacter freundii</i>
Total length (bp)	165,906	137,952	175,254	190,310
Mean G + C content (%) of the whole sequence	52.5	52.0	52.3	52.5
Total number of ORFs	198	170	209	234
Length (bp) of the backbone	127,803	113,439	117,080	116,810
Mean G + C content (%) of the backbone	51.2	51.2	51.2	51.3
Resistance islands				
ARI-A	Tn6358	The Tn1696-In27-Tn125 region	Tn6359	The Tn1696-In1212 region
ARI-B	–	–	The <i>aphA1</i> -In641- <i>floR</i> - <i>tetAR</i> - <i>strAB</i> - <i>sul2</i> region	The <i>floR</i> - <i>tetAR</i> - <i>strAB</i> - <i>sul2</i> - <i>bla</i> _{CMY-2} region
The <i>ISEcp1</i> - <i>bla</i> _{CMY} unit	–	The <i>ISEcp1</i> - <i>bla</i> _{CMY-6} unit	The <i>ISEcp1</i> - <i>bla</i> _{CMY-2} unit	†
Additional	–	–	–	The <i>bla</i> _{KPC-2} region

† Segmented and merged into the *floR*-*tetAR*-*strAB*-*sul2*-*bla*_{CMY-2} region and the *bla*_{KPC-2} region.

–: Absence; ORF: Open reading frame.

remaining two could not be defined as either type 1 or type 2 because none of the above signature sequences could be identified (Table 1). In addition to *rhs1*, *orf1832* and ARI-A characteristic of type 1, p112298-tetA still carried the *i2* insert characteristic of type 2, which might result from homologous recombination between type 1 and type 2 backbones.

The three sequenced type 1 IncC plasmids pHS36-NDM [28], pVAS3-1 [7] and p112298-tetA (this study) from China, together with the type 1 IncC reference plasmid pR148 [10], were included in a genomic comparison. The modular structure of each plasmid was discriminated as the IncC backbone with insertion of one or more separate accessory modules (Supplementary Figures 1 & 2). These accessory modules were further dissected as resistance islands including the *ISEcp1*-*bla*_{CMY} unit, ARI-A and ARI-B and the *bla*_{KPC-2} regions and each plasmid contained one or more of these resistance islands (Table 2), which were inserted at various sites in the plasmid backbones (see below). The drug resistance genes identified were exclusively located within the resistance islands and these four plasmids displayed dramatically different profiles of resistance genes (Table 3).

Pairwise sequence comparison using BLAST indicated that these four plasmids had 99% nucleotide identity over >87% of their backbone sequences (Supplementary Table 1), denoting considerable modular differences in the backbones of these four plasmids. Nevertheless, the master IncC backbone genes or gene loci [3], involved in replication (*repA*), conjugative transfer (*tra1* and *tra2*), partitioning and stability (*parAB*, *parM* and *stbA*) and DNA metabolism (*ssb*-*bet*-*exo*, *ter*-*kfrA*, *int*-*yacC*, *nuc*, *uvrD* and *pri*), were shared by these four plasmids.

Major modular differences in type 1 IncC plasmids from China

Linear sequence comparison revealed at least six major modular differences in pR148, pHS36-NDM, pVAS3-1 and p112298-tetA (Supplementary Figure 1). First, the ARI-B islands were inserted at a site (Figure 1A) downstream of *orf498* in pVAS3-1 and p112298-tetA, but not found in pR148 and pHS36-NDM (Supplementary Figure 2). The *dcm1* region, which was composed of 16 IncC backbone genes (from *orf933* to *orf450*) as observed in pR148 and pHS36-NDM, was lost due to an IS26-mediated deletion event arising from within ARI-B. Second, the *ISEcp1*-*bla*_{CMY-6}/*bla*_{CMY-2} units were inserted at a site (Figure 1B) between *traA* and *orf1832* in pHS36-NDM and pVAS3-1, respectively, but none of the *ISEcp1*-*bla*_{CMY} units were found in pR148 (Supplementary Figure 2). The *ISEcp1*-*bla*_{CMY-2} unit was segmented into two separate parts in p112298-tetA (see below). Third, the *dcm3* region, which was composed of a partial region of *orf540* together with 15 consecutive IncC backbone genes (from *orf501* to *orf2454*) as observed in pR148, pVAS3-1 and p112298-tetA, were not identified in pHS36-NDM. Fourth, the ARI-A islands were inserted at a site (Figure 1C) within *orf240* in all these four plasmids, splitting *orf240* into two separate parts *orf240-5'* and *orf240-3'* in pR148 (Supplementary Figure 2). Compared with the prototype structure *orf240-3'*-*orf375*-*orf726*-*rhs1* (designated the *rhs1* region; 5965 bp in length) in pR148, *rhs1* was pseudogenized

Table 3. Drug resistance genes in plasmids analyzed.

Plasmid	Resistance marker	Resistance phenotype	Nucleotide position	Region located
pR148	<i>qacH4</i>	Quaternary ammonium compound resistance	106328.106660	Tn6358 (ARI-A)
	<i>bla_{OXA-10}</i>	β -lactam resistance	106906.107706	
	<i>aadA1e</i>	Aminoglycoside resistance	107723.108514	
	<i>qacED1</i>	Quaternary ammonium compound resistance	110516.110863	
	<i>sul1</i>	Sulfonamide resistance	110857.111696	
	<i>catA2</i>	Phenicol resistance	114393.115034	
	<i>sul1</i>	Sulfonamide resistance	115981.116820	
	<i>chrA</i>	Chromate resistance	118635.119840	
	<i>tetA(A)</i>	Tetracycline resistance	125446.126645	
	The <i>mer</i> locus	Mercuric resistance	132315.136582	
pHS36-NDM	<i>dfrA12</i>	Trimethoprim resistance	99867.100364	The Tn1696-In27-Tn125 region (ARI-A)
	<i>aadA2</i>	Aminoglycoside resistance	100784.101563	
	<i>qacED1</i>	Quaternary ammonium compound resistance	101727.102074	
	<i>sul1</i>	Sulfonamide resistance	102068.102907	
	<i>bla_{NDM-1}</i>	Carbapenem resistance	107326.108138	
	<i>ble_{MBL}</i>	Bleomycin resistance	108142.108507	
	<i>bla_{CMY-6}</i>	β -lactam resistance	56967.58112	
	<i>sugE</i>	Quaternary ammonium compound resistance	58736.59176	
pVAS3-1	<i>qnrVC4</i>	Quinolone resistance	128681.129337	The <i>aphA1</i> -In641- <i>floR</i> - <i>tetAR</i> - <i>strAB</i> - <i>sul2</i> region (ARI-B)
	<i>qacF</i>	Quaternary ammonium compound resistance	129594.129926	
	<i>aacA4'-17</i>	Aminoglycoside resistance	130015.130569	
	<i>cmlA1g</i>	Phenicol resistance	130784.132043	
	<i>bla_{OXA-10}</i>	β -lactam resistance	132308.133108	
	<i>aadA1e</i>	Aminoglycoside resistance	133125.133916	
	<i>dfrA14b</i>	Trimethoprim resistance	133942.134424	
	<i>mph(A)</i>	Macrolide resistance	138129.139034	
	The <i>mer</i> locus	Mercuric resistance	140659.145912	
	<i>aphA1</i>	Aminoglycoside resistance	18862.19677	
	<i>sul3</i>	Sulfonamide resistance	23672.24463	
	<i>qacH2</i>	Quaternary ammonium compound resistance	25643.25975	
	<i>aadA1a</i>	Aminoglycoside resistance	26145.26936	
	<i>cmlA1</i>	Phenicol resistance	27029.28288	
	<i>aadA2</i>	Aminoglycosideresistance	28550.29329	
	<i>estX3</i>	Aminoglycoside resistance	30103.30945	
	<i>floR</i>	Phenicol resistance	39543.40757	
	<i>tetA(A)</i>	Tetracycline resistance	41357.42631	
	<i>strB</i>	Aminoglycoside resistance	43892.44728	
	<i>strA</i>	Aminoglycoside resistance	44728.45531	
	<i>sul2</i>	Sulfonamide resistance	45592.46407	
	<i>bla_{CMY-2}</i>	β -lactam resistance	74748.75893	
	<i>sugE</i>	Quaternary ammonium compound resistance	76517.76957	
	<i>aacA7</i>	Aminoglycoside resistance	91328.91786	
	<i>qacED1</i>	Quaternary ammonium compound resistance	91916.92263	
	<i>sul1</i>	Sulfonamide resistance	92257.93096	
p112298-tetA	<i>aacA7</i>	Aminoglycoside resistance	91328.91786	The Tn1696-In1212 region (ARI-A)
	<i>qacED1</i>	Quaternary ammonium compound resistance	91916.92263	
	<i>sul1</i>	Sulfonamide resistance	92257.93096	

Table 3. Drug resistance genes in plasmids analyzed (cont.).

Plasmid	Resistance marker	Resistance phenotype	Nucleotide position	Region located
	<i>floR</i>	Phenicol resistance	24388.25602	The <i>floR</i> - <i>tetAR</i> - <i>strAB</i> - <i>sul2</i> - <i>bla</i> _{CMY-2} region (ARI-B)
	<i>tetA(A)</i>	Tetracycline resistance	26202.27476	
	<i>strB</i>	Aminoglycoside resistance	28737.29573	
	<i>strA</i>	Aminoglycoside resistance	29573.30376	
	<i>sul2</i>	Sulfonamide resistance	30437.31252	
	<i>bla</i> _{CMY-2}	β-lactam resistance	37573.38718	
	<i>sugE</i>	Quaternary ammonium compound resistance	39342.39782	
	<i>bla</i> _{KPC-2}	β-lactam resistance	159597.160478	The <i>bla</i> _{KPC-2} region

Figure 1. Sites of insertion of resistance islands. Genes are denoted by arrows. Genes, mobile elements and other features are colored based on function classification. Shading denotes regions of homology (>95% nucleotide identity). The accession numbers of pR148 [10], pHS36-NDM [28] and pVAS3-1 [7] are JX141473, KU726616 and KU160531, respectively.

due to 1-bp in-frame deletion in pVAS3-1, a 3473-bp region (composed of *orf240-3'-orf375-orf726* and a 5'-end fragment of *rhs1*) was deleted in pHS36-NDM, while the 788-bp *orf240-3'-orf375* region was deleted in p112298-tetA (Figure 1C). Various types of genetic modification in the *rhs1* regions immediately downstream of ARI-A occurred along with and/or after insertion of ARI-A and notably, those found in pHS36-NDM and p112298-tetA were characteristic of two corresponding sublineages of type 1 IncC plasmid being reported outside of Australia and USA. Fifth, the *bla*_{KPC-2} region was inserted at a site upstream of *orf726* in p112298-tetA. Sixth, the *orf312* to *tnpA*_{ISEcp1} region was translocated in p112298-tetA relative to pHS36-NDM and pVAS3-1 (Supplementary Figure 2) and this translocation split the *ISEcp1-bla*_{CMY-2} unit into two segments (one emerged as a part of ARI-B and another as a portion of the *bla*_{KPC-2} region) and the breaking point was located within *tnpA*_{ISEcp1} (Figure 1B).

The ARI-A islands

The ARI-A islands manifested as Tn6358, the Tn1696-In27-Tn125 region, Tn6359 and the Tn1696-In1212 region in pR148, pHS36-NDM, pVAS3-1 and p112298-tetA, respectively (Figure 2), all of which were the derivatives of Tn1696, a unit transposon belonging to the Tn21 subgroup of the Tn3 family [29]. Tn1696 was generated from insertion of class 1 integron In4 at a site within the resolution (*res*) site of a core structure IRL (inverted repeat left)-*tnpA* (transposase)-*tnpR* (resolvase)-*res*-*mer* (mercury resistance locus)-IRR (inverted repeat

Figure 2. The ARI-A islands and comparison with related regions. Genes are denoted by arrows. Genes, mobile elements and other features are colored based on their functional classification. Shading denotes regions of homology (nucleotide identity >95%). Numbers in brackets indicate the nucleotide positions within the corresponding plasmids. Tn1721 [31], Tn1696 [29] and Tn125 [30] are prototype mobile elements as the references for sequence comparison and their accession numbers are X61367, U12338 and JN872328, respectively.

right). The four Tn1696 derivatives in this study differed from Tn1696 primarily by insertion of distinct integron or integron-associated regions, in other words In834, In27 together with a truncated Tn125 carrying *bla*_{NDM-1} [30], In1222 and In1212, respectively, at the same site within *res* as observed for Tn1696 (Figure 2).

In834 (Figure 2) was a complex class 1 integron [32], which harbored two intrinsic resistance regions (the cassette array *qacH4-bla*_{OXA-10}-*aadA1e* and *catA2*) and two additional integrated resistance regions (the *chrA* region and a truncated Tn1721 carrying the class A tetracycline module *tetA(A)-tetR(A)* [33]). The three class 1 integrons In27, In1212 and In1222 (Figure 2) carried the cassette arrays *dfrA12-gcuF-aadA2*, *aacA* and *qnrVC4-qacF-aacA4-cmlA1g-bla*_{OXA-10}-*aadA1e-dfrA14b*, respectively; moreover, In1222 integrated an additional macrolide resistance unit IS26-*mph(A)-mrx-mphR(A)*-IS6100 [33]. In834 and In1222 were bordered by paired inverted repeats (IRi, integrase end; IRt, *tmi* end), while In27 and In1212 lacked IRt.

Tn6358 and Tn6359 (Figure 2) had paired terminal IRL and IRR sequences, which were further bracketed by 5-bp direct repeats (DRs; target site duplication signals for transposition). By contrast, the Tn1696-In27-Tn125 region and the Tn1696-In1212 region carried only IRLs but lacked IRRs, making them could not be discriminated as naive transposons. All these IRL and IRR elements were interrupted by IS5075 or its close variant IS4321 or IS4321R, all of which targeted the IRL and IRR sequences of Tn21 subgroup transposons [34]. Tn6358 and Tn6359 harbored the identical *mer* locus, which was found in several other type 1 IncC plasmids such as pRMH760 [14] and pSH696.135 [35] but genetically distant from the counterpart in Tn1696.

In the Tn1696-In1212 region of p112298-tetA (Figure 2), downstream of In1212 was a Δ IS1326 element, together with the IS26 to *orf124* region presented in various plasmids such as p0716-KPC (accession number KY270849). The *orf633* to *orf124* region (a portion of the IS26 to *orf124* region) contained several genes (*parA2-parB2*, *umuCD*, *psiAB* and *ardA*) for plasmid maintenance and was found in only p112298-tetA but not pR148, pHS36-NDM and pVAS3-1, and this region did not belong to the IncC backbone.

The ARI-B islands

The ARI-B islands from p112298-tetA and pVAS3-1 were identified as the derivatives of the *floR-tetAR-strAB-sul2* region, a 13.6-kb IS26- and ISCR2-associated MDR module widely found in IncC plasmids [36] and herein designated the *floR-tetAR-strAB-sul2-bla*_{CMY-2} region and the *aphA1*-In641-*floR-tetAR-strAB-sul2* region, respectively (Figure 3A).

The ARI-B island from p112298-tetA (Figure 3A) was organized in order of the *floR-tetAR-strAB-sul2* region, the *repC-repA*-IS26 remnant of the composite transposition Tn6029 [37], the IS903D-*iroN* remnant of the ISEcp1-*bla*_{CTX-M}-IS903D-*iroN* transposition unit [38] and the truncated ISEcp1-*bla*_{CMY-2} unit with insertion of IS1R.

The ARI-B island from pVAS3-1 (Figure 3A) was generated from the connection of the *aphA1*-carrying composite transposition Tn4352 [37] with the *floR-tetAR-strAB-sul2* region, which was further interrupted and truncated by insertion of a 15.7-kb exogenous region that was composed of IS1006, the *tnpAR-res-tnpM* remnant of Tn21 [40] and a *sul3*-associated class 1 integron In641 [41] carrying the cassette array *estX3-psp-aadA2-cmlA-aadA1a-qacH2*.

The assembly of these two ARI-B islands involved complex homologous recombination events based on the common regions (e.g., IS26 and ISEcp1) of the above prototype mobile elements.

The *bla*_{KPC-2} region from p112298-tetA

Insertion of a core *bla*_{KPC} platform (the Tn6376 to Δ *repB* region) into the cryptic unit transposon Tn1722 generated Tn6296 (Figure 3B), as initially found in pKP048 [39]. Tn6296 and its various derivatives represent the major *bla*_{KPC} vehicles in KPC-encoding plasmids from China [42,43].

The *bla*_{KPC-2} region from p112298-tetA was organized sequentially as the 5'-terminal small remnant of the ISEcp1-*bla*_{CMY-2} unit, an 8.4-kb region composed of multiple IS elements, a 6.8-kb truncated version of Tn6296 designated Δ Tn6296 and IS26 (Figure 3B). Δ Tn6296 carried the *bla*_{KPC-2} gene and was flanked by two directly oriented copies of IS26 in p112298-tetA, displaying a composite transposon-like structure. The almost identical element was also found in the MDR region of p112298-KPC [15], which coexisted with p112298-tetA in the 112298 isolate.

The presence of the Δ Tn6296-carrying composite transposon-like elements and their surrounding regions within p112298-tetA and p112298-KPC were verified by PCR amplifications that targeted a set of jointing fragments using genomic DNAs of strains 112298, 112298-tetA-DH10B and 112298-KPC-TOP10 (an *E. coli* TOP10 electroporant containing p112298-KPC [15]) as templates (Supplementary Figure 3).

Figure 3. The ARI-B islands and the *ISEcp1*-*bla_{CMY}* units, and comparison with related regions. Shown is the comparison of related regions with the ARI-B islands (A) and the *ISEcp1*-*bla_{CMY}* units (B). Genes are denoted by arrows. Genes, mobile elements and other features are colored based on their functional classification. Shading denotes regions of homology (nucleotide identity >95%). Numbers in brackets indicate the nucleotide positions within the corresponding plasmids. The arrowheads indicate the location of PCR primers and the expected amplicons. See Supplementary Figure 3 for the PCR results. Tn6029 [37], the *ISEcp1*-*bla_{CMY-2}* transposition unit [38] and Tn6269 [39] are prototype mobile elements as the references for sequence comparison and their accession numbers are GQ150541, KR337637 and FJ628167, respectively.

Table 4. Antimicrobial drug susceptibility profiles.

Antibiotics	MIC (mg/l)/antimicrobial susceptibility		
	112298	112298-tetA-DH10B	DH10B
Ampicillin	>1024/R	1024/R	<4/S
Ceftazidime	>512/R	16/R	<4/S
Meropenem	64/R	4/R	<1/S
Cefoxitin	>1024/R	64/R	<8/S
Aztreonam	>512/R	32/R	<4/S
Tetracycline	>128/R	16/R	<1/S
Chloramphenicol	1024/R	128/R	<8/S
Gentamicin	128/R	64/R	<4/S
Ciprofloxacin	8/R	<1/S	<1/S
Azithromycin	64/R	<4/S	<4/S
Fosfomycin	256/R	<64/S	<64/S
Tigecycline	<1/S	<1/S	<1/S
Colistin	<1/S	<1/S	<1/S

R: Resistant; S: Sensitive.

Plasmid transfer & resistance phenotypes

p112298-tetA could be transferred from the 112298 isolate into *E. coli* DH10B through electroporation, generating an electroporant 112298-tetA-DH10B. Repeated attempts failed to transfer p112298-tetA from the 112298 isolate into *E. coli* EC600 through conjugation transfer and p112298-tetA was nonconjugative due to lesion of conjugation transfer gene function (see above). Both 112298 and 112298-tetA-DH10B had the activity class A carbapenemase (data not shown), which was attributable to the production of KPC enzyme. Strain 112298 was resistant to ampicillin, ceftazidime, meropenem, cefoxitin, aztreonam, tetracycline, chloramphenicol, gentamicin, ciprofloxacin, azithromycin and fosfomycin, but remained susceptible to tigecycline and colistin; 112298-tetA-DH10B was resistant to the first eight drugs but susceptible to all the other ones tested (Table 4).

Discussion

IncC plasmids often contain one or more exogenous resistance islands, which are inserted at various sites in the plasmid backbones. pR148 carries a single island ARI-A and represents a precursor of IncC plasmids [3]. pHS36-NDM harbors not only ARI-A but also the *ISEcp1-bla_{CMY-6}* unit that encodes CMY-6, a Trp221Leu variant of CMY-2. In addition to ARI-A and the *ISEcp1-bla_{CMY-2}* unit, pVAS3-1 has gained an additional island ARI-B. p112298-tetA contains a total of three complete islands, namely ARI-A, ARI-B and the *bla_{KPC-2}* region. The ARI-A or ARI-B islands from different plasmids display distinct profiles of resistance genes and associated mobile elements such as insertion sequences, integrons and transposons. pR148, pHS36-NDM, pVAS3-1 and p112298-tetA might represent the relatively 'oldest' to 'youngest' lineages, under the assumption that IncC plasmids have evolved to consecutively acquire exogenous resistance regions.

IncC plasmids contain two separate conjugal transfer regions *tra1* and *tra2*. As observed in conjugative pR148, *tra1* is composed of 24 genes that are oriented in the same transcriptional direction and code for a F-type type IV secretion system (T4SS) necessary for conjugative transfer [44], while *tra2* consist of 13 genes involved in not only T4SS assembly (*traFHG*) but also transcriptional regulation (*acr1*, *acr2* and *acaCD*) of conjugal transfer genes [45,46]. The *ISEcp1-bla_{CMY}* unit lies within an intergenic region of *tra1* of pHS36-NDM and pVAS3-1, and these two plasmids retain the ability to conjugal transfer [7,28]. By contrast, the *tra1* region of p112298-tetA is interrupted into two separate parts *tra1-3'* and *tra1-5'* and moreover, *tra1-5'* is translocated to a position far away from *tra1-3'*, rendering p112298-tetA nonconjugative. Rearrangements in the *tra1* region and thereby the resulting loss of *tra* genes due to the insertion of the *ISEcp1-bla_{CMY}* units have been shown to correlate with the conjugal transfer deficiency in IncC plasmids [36].

Compared with pR148, pHS36-NDM and pVAS3-1, p112298-tetA is quite unique due to the presence of: the *orf633* to *orf375* backbone region encoding plasmid maintenance functions; the *i2* insert characteristic of type 2 IncC; a third island (the *bla_{KPC-2}* region) containing a Δ Tn6296-carrying composite transposon-like element, which is shared by p112298-KPC coexisting with p112298-tetA; the segmented *ISEcp1-bla_{CMY-2}* unit further

merged with ARI-B and the *bla*_{KPC-2} region; and a lesion in modular organization of *traI* making p112298-tetA nonconjugative. Complex rearrangement and homologous recombination events have occurred during evolution of p112298-tetA.

Conclusion & future perspective

The plasmid p112298-tetA from clinical *C. freundii* 112298 was fully sequenced and compared with the two available sequenced type 1 IncC plasmids pHS36-NDM and pVAS3-1 from China and the type 1 IncC reference plasmid pR148. All of them are MDR plasmids and contain one or more exogenous islands encoding antimicrobial resistance mechanisms, which are inserted at various sites in the plasmid backbones. These plasmids display significant modular differences in both plasmid backbones and accessory regions and, thereby, dramatically different profiles of antimicrobial resistance genes. Epidemiological studies and routine surveillance of IncC plasmids in China are needed. Comparative genomics analysis of a large collection of IncC plasmids will provide a deeper understanding of evolution and diversification of IncC plasmids.

Summary points

- This study presented the complete sequence of p112298-tetA, a type 1 IncC plasmid in *Citrobacter freundii* from a Chinese teaching hospital.
- Comparative genomics analysis was performed with the three available sequenced type 1 IncC plasmids pHS36-NDM, pVAS3-1 and p112298-tetA from China, and the type 1 IncC reference plasmid pR148.
- These plasmids contain one or more exogenous resistance islands, which are inserted at various sites in the plasmid backbones and displayed dramatically different profiles of antimicrobial resistance genes.
- pR148, pHS36-NDM, pVAS3-1 and p112298-tetA represents the relatively 'oldest' to 'youngest' lineages from an evolutionary point of view of acquisition of exogenous resistance regions.
- Complex rearrangement and homologous recombination events have occurred during evolution of p112298-tetA, making it significantly differ modularly from pR148, pHS36-NDM and pVAS3-1 with respect to both plasmid backbones and exogenous resistance regions.

Supplementary data

To view the supplementary data that accompany this paper please visit the journal website at: www.futuremedicine.com/doi/suppl/10.2217/fmb-2017-0072.

Financial & competing interests disclosure

This work was funded by the China Postdoctoral Science Foundation (2015M582803) and the Foundation of the State Key Laboratory of Pathogen and Biosecurity of China (SKLPBS1520). The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Ethical conduct of research

The authors state that they have obtained appropriate institutional review board approval or have followed the principles outlined in the Declaration of Helsinki for all human or animal experimental investigations. In addition, for investigations involving human subjects, informed consent has been obtained from the participants involved.

References

1. Fricke WF, Welch TJ, McDermott PF *et al.* Comparative genomics of the IncA/C multidrug resistance plasmid family. *J. Bacteriol.* 191(15), 4750–4757 (2009).
2. Carattoli A, Miriagou V, Bertini A *et al.* Replicon typing of plasmids encoding resistance to newer beta-lactams. *Emerg. Infect. Dis.* 12(7), 1145–1148 (2006).
3. Harmer CJ, Hall RM. The A to Z of A/C plasmids. *Plasmid* 80, 63–82 (2015).
4. Datta N, Hedges RW. R factors of compatibility group A. *J. Gen. Microbiol.* 74(2), 335–337 (1973).
5. Harmer CJ, Hamidian M, Hall RM. pIP40a, a type 1 IncC plasmid from 1969 carries the integrative element GIsul2 and a novel class II mercury resistance transposon. *Plasmid* 92 17–25 (2017).

6. Johnson TJ, Lang KS. IncA/C plasmids: an emerging threat to human and animal health? *Mob. Genet. Elements* 2(1), 55–58 (2012).
7. Ye L, Li R, Lin D *et al.* Characterization of an IncA/C multidrug resistance plasmid in *Vibrio alginolyticus*. *Antimicrob. Agents Chemother.* 60(5), 3232–3235 (2016).
8. Li R, Lin D, Chen K, Wong MH, Chen S. First detection of AmpC beta-lactamase bla(CMY-2) on a conjugative IncA/C plasmid in a *Vibrio parahaemolyticus* isolate of food origin. *Antimicrob. Agents Chemother.* 59(7), 4106–4111 (2015).
9. Folster JP, Katz L, McCullough A *et al.* Multidrug-resistant IncA/C plasmid in *Vibrio cholerae* from Haiti. *Emerg. Infect. Dis.* 20(11), 1951–1953 (2014).
10. Del Castillo CS, Hikima J, Jang HB *et al.* Comparative sequence analysis of a multidrug-resistant plasmid from *Aeromonas hydrophila*. *Antimicrob. Agents Chemother.* 57(1), 120–129 (2013).
11. McIntosh D, Cunningham M, Ji B *et al.* Transferable, multiple antibiotic and mercury resistance in Atlantic-Canadian isolates of *Aeromonas salmonicida* subsp. *Salmonicida* is associated with carriage of an IncA/C plasmid similar to the *Salmonella enterica* plasmid pSN254. *J. Antimicrob. Chemother.* 61(6), 1221–1228 (2008).
12. Kim MJ, Hirono I, Kurokawa K *et al.* Complete DNA sequence and analysis of the transferable multiple-drug resistance plasmids (R Plasmids) from *Photobacterium damsela* subsp. *Piscicida* isolates collected in Japan and the United States. *Antimicrob. Agents Chemother.* 52(2), 606–611 (2008).
13. Doublet B, Boyd D, Douard G, Praud K, Cloeckaert A, Mulvey MR. Complete nucleotide sequence of the multidrug resistance IncA/C plasmid pR55 from *Klebsiella pneumoniae* isolated in 1969. *J. Antimicrob. Chemother.* 67(10), 2354–2360 (2012).
14. Harmer CJ, Hall RM. pRMH760, a precursor of A/C(2) plasmids carrying blaCMY and blaNDM genes. *Microb. Drug Resist.* 20(5), 416–423 (2014).
15. Feng J, Qiu Y, Yin Z *et al.* Coexistence of a novel KPC-2-encoding MDR plasmid and an NDM-1-encoding pNDM-HN380-like plasmid in a clinical isolate of *Citrobacter freundii*. *J. Antimicrob. Chemother.* 70(11), 2987–2991 (2015).
16. Liang Q, Yin Z, Zhao Y *et al.* Sequencing and comparative genomics analysis of the IncHI2 plasmids pT5282-mphA and p112298-catA and the IncHI5 plasmid pYNKP001-dfrA. *Int. J. Antimicrob. Agents* 49(6), 709–718 (2017).
17. CLSI. Performance standards for antimicrobial susceptibility testing: twenty-fifth informational supplement M100-S25. CLSI, PA, USA (2015).
18. Nederbragt AJ. On the middle ground between open source and commercial software – the case of the Newbler program. *Genome Biol.* 15(4), 113 (2014).
19. Brettin T, Davis JJ, Disz T *et al.* RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci. Rep.* 5, 8365 (2015).
20. Boratyn GM, Camacho C, Cooper PS *et al.* BLAST: a more efficient report with usability improvements. *Nucleic Acids Res.* 41, W29–W33 (2013).
21. Boutet E, Lieberherr D, Tognolli M *et al.* UniProtKB/Swiss-Prot, the manually annotated section of the UniProt KnowledgeBase: how to use the entry view. *Methods Mol. Biol.* 1374, 23–54 (2016).
22. O’leary NA, Wright MW, Brister JR *et al.* Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 44(D1), D733–D745 (2016).
23. Jia B, Raphenya AR, Alcock B *et al.* CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res.* 45, D566–D573 (2016).
24. Zankari E, Hasman H, Cosentino S *et al.* Identification of acquired antimicrobial resistance genes. *J. Antimicrob. Chemother.* 67(11), 2640–2644 (2012).
25. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. ISfinder: the reference centre for bacterial insertion sequences. *Nucleic Acids Res.* 34, D32–D36 (2006).
26. Moura A, Soares M, Pereira C, Leitao N, Henriques I, Correia A. INTEGRALL: a database and search engine for integrons, integrases and gene cassettes. *Bioinformatics* 25(8), 1096–1098 (2009).
27. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32(5), 1792–1797 (2004).
28. Huang J, Wang M, Ding H *et al.* New Delhi metallo-beta-lactamase-1 in carbapenem-resistant *Salmonella* strain, China. *Emerg. Infect. Dis.* 19(12), 2049–2051 (2013).
29. Partridge SR, Brown HJ, Stokes HW, Hall RM. Transposons Tn1696 and Tn21 and their integrons In4 and In2 have independent origins. *Antimicrob. Agents Chemother.* 45(4), 1263–1270 (2001).
30. Poiriel L, Bonnin RA, Boulanger A, Schrenzel J, Kaase M, Nordmann P. Tn125-related acquisition of blaNDM-like genes in *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* 56(2), 1087–1089 (2012).
31. Allmeier H, Cresnar B, Greck M, Schmitt R. Complete nucleotide sequence of Tn1721: gene organization and a novel gene product with features of a chemotaxis protein. *Gene* 111(1), 11–20 (1992).

32. Quiroga MP, Andres P, Petroni A *et al.* Complex class 1 integrons with diverse variable regions, including aac(6')-Ib-cr, and a novel allele, qnrB10, associated with ISCR1 in clinical enterobacterial isolates from Argentina. *Antimicrob. Agents Chemother.* 51(12), 4466–4470 (2007).
33. Partridge SR. Analysis of antibiotic resistance regions in Gram-negative bacteria. *FEMS Microbiol. Rev.* 35(5), 820–855 (2011).
34. Partridge SR, Hall RM. The IS1111 family members IS4321 and IS5075 have subterminal inverted repeats and target the terminal inverted repeats of Tn21 family transposons. *J. Bacteriol.* 185(21), 6371–6384 (2003).
35. Han J, Lynne AM, David DE *et al.* DNA sequence analysis of plasmids from multidrug resistant *Salmonella enterica* serotype Heidelberg isolates. *PLoS ONE* 7(12), e51160 (2012).
36. Fernandez-Alarcon C, Singer RS, Johnson TJ. Comparative genomics of multidrug resistance-encoding IncA/C plasmids from commensal and pathogenic *Escherichia coli* from multiple animal sources. *PLoS ONE* 6(8), e23415 (2011).
37. Cain AK, Liu X, Djordjevic SP, Hall RM. Transposons related to Tn1696 in IncHI2 plasmids in multiply antibiotic resistant *Salmonella enterica* serovar Typhimurium from Australian animals. *Microb. Drug Resist.* 16(3), 197–202 (2010).
38. Zong Z, Yu R, Wang X, Lu X. blaCTX-M-65 is carried by a Tn1722-like element on an IncN conjugative plasmid of ST131 *Escherichia coli*. *J. Med. Microbiol.* 60(Pt 4), 435–441 (2011).
39. Shen P, Wei Z, Jiang Y *et al.* Novel genetic environment of the carbapenem-hydrolyzing beta-lactamase KPC-2 among Enterobacteriaceae in China. *Antimicrob. Agents Chemother.* 53(10), 4333–4338 (2009).
40. Liebert CA, Hall RM, Summers AO. Transposon Tn21, flagship of the floating genome. *Microbiol. Mol. Biol. Rev.* 63(3), 507–522 (1999).
41. Chang CY, Lu PL, Lin CC, Lee TM, Tsai MY, Chang LL. Integron types, gene cassettes, antimicrobial resistance genes and plasmids of *Shigella sonnei* isolates from outbreaks and sporadic cases in Taiwan. *J. Med. Microbiol.* 60(Pt 2), 197–204 (2011).
42. Wang L, Fang H, Feng J *et al.* Complete sequences of KPC-2-encoding plasmid p628-KPC and CTX-M-55-encoding p628-CTXM coexisted in *Klebsiella pneumoniae*. *Front. Microbiol.* 6, 838 (2015).
43. Dai X, Zhou D, Xiong W *et al.* The IncP-6 plasmid p10265-KPC from *Pseudomonas aeruginosa* carries a novel deltaiseC33-associated bla KPC-2 gene cluster. *Front. Microbiol.* 7, 310 (2016).
44. Christie PJ. The mosaic type IV secretion systems. *EcoSal Plus* 7(1), doi:10.1128/ecosalplus.ESP-0020-2015 (2016) (Epub ahead of print).
45. Carraro N, Matteau D, Luo P, Rodrigue S, Burrus V. The master activator of IncA/C conjugative plasmids stimulates genomic islands and multidrug resistance dissemination. *PLoS Genet.* 10(10), e1004714 (2014).
46. Lang KS, Johnson TJ. Characterization of Acr2, an H-NS-like protein encoded on A/C2-type plasmids. *Plasmid* 87–88, 17–27 (2016).

Susceptibility testing and reporting of new antibiotics with a focus on tedizolid: an international working group report

Mark H Wilcox¹, Natalia Dmitrieva², Ana Cristina Gales³, Irina Petukhova², Suleiman Al-Obeid⁴, Flavia Rossi⁵ & Joseph M Blondeau^{*,6}

¹Leeds Teaching Hospitals NHS Trust & University of Leeds, Leeds, UK

²N.N. Blokhin Cancer Research Centre, Moscow, Russia

³Division of Infectious Diseases, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, Brazil

⁴Microbiology Department, Security Forces Hospital, Riyadh, Saudi Arabia

⁵Hospital das Clínicas da Faculdade de Medicina, Seção de Microbiologia, Divisão de Laboratório Central LIM03, Universidade de São Paulo, São Paulo, Brazil

⁶Royal University Hospital, Saskatchewan, Canada

* Author for correspondence: Tel.: +1 306 655 6943; Fax: +1 306 655 6947; joseph.blondeau@saskatoonhealthregion.ca

Inappropriate use and overuse of antibiotics are among the most important factors in resistance development, and effective antibiotic stewardship measures are needed to optimize outcomes. Selection of appropriate antimicrobials relies on accurate and timely antimicrobial susceptibility testing. However, the availability of clinical breakpoints and *in vitro* susceptibility testing often lags behind regulatory approval by several years for new antimicrobials. A Working Group of clinical/medical microbiologists from Brazil, Canada, Mexico, Saudi Arabia, Russia and the UK recently examined issues surrounding antimicrobial susceptibility testing for novel antibiotics. While commercially available tests are being developed, potential surrogate antibiotics may be used as marker of susceptibility. Using tedizolid as an example of a new antibiotic, this special report makes recommendations to optimize routine susceptibility reporting.

First draft submitted: 9 June 2017; Accepted for publication: 26 July 2017; Published online: 18 August 2017

Keywords: antimicrobial susceptibility testing • surrogate antibiotic • tedizolid

Widespread use and misuse of antibiotics have been associated with the selection and spread of antibiotic-resistant strains in both humans and animals. Antimicrobial resistance now poses a significant threat to public health across the world, with alarming increases in the number of infections due to multidrug-resistant pathogens [1]. Consequently, availability of new antimicrobials is now a critical unmet need. In an effort to promote the development of new, more active antibiotics, the WHO recently published its first list of antibiotic-resistant 'priority pathogens' divided into three categories – critical, high and medium. It is hoped that the publication of this list will help guide research toward pathogens that present the greatest risks to public health [2].

While research on antibiotics to combat multidrug-resistant Gram-negative pathogens (e.g., carbapenem-resistant Enterobacteriaceae) is recognized as being a critical need, high priority is also given to Gram-positive bacteria, which remain a major cause of nosocomial infections [2,3]. Methicillin-resistant *Staphylococcus aureus* (MRSA), in particular, is a prevalent threat in many parts of the world, and is often associated with significant morbidity and mortality, particularly in the elderly [4,5]. While vancomycin has been the mainstay of treatment for the management of MRSA infections [6], therapeutic failures have been reported for isolates possessing minimum inhibitory concentrations (MICs) >1.5 µg/ml [7–9]. In addition, a loading dose of vancomycin is recommended and therapeutic drug monitoring required to optimize outcomes, so complicating clinical use [10], particularly in low-resource settings. The high activity of the novel oxazolidinone antibiotic tedizolid against Gram-positive bacteria, including MRSA and vancomycin-resistant enterococci, presents a new treatment option for these challenging pathogens, providing more potent activity than linezolid *in vitro* [11–14]. Although the *optrA* gene in some strains of Enterococci may confer elevated MICs to both tedizolid and linezolid [15–17], this mechanism has not yet been identified in MRSA.

Noninferiority of tedizolid phosphate (given for 6 days) versus linezolid (10 days) has been demonstrated in two Phase III studies in patients with acute bacterial skin and skin structure infections [18,19].

Inappropriate use and overuse of antibiotics are among the most important factors in the development of resistance, and effective antibiotic stewardship measures are needed to optimize the use of antimicrobials [20]. This includes use of restrictive reporting, and/or encouraging diverse prescribing to avoid overuse of valuable antibiotics and help control the increase in antimicrobial resistance. Clinicians' ability to select appropriate antimicrobials relies on accurate and timely antimicrobial susceptibility testing (AST), a process designed to predict clinical efficacy [20]. However, performing effective AST is challenging, particularly for newly approved antibiotics, since availability of clinical breakpoints and inclusion on commercial panels and/or access to other *in vitro* susceptibility materials often lag behind regulatory approval by several years [1]. Obtaining sufficient clinical data to enable clinical breakpoints for new agents to be established is also problematic. A Working Group of clinical/medical microbiologists from Brazil, Canada, Mexico, Saudi Arabia, Russia and the UK was convened to examine issues surrounding AST for novel antibiotics. This article summarizes their recommendations for optimal routine reporting of susceptibility to new antibiotics, using the novel oxazolidinone antibiotic tedizolid as an exemplar.

Overview of testing methodology

In recent years, few new Gram-positive directed antibiotics have become available, with only dalbavancin hydrochloride, oritavancin and tedizolid phosphate being approved over the past 5 years [21–26]. In order to select the optimal therapy for their patients, clinicians must be aware of changes in resistance patterns and the effectiveness of the antibiotics used to treat particular organisms over time. For this reason, a number of global antimicrobial surveillance programs, including STAR (Surveillance of Tedizolid Activity and Resistance; tedizolid), SOAR (Survey of Antibiotic Resistance; amoxicillin/clavulanic acid, cefuroxime, cefaclor, azithromycin and ofloxacin), ZAAPS (Zyvox® Annual Appraisal of Potency and Spectrum; linezolid) and TEST (Tigecycline Evaluation Surveillance Trial; tigecycline), have been initiated [27–31].

Susceptibility testing can be either automated (e.g., Vitek2, Microscan and Phoenix), or manual (e.g., disk diffusion [as recommended by the European Committee on Antimicrobial Susceptibility Testing – EUCAST – or Clinical and Laboratory Standards Institute – CLSI], broth microdilution or gradient testing [e.g., E-test®]). Each system has advantages and limitations, with some providing quality control ranges [1,32–35]. Working Group members report that the majority of laboratories in their countries use automated testing systems routinely for susceptibility testing, most of which provide MICs. Manual testing is frequently used to confirm automated AST results and/or MICs; E-test® and disk diffusion are the preferred methods, with the latter being the more favored choice in cost-conscious environments. Of note, in routine susceptibility testing, vancomycin MICs may not reflect accurately the actual MICs and their confirmation by E-test® has recently been recommended to minimize the risk of treatment failure [36]. Determination of MICs is important for certain strains not identified by an automated system or disk diffusion (e.g., vancomycin-intermediate *S. aureus* [VISA]). Failure with empiric vancomycin treatment in a Saudi 69-year-old male patient, for example, prompted physicians to obtain more accurate susceptibility data on the pathogen isolated from blood. This led to the discovery of the first heterogeneous vancomycin-intermediate *Staphylococcus aureus* (hVISA), a threatening signal to the effectiveness of empiric vancomycin treatment in hospitalized patients with severe infections [37].

Effective performance of AST by clinical microbiology laboratories is essential in order to determine susceptibility to the chosen empirical antimicrobial agents, and to detect the emergence of resistance [38]. Use of susceptibility breakpoints (e.g., those provided by the CLSI and EUCAST) is important, both for consistent reporting of antimicrobial susceptibility and for consistency and comparability of international surveillance schemes [39]. EUCAST and CLSI also advocate use of epidemiological cut-off values as one of several tools in the process of establishing clinical breakpoints; determination of such cut-off values may promote more susceptibility testing and allow MIC values to be put into context [40]. While most laboratories follow either CLSI or EUCAST breakpoints for susceptibility testing, some may use both if a pathogen is not included in the guidelines they normally follow. When there is no available information on appropriate clinical breakpoints for certain bacteria, physicians may rely on their best judgment with regard to MIC findings and experience in eradicating those pathogens, and/or on consultation with clinical/medical microbiologists or infectious disease specialists.

Most laboratories in the Working Group members' countries report an isolate's susceptibility to be 'susceptible', 'resistant' or 'intermediate' according to either CLSI or EUCAST criteria. The 'intermediate' category has been used in a number of various ways, for example as a buffer to prevent very major and major errors, to be used alongside

the 'resistant' category to indicate results that are 'nonsusceptible' and to indicate pathogen or drug combinations where an increased drug exposure may be necessary for optimal results [40]. However, antibiotic decision-making is usually based on 'susceptible' or 'resistant' categories, since treatment choices for 'intermediate' susceptibility are not clear-cut. 'Intermediate' should be interpreted as 'nonsusceptible' and physicians should thus choose an alternative antibiotic to which the pathogen is susceptible. Nevertheless, reporting susceptibility as 'intermediate' can be useful in some cases, for example if two antibiotics are used synergistically (e.g., meropenem and colistin) or to optimize the antimicrobial dosage regimen [41].

Selection of antibiotics for routine susceptibility testing depends primarily on the type of AST method, particularly if an automated panel is used. All antibiotics included on the panel are routinely tested, although not all of those tested may be reported; reports are usually issued only for first-line antibiotics. Until the cause of an infection is known, initial therapy is generally empiric and guided by clinical presentation [20]. Clinical/medical microbiologists and infectious disease specialists have an important role in advising physicians on the most appropriate antibiotics to use, although decision-making can also involve infectious disease specialists and clinical pharmacologists. The number and selection of antimicrobials tested is dependent on the organism isolated, infection site, the institution's formulary, physician requests and the automated panel or other testing methodology used [38]. Notably, the choice of antibiotics to be tested via automated methods can be limited, including for new antibiotics that have yet to be adopted by automated systems. While inclusion of an antibiotic on the hospital formulary is a key factor in selection for susceptibility testing, availability on the automated panel, supply of the manual testing equipment (disks or gradient strips) and requests from physicians are also important.

The inclusion of new antibiotics on commercial, automated test panels is often delayed for a considerable time after approval [38,42]. However, agents from the same class with similar activities can be used as surrogate markers (i.e., class representatives) to predict susceptibility of clinical isolates to new agents and/or to those not included in routine testing. Clinical laboratories have used surrogate testing successfully for decades (see Table 1). Reliability of a surrogate marker is typically analyzed by testing the categorical agreement between the susceptibility results for the two agents, defining errors as very major (i.e., false-susceptibility), major (i.e., false-resistance) or minor (i.e., the result for one agent was intermediate while the other agent was susceptible, nonsusceptible or resistant) [43]. However, it should be noted that the chance of a very major error is extremely low when the occurrence of resistant isolates is rare or absent. Furthermore, in rare cases, susceptibility to a surrogate antibiotic might not reveal resistance to the agent in question, for example a recently recognized, uncommon variant of *fexA* conferring resistance to florfenicol, but not to chloramphenicol [44]. Such reports highlight the need to review the utility of surrogate susceptibility testing as new data emerge.

Testing of new antibiotics: a focus on tedizolid

An alternative approach to susceptibility testing is needed for new antibiotics as their inclusion on commercial AST panels is often delayed for several years after approval, which complicates testing and reporting [1]. Despite inclusion on hospital formulary, the use of new antibiotics may be restricted for some months, which limits the opportunity for physicians to gain clinical experience. This may be difficult to explain to physicians if the antibiotic is already included in clinical practice guidelines. Consequently, clinical/medical microbiologists have a responsibility to provide clinically relevant information to physicians and explain why it is or is not appropriate to use an antibiotic. Additional barriers to susceptibility testing of new antibiotics include: problems with the availability and regular supply of materials required for manual testing; not being included on the hospital formulary; lack of breakpoints for certain pathogens; and the increased workload and costs associated with supplementary testing. Working Group members report that novel antibiotics will most likely be routinely tested once they become available on an automated panel(s). Until then, susceptibility testing for new agents is performed manually, but often undertaken only upon physician request. In the short term, surrogate agents (that are included on automated panels) can be used to predict pathogen susceptibility to new antibiotics [1].

Several methods are currently available to test tedizolid susceptibility, including the use of linezolid as a surrogate antibiotic. Linezolid susceptibility is considered a highly reliable surrogate for tedizolid susceptibility; a high categorical agreement has been reported between the susceptibility of tedizolid and linezolid; and the very major error rates were low ($\leq 0.2\%$) for all organisms tested [42]. Based on these findings, EUCAST recommends that isolates susceptible to linezolid can be reported as susceptible to tedizolid [47]. For isolates intermediate/resistant to linezolid, an MIC test must be performed to confirm susceptibility or resistance to tedizolid. Most laboratories in Working Group members' countries would add to the AST report a 'tedizolid susceptible' result based on a

Table 1. Antibiotics used as surrogate agents in clinical practice.

Surrogate agent	Pathogen	Antibiotic(s) for which susceptibility is reported	Ref.
Ampicillin	Enterococci	Amoxicillin-clavulanic acid, ampicillin-sulbactam, piperacillin, piperacillin-tazobactam	[45]
Cefepime	<i>H. influenzae</i>	Ceftaroline	[46]
Cefoxitin	Staphylococci	Oxacillin	[45]
		Cephalosporins [†]	[47]
Ceftazidime	<i>H. influenzae</i>	Ceftaroline	[46]
Ceftriaxone	<i>S. pneumoniae</i>	Ceftaroline	[46]
	<i>H. influenzae</i>	Ceftaroline	[46]
Cephalothin	Enterobacteriaceae	Cefapirin, cephradine, cefaclor, cefadroxil, cefpodoxime, cephalexin and loracarbef	[45,48]
Ertapenem	Enterobacteriaceae <i>Haemophilus</i> spp. <i>S. pneumoniae</i>	Doripenem	[49]
Erythromycin	Streptococci	Azithromycin, clarithromycin and dirithromycin	[45]
Imipenem	<i>S. aureus</i>	Ceftaroline	[46]
	<i>P. aeruginosa</i>	Doripenem	[50]
Levofloxacin	<i>S. pneumoniae</i>	Fluoroquinolones	[51]
Linezolid	Staphylococci Enterococci Streptococci <i>Streptococcus anginosus</i> group	Tedizolid	[42]
	Staphylococci Streptococci		[47]
Meropenem	<i>S. aureus</i>	Ceftaroline	[46]
	<i>P. aeruginosa</i>	Doripenem	[50]
Nalidixic acid	<i>H. influenzae</i> <i>M. catarrhalis</i>	Ciprofloxacin, levofloxacin, moxifloxacin and ofloxacin	[47]
Norfloxacin	Staphylococci Streptococci	Ciprofloxacin, levofloxacin, moxifloxacin and ofloxacin	[47]
Oxacillin	<i>S. aureus</i>	Cefazolin, ceftriaxone, ceftaroline and nafcillin	[52]
Penicillin	Enterococci	Ampicillin, amoxicillin, amoxicillin-clavulanic acid, ampicillin-sulbactam, piperacillin, piperacillin-tazobactam	[45]
Teicoplanin	Staphylococci Streptococci	Dalbavancin	[53]
Tetracycline	<i>Vibrio cholerae</i>	Doxycycline [‡]	[45]
	Staphylococci Streptococci <i>H. influenzae</i> <i>M. catarrhalis</i> <i>N. gonorrhoeae</i>	Doxycycline and minocycline	[47]
Vancomycin	Staphylococci Streptococci	Dalbavancin	[53,54]
	Staphylococci Streptococci	Dalbavancin and oritavancin	[47]
	Staphylococci Streptococci Enterococci	Oritavancin	[55]

^{††}Not cefixime, ceftazidime, ceftibuten and ceftolozane-tazobactam.

Disk diffusion should not be used for doxycycline due to poor correlation with minimum inhibitory concentration results.

surrogate 'linezolid susceptible' actual testing result, but would not report a 'tedizolid resistance' result without knowledge of the results of actual tedizolid susceptibility testing. AST results are typically discussed with the treating physician before reporting so as not to delay clinical decision-making. The CLSI recommends that tedizolid and linezolid are included in the Group B optional primary test and reported selectively [45]. Additional methods available for tedizolid susceptibility testing include MIC test strips (Liofilchem[®], Liofilchem s.r.l., Roseto, Italy) and a broth microdilution device (Sensititre[™] Thermo Scientific[™], Thermo Fisher Scientific Inc., Cleveland, OH, USA) developed primarily for research purposes [56]. Use of 20 µg disk diffusion has been approved by the CLSI for measurement of tedizolid susceptibility as a quality control measure [45]. To obtain susceptibility results in microbiology laboratories, currently a more practicable 2 µg disk diffusion method is under development, in addition to automated susceptibility testing panels. It should be noted that methodological testing issues have been observed with tedizolid and linezolid which suggest that 80% inhibition MIC end point criterion should be employed for testing both agents [57].

Tedizolid has demonstrated more potent activity *in vitro* than linezolid against Gram-positive bacteria, including MRSA and vancomycin-resistant enterococci, as well as noninferiority in patients with skin and skin structure infections [11–14,18–19]. Of note, tedizolid retains *in vitro* activity against *S. aureus* and other Gram-positive bacterial strains that harbor the *cfi*-gene encoded methylase enzyme [58,59]. This methylase enzyme confers resistance against five structurally different antibiotic classes (e.g., clindamycin in the lincosamide class, chloramphenicol in the phenicol class, and linezolid, but not tedizolid, in the oxazolidinone class) [58]. The encoded enzyme methylates the A2053 nucleotide in the peptidyl-transferase center of the 23S ribosomal RNA, which is a very highly conserved site, and this methylation prevents binding of antibiotics to peptidyl-transferase center [60]. Importantly, chromosomal mutations in Domain V of rRNA or ribosomal L3 protein identified to date were demonstrated to confer resistance to both linezolid and tedizolid, stressing the need to test tedizolid susceptibility [59,61]. The Working Group stresses the importance of providing tedizolid susceptibility results in order to guide clinicians in selecting the most effective agent for their patients, and to provide an alternative option in cases of resistance development. They recommend the adoption of EUCAST guidelines on use of linezolid as a surrogate to predict tedizolid susceptibility [47], in order to enable routine inclusion of tedizolid in AST reports by microbiology laboratories. The Working Group also advocates having tedizolid MIC test strips in place in laboratories so that it is possible to perform tedizolid susceptibility testing, particularly when nonsusceptibility to linezolid is detected. Furthermore, they highlight the importance of a surveillance program for monitoring resistance and appropriate antibiotic use of all new antibiotics.

Conclusion & recommendations

The Working Group has highlighted the difficulty in acquiring sufficient clinical data to enable clinical breakpoints for new antibiotics to be established and the time taken (possibly up to 5 years) for such agents to be included in automated testing panels, which are often used for routine susceptibility testing. Policy makers should aim to minimize delays in the adoption of new breakpoints for antibiotics against emerging pathogens, particularly when containment of spread is vital; delays should be reduced to less than 1.5 years whenever possible [62]. However, to guide clinicians on appropriate treatment, it is critical that the susceptibility of new antibiotics is reported. Using tedizolid as an example, if it is not possible to determine susceptibility, data may be extrapolated by using linezolid as a surrogate, allowing subsequent appropriate use of tedizolid. In situations identified by a clinical/medical microbiologist (e.g., linezolid nonsusceptible strains of Gram-positive bacteria) an appropriate manual test for tedizolid will be necessary (Figure 1). Susceptibility testing and routine reporting of selected new antibiotics can be desirable to ensure that clinicians make the appropriate choices for the management of infection, and that development of resistance is closely monitored as new agents become available.

Future perspective

At present, there is a considerable time lag between approval of new antibiotics, availability of clinical breakpoints and inclusion on commercial AST panels. It is hoped that closer coordination in the coming years between those involved in drug development and AST panels and regulatory authorities will shorten this delay, enabling routine testing of new agents at the time of approval. In the meantime, to ensure appropriate antibiotic choice, it is essential that laboratories use the available tools to enable them to report susceptibility of new antibiotics.

Figure 1. Recommended approach for routine reporting of susceptibility of new antibiotics, for example, tedizolid.

†Use of broth microdilution is also applicable.

As recommended by published evidence/according to susceptibility testing guidance [42,47].

NS: Nonsusceptible; S: Susceptible.

Acknowledgements

The authors acknowledge J Alder, Bayer, Whippany, NJ, USA; M Glenscek-Sieberth, Bayer, Wuppertal, Germany; and L Li, Bayer, Berlin, Germany for their constructive contributions to the discussions of the Working Group at the virtual meeting.

Financial & competing interest disclosure

The authors received no payment for writing, preparation and approval of this manuscript. Each author received a consultancy honorarium for participation in a virtual advisory board meeting where the issues prompting these recommendations of the Working Group were discussed. M Wilcox has received consultancy fees from Aicuris, Allergan, Astra Zeneca, Basilea, Bayer, Durata, European Tissue Symposium, Merck, Meridian, Nabriva, Roche, Synthetic Biologics, Valneva, The Medicine Company; grants from Micropharm; and grants and consultancy fees from Abbott, Actelion, Alere, Astellas, Biomerieux, Da Volterra, Qiagen, MotifBio, Pfizer, Sanofi Pasteur, Seres, and Summit. N Dmitrieva has no other conflict of interest. AC Gales has received consultancy fees and/or research grants from Astra Zeneca, Bayer, BD, Merck Sharp & Dohme, and Pfizer. I Petukhova has no other conflict of interest. S Al-Obeid has no other conflict of interest. F Rossi has no other conflict of interest. J Blondeau has received consultancy fees and/or research grants from Abbott, Alcon, Allergan, Astellas, Bayer, Cooper Vision, Merck, Pfizer, Roche, Sanofi, Summit, and Wyeth. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

Editorial support was provided by Highfield Communication, Oxford, UK, sponsored by Bayer AG, Berlin, Germany.

Executive summary

Background

- A Working Group of clinical/medical microbiologists from Brazil, Canada, Mexico, Saudi Arabia, Russia and the UK has made recommendations for optimal routine testing and reporting of susceptibility to new antibiotics, using the novel oxazolidinone antibiotic tedizolid as an example.

Overview of testing methodology

- Most laboratories use automated testing systems (e.g., Vitek2, Microscan and Phoenix) routinely for susceptibility testing, with manual testing (e.g., EUCAST or CLSI disk diffusion, broth microdilution or gradient testing [e.g., E-test[®]]) used to confirm the results and minimum inhibitory concentrations (MICs).
- The inclusion of new antibiotics on commercial, automated test panels and/or the availability of susceptibility testing materials, are often delayed for a considerable time after approval; however, agents from the same class with similar activities can be used as surrogate markers to predict susceptibility of clinical isolates to new agents and/or to those not included in routine testing.

Testing of new antibiotics: a focus on tedizolid

- Several methods are currently available to test tedizolid susceptibility: use of linezolid as a surrogate antibiotic, MIC test strips (Liofilchem[®], Liofilchem s.r.l., Roseto, Italy) and a broth microdilution device (Sensititre[™] Thermo Scientific[™], Thermo Fisher Scientific Inc., Cleveland, OH, USA).
- Adoption of EUCAST guidelines on use of linezolid as a surrogate to predict tedizolid susceptibility is recommended in order to enable routine inclusion of tedizolid in antimicrobial susceptibility testing reports; having tedizolid MIC test strips in place in laboratories is also advocated so that tedizolid susceptibility testing can be performed when nonsusceptibility to linezolid is detected.

Conclusion & recommendations

- It can take up to 5 years for new antibiotics to be included in automated testing panels yet it is critical that the susceptibility of these agents is reported.
- If it is not possible to determine susceptibility, data may be extrapolated by using a surrogate, for example, linezolid susceptibility as a reliable surrogate for tedizolid.
- Susceptibility testing and routine reporting of selected new antibiotics is desirable to ensure that clinicians make the appropriate choices for the management of infection.

Open access

This work is licensed under the Attribution-NonCommercial-NoDerivatives 4.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>

References

Papers of special note have been highlighted as: ●● of considerable interest

1. Humphries R, Hindler J. Emerging resistance, new antimicrobial agents. . . but no tests! The challenge of antimicrobial susceptibility testing in the current US regulatory landscape. *Clin. Infect. Dis.* 63, 83–88 (2016).
2. World Health Organization. Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics. February 27, 2017. www.who.int/medicines/publications/WHO-PPL-Short-Summary_25Feb-ET_NM_WHO.pdf?ua=1
- 3. This first list of antibiotic-resistant ‘priority pathogens’ published by WHO will help guide research toward pathogens that present the greatest risks to public health.
3. Liu C, Bayer A, Cosgrove SE *et al.* Clinical practice guidelines by the Infectious Diseases Society of America for the treatment of methicillin-resistant *Staphylococcus aureus* infections in adults and children. *Clin. Infect. Dis.* 52, e18–e55 (2011).
4. Rollnik JD. Outcome of MRSA carriers in neurological early rehabilitation. *BMC Neurol.* 14, 34 (2014).
5. Schmid H, Romanos A, Schiff H, Lederer S. Persistent nasal methicillin-resistant *Staphylococcus aureus* carriage in hemodialysis outpatients: a predictor of worse outcome. *BMC Nephrol.* 14, 93 (2013).
6. Appelbaum PC. Reduced glycopeptide susceptibility in methicillin-resistant *Staphylococcus aureus* (MRSA). *Int. J. Antimicrob. Agents* 30(5), 398–408 (2007).
7. Rybak MJ, Lomaestro BM, Rotschafer JC *et al.* Vancomycin therapeutic guidelines: a summary of consensus recommendations from the Infectious Diseases Society of America, the American Society of Health-System Pharmacists, and the Society of Infectious Diseases Pharmacists. *Clin. Infect. Dis.* 49, 325–327 (2009).
8. Choi EY, Huh JW, Lim CM *et al.* Relationship between the MIC of vancomycin and clinical outcome in patients with MRSA nosocomial pneumonia. *Intensive Care Med.* 37, 639–647 (2011).
9. Kullar R, Davis SL, Levine DP, Rybak MJ. Impact of vancomycin exposure on outcomes in patients with methicillin-resistant *Staphylococcus aureus* bacteremia: support for consensus guidelines suggested targets. *Clin. Infect. Dis.* 52, 975–981 (2011).

10. Ye ZK, Tang HL, Zhai SD. Benefits of therapeutic drug monitoring of vancomycin: a systematic review and meta-analysis. *PLoS ONE* 8(10), e77169 (2013).
11. Chen KH, Huang YT, Liao CH, Sheng WH, Hsueh PR. *In vitro* activities of tedizolid and linezolid against gram-positive cocci associated with acute bacterial skin and skin structure infections and pneumonia. *Antimicrob. Agents Chemother.* 59(10), 6262–6265 (2015).
12. Flanagan S, Bartizal K, Minassian SL, Fang E, Prokocimer P. *In vitro*, *in vivo*, and clinical studies of tedizolid to assess the potential for peripheral or central monoamine oxidase interactions. *Antimicrob. Agents Chemother.* 57(7), 3060–3066 (2013).
13. Prokocimer P, Bien P, DeAnda C, Pillar CM, Bartizal K. *In vitro* activity and microbiological efficacy of tedizolid (TR-700) against Gram-positive clinical isolates from a Phase II study of oral tedizolid phosphate (TR-701) in patients with complicated skin and skin structure infections. *Antimicrob. Agents Chemother.* 56(9), 4608–4613 (2012).
14. Shaw KJ, Poppe S, Schaadt R *et al.* *In vitro* activity of TR-700, the antibacterial moiety of the prodrug TR-701, against linezolid-resistant strains. *Antimicrob. Agents Chemother.* 52(12), 4442–4447 (2008).
15. Wang Y, Lv Y, Cai J *et al.* A novel gene, *optrA*, that confers transferable resistance to oxazolidinones and phenicols and its presence in *Enterococcus faecalis* and *Enterococcus faecium* of human and animal origin. *J. Antimicrob. Chemother.* 70(8), 2182–2190 (2015).
16. Gawryszevska I, Zabicka D, Hryniewicz W, Sadowy E. Linezolid-resistant enterococci in Polish hospitals: species, clonality and determinants of linezolid resistance. *Eur. J. Clin. Microbiol. Infect. Dis.* 36(7), 1279–1286 (2017).
17. Pfaller MA, Mendes RE, Streit JM, Hogan PA, Flamm RK. Five-year summary of *in vitro* activity and resistance mechanisms of linezolid against clinically important Gram-positive cocci in the United States from the LEADER Surveillance Program (2011 to 2015). *Antimicrob. Agents Chemother.* 61(7), pii:e00609–17 (2017).
18. Prokocimer P, De Anda C, Fang E, Mehra P, Das A. Tedizolid phosphate vs linezolid for treatment of acute bacterial skin and skin structure infections: the ESTABLISH-1 randomized trial. *JAMA* 309(6), 559–569 (2013).
19. Moran GJ, Fang E, Corey GR, Das AF, De Anda C, Prokocimer P. Tedizolid for 6 days versus linezolid for 10 days for acute bacterial skin and skin-structure infections (ESTABLISH-2): a randomised, double-blind, phase 3, non-inferiority trial. *Lancet Infect. Dis.* 14(8), 696–705 (2014).
20. Leekha S, Terrell CL, Edson RS. General principles of antimicrobial therapy. *Mayo Clin. Proc.* 86, 156–167 (2011).
21. Food and Drug Administration. Dalvance: FDA News Release. (2014). www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm398724.htm
22. Food and Drug Administration. Orbactiv: FDA News Release. (2014). www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm408475.htm
23. Food and Drug Administration. Sivextro: FDA News Release. (2014). www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm402174.htm
24. European Medicines Agency. Xydalba. (2015). www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/002840/smops/Positive/human_smop.000765.jsp&mid=WC0b01ac058001d127
25. European Medicines Agency. Orbactiv. (2015). www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/003785/smops/Positive/human_smop.000781.jsp&mid=WC0b01ac058001d127
26. European Medicines Agency. Sivextro. (2015). www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/002846/smops/Positive/human_smop.000780.jsp&mid=WC0b01ac058001d127
27. Sahm DF, Deane J, Bien PA *et al.* Results of the surveillance of tedizolid activity and resistance program: *in vitro* susceptibility of gram-positive pathogens collected in 2011 and 2012 from the United States and Europe. *Diagn. Microbiol. Infect. Dis.* 81(2), 112–118 (2015).
28. Van PH, Binh PT, Minh NH, Morrissey I, Torumkuney D. Results from the Survey of Antibiotic Resistance (SOAR) 2009–11 in Vietnam. *J. Antimicrob. Chemother.* 71(Suppl. 1), i93–i102 (2016).
29. Mendes RE, Hogan PA, Streit JM, Jones RN, Flamm RK. Zyvox® Annual Appraisal of Potency and Spectrum (ZAAPS) program: report of linezolid activity over 9 years (2004–12). *J. Antimicrob. Chemother.* 69, 1582–1528 (2014).
30. Sader HS, Castanheira M, Farrell DJ, Flamm RK, Mendes RE, Jones RN. Tigecycline antimicrobial activity tested against clinical bacteria from Latin American medical centres: results from SENTRY Antimicrobial Surveillance Program (2011–2014). *Int. J. Antimicrob. Agents* 48, 144–150 (2016).
31. Giammanco A, Calà C, Fasciana T, Dowzicky MJ. Global assessment of the activity of tigecycline against multidrug-resistant Gram-negative pathogens between 2004 and 2014 as part of the Tigecycline Evaluation and Surveillance Trial. *mSphere* 2(1), pii:e00310–16 (2017).
32. Matuschek E, Brown DF, Kahlmeter G. Development of the EUCAST disk diffusion antimicrobial susceptibility testing method and its implementation in routine microbiology laboratories. *Clin. Microbiol. Infect.* 20, O255–O266 (2014).
33. Cantón R, Livermore DM, Morosini MI, Díaz-Regañón J, Rossolini GM, PREMIUM Study Group. Etest® versus broth microdilution for ceftaroline MIC determination with *Staphylococcus aureus*: results from PREMIUM, a European multicentre study. *J. Antimicrob. Chemother.* 72(2), 431–436 (2017).

34. Steward CD, Stocker SA, Swenson JM *et al.* Comparison of agar dilution, disk diffusion, MicroScan, and Vitek antimicrobial susceptibility testing methods to broth microdilution for detection of fluoroquinolone-resistant isolates of the family Enterobacteriaceae. *J. Clin. Microbiol.* 37, 544–547 (1999).
35. Karatuna O. *Quality Assurance In Antimicrobial Susceptibility Testing, Latest Research Into Quality Control.* Akyar I (Ed.). InTech (2012).
36. Van Hal SJ, Lodise TP, Paterson DL. The clinical significance of vancomycin minimum inhibitory concentration in *Staphylococcus aureus* infections: a systematic review and meta-analysis. *Clin. Infect. Dis.* 54, 755–771 (2012).
37. Al-Obeid S, Haddad Q, Cherkaoui A, Schrenzel J, Francois P. First detection of an invasive *Staphylococcus aureus* strain (D958) with reduced susceptibility to glycopeptides in Saudi Arabia. *J. Clin. Microbiol.* 48(6), 2199–2204 (2010).
38. Jorgensen J, Ferraro M. Antimicrobial susceptibility testing: a review of general principles and contemporary practices. *Clin. Infect. Dis.* 49, 1749–1755 (2009).
39. Brown D, Canton R, Dubreuil L *et al.* Widespread implementation of EUCAST breakpoints for antibacterial susceptibility testing in Europe. *Euro Surveill.* 20, pii:21008 (2015).
40. Silley P. Susceptibility testing methods, resistance and breakpoints: what do these terms really mean? *Rev. Sci. Tech.* 31(1), 33–41 (2012).
41. Daoud Z, Mansour N, Masri K. Synergistic combination of carbapenems and colistin against *P. aeruginosa* and *A. baumannii*. *Open J. Med. Microbiol.* 3, 253–258 (2013).
42. Zurenko G, Bien P, Bensaci M, Patel HN, Thorne G. Use of linezolid susceptibility test results as a surrogate for the susceptibility of Gram-positive pathogens to tedizolid, a novel oxazolidinone. *Ann. Clin. Microbiol. Antimicrob.* 13, 46 (2014).
- **The high categorical agreement between MIC values for tedizolid and linezolid, and low very major error rates reported in this article for all organism groups tested support the use of linezolid as a reliable surrogate for tedizolid susceptibility testing.**
43. Clinical and Laboratory Standards Institute (CLSI). Development of *In vitro* Susceptibility Testing Criteria and Quality Control Parameters; Approved Guideline. (document M23–A3, 3rd Edition). CLSI, Wayne, PA (2008).
44. Gómez-Sanz E, Kadlec K, Feßler AT, Zarazaga M, Torres C, Schwarz S. A novel *fexA* variant from a canine *Staphylococcus pseudintermedius* isolate that does not confer florfenicol resistance. *Antimicrob. Agents Chemother.* 57(11), 5763–5766 (2013).
45. Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Susceptibility Testing (document M100-S26); CLSI (2016).
- **Susceptibility breakpoints for antimicrobial susceptibility testing.**
46. Jones RN, Flamm RK, Sader HS, Stilwell MG. Interim susceptibility testing for ceftaroline, a new MRSA-active cephalosporin: selecting potent surrogate β -lactam markers to predict ceftaroline activity against clinically indicated species. *Diagn. Microbiol. Infect. Dis.* 75(1), 89–93 (2013).
47. European Committee on Antimicrobial Susceptibility Testing. Breakpoint tables for interpretation of MICs and zone diameters. Version 6.0, (2016). www.eucast.org.
- **Susceptibility breakpoints for antimicrobial susceptibility testing.**
48. Turnidge JD, Subcommittee on Antimicrobial Susceptibility Testing of the Clinical and Laboratory Standards Institute. Cefazolin and enterobacteriaceae: rationale for revised susceptibility testing breakpoints. *Clin. Infect. Dis.* 52(7), 917–924 (2011).
49. Jones RN, Sader HS, Fritsche TR, Janeček MJ. Selection of a surrogate beta-lactam testing agent for initial susceptibility testing of doripenem, a new carbapenem. *Diagn. Microbiol. Infect. Dis.* 59(4), 467–472 (2007).
50. Hagihara M, Kuti JL, Nicolau DP. Predicting doripenem susceptibility based on meropenem and imipenem interpretation for *Pseudomonas aeruginosa*. *Diagn. Microbiol. Infect. Dis.* 72(3), 258–262 (2012).
51. Sader HS, Koeth LM, Poupard JA, Jones RN. Interpretive categorical accuracy of fluoroquinolone reference broth microdilution MIC results when testing *Streptococcus pneumoniae*: selection of a surrogate testing agent. *Diagn. Microbiol. Infect. Dis.* 62, 460–463 (2008).
52. Kang N, Housman ST, Nicolau DP. Assessing the surrogate susceptibility of oxacillin and cefoxitin for commonly utilized parenteral agents against methicillin-susceptible *Staphylococcus aureus*: focus on ceftriaxone discordance between predictive susceptibility and *in vivo* exposures. *Pathogens* 4(3), 599–605 (2015).
53. Jones RN, Sader HS, Fritsche TR, Hogan PA, Sheehan DJ. Selection of a surrogate agent (vancomycin or teicoplanin) for initial susceptibility testing of dalbavancin: results from an international antimicrobial surveillance program. *J. Clin. Microbiol.* 44, 2622–2625 (2006).
54. Dunne MW, Sahn D, Puttagunta S. Use of vancomycin as a surrogate for dalbavancin *in vitro* susceptibility testing: results from the DISCOVER studies. *Ann. Clin. Microbiol. Antimicrob.* 14, 19 (2015).
55. Jones RN, Moeck G, Arhin FF, Dudley MN, Rhomberg PR, Mendes RE. Results from oritavancin resistance surveillance programs (2011 to 2014): clarification for using vancomycin as a surrogate to infer oritavancin susceptibility. *Antimicrob. Agents Chemother.* 60(5), 3174–3177 (2016).
56. Jones RN, Holliday NM, Rhomberg PR. Validation of a commercial dry-form broth microdilution device (Sensititre) for testing tedizolid, a new oxazolidinone. *J. Clin. Microbiol.* 53(2), 657–659 (2015).

57. Pfaller MA, Flamm RK, Jones RN, Farrell DJ, Mendes RE. Activities of tedizolid and linezolid determined by the reference broth microdilution method against 3,032 Gram-positive bacterial isolates collected in Asia-Pacific, Eastern Europe, and Latin American countries in 2014. *Antimicrob. Agents Chemother.* 60(9), 5393–5399 (2016).
58. Long KS, Poehlsgaard J, Kehrenberg C, Schwarz S, Vester B. The Cfr rRNA methyltransferase confers resistance to Phenicol, Lincosamides, Oxazolidinones, Pleuromutilins, and Streptogramin A antibiotics. *Antimicrob. Agents Chemother.* 50(7), 2500–2505 (2006).
59. Locke JB, Finn J, Hilgers M *et al.* Structure-activity relationships of diverse oxazolidinones for linezolid-resistant *Staphylococcus aureus* strains possessing the *efr* methyltransferase gene or ribosomal mutations. *Antimicrob. Agents Chemother.* 54(12), 5337–5343 (2010).
60. Stefani S, Bongiorno D, Mongelli G, Campanile F. Linezolid resistance in Staphylococci. *Pharmaceuticals* 3, 1988–2006 (2010).
61. Zhanel GG, Love R, Adam H *et al.* Tedizolid: a novel oxazolidinone with potent activity against multidrug-resistant Gram-positive pathogens. *Drugs* 75(3), 253–270 (2015).
62. Bartsch SM, Huang SS, Wong KF *et al.* Impact of delays between Clinical and Laboratory Standards Institute and Food and Drug Administration revisions of interpretive criteria for carbapenem-resistant enterobacteriaceae. *J. Clin. Microbiol.* 54(11), 2757–2762 (2016).

The hidden pathogenic potential of environmental fungi

Glauber R de S Araújo¹, Wanderley de Souza¹ & Susana Frases^{*,1}

¹Laboratório de Ultraestrutura Celular Hertha Meyer, Instituto de Biofísica Carlos Chagas Filho, Federal University of Rio de Janeiro, Rio de Janeiro, RJ, Brazil

* Author for correspondence: Tel.: +55 21 3938 6593; susanafrases@biof.ufrj.br

Invasive fungal infections are a growing threat to immunocompromised patients, highlighting the importance of monitoring fungal pathogens. Global warming (including climatic oscillations) may select for environmental species that have acquired thermotolerance, a key step toward pathogenesis to humans. Also, important virulence factors have developed in environmental fungi, because they are essential for yeast survival in the environment. Thus, fungi traditionally regarded as nonpathogenic to humans have virulence factors similar to those of their pathogenic relatives. Here, we highlight the emergence of saprophytic environmental fungi – including species of *Cryptococcus*, *Aspergillus*, *Penicillium*, *Candida* and *Scedosporium* – as new human pathogens. Emerging pathogens are, in some cases, resistant to the available antifungals, potentiating the threat of novel fungal diseases.

First draft submitted: 27 June 2017; Accepted for publication: 29 August 2017; Published online: 23 November 2017

Keywords: antifungal resistance • *Aspergillus* • *Candida* • *Cryptococcus* • environmental fungi • fungal virulence • global warming • *Penicillium* • *Scedosporium* • thermotolerance

Fungal infections on the rise

Over the last decades, the world has experienced an increase in the incidence and spread of emerging fungal infections. These events may be associated with several factors, including the increase in life expectancy, which generates older individuals with a weaker immune response, as well as increased epigenetic abnormalities that may favor disease establishment. In the case of fungal infections, the reduction in the host's natural defenses (caused by disease or immunosuppressive medication) meant that numerous pathogens previously considered harmless gradually became disease agents, causing serious infections that are both resistant to antifungal and fatal. The incidence of pathogenic fungal infections has risen enough to constitute a fundamental change the epidemiology of invasive fungal diseases, especially in immunocompromised individuals, such as those with HIV, cancer or subjected to transplants [1–5].

The distribution of fungal species on the planet was determined, broadly, by the following effects: the separation of fungal species through continental drift; the European colonial expansion, when various fungal diseases were brought from the old world to the new world; and human displacement (emigration, migration, immigration) [6]. Environmental fungi that are pathogenic to humans exist in a broad range of geographic areas, but they are more common in warmer regions of the planet, probably due to temperature and humidity restrictions on their growth or propagation. This is the case for endemic species such as *Blastomyces* spp., *Coccidioides* spp., *Histoplasma* spp. and the species *Cryptococcus gattii* [7]. In contrast, environmental fungi capable of growing at wide temperature ranges are ubiquitous and cosmopolitan [8].

Natural disasters are clear examples of environmental conditions changing the dynamics of fungal populations. The link between natural disasters – which alter the local environmental conditions – and the subsequent occurrence of a typical fungal infections in disaster-affected individuals has been increasingly recognized [8], and highlights the importance of environmental factors on the dispersion of fungal diseases. For example, after the Japan earthquake and tsunami in 2011, fatal cases of pulmonary multiorgan dissemination by *Aspergillus fumigatus* and *Scedosporium apiospermum* were described, while these fungi do not typically cause invasive infections in immunocompetent hosts [9,10].

Global warming & the emergence of novel fungal pathogens

The emergence of new diseases most frequently results from changes in host or pathogen ecology, from environmental alterations, or from interactions of all of these components. Increased environmental pressures – such as global warming and higher ultraviolet radiation levels – may result in stronger pathogens with advanced virulence mechanisms that favor survival in harsher environmental conditions [11].

Climate change is a consequence of biotic and abiotic processes, such as variations in solar radiation and human activities, which have been identified as the primary causes of global warming. Evidence suggests that the increase in the Earth's temperature over the last centuries has already resulted in marked ecological changes, including alterations in the seasons and in the distribution of a variety of species [12,13]. Climate change is expected to widen the invasion niche of many fungal species [14], and it has already driven shifts in the composition of saprotrophic fungal populations [15]. Most fungi grow at 12–30°C, suffering a rapid decline in growth and viability at temperatures above 30°C [16]. Consequently, the high mammalian body temperatures are sufficient to inhibit the growth and replication of the majority of fungal strains [17–19]. However, a warmer climate could change the distribution of both heat-tolerant and susceptible species, by selecting for those that are more thermotolerant. This effect is expected not only to widen the geographic area of pathogenic (i.e., intrinsically heat-tolerant) species, but also to facilitate the close interaction of environmental fungi with human populations, leading to the emergence of novel fungal pathogens.

Virulence factors in environmental fungi: a reservoir of potential pathogenesis tools

The majority of fungi are facultative parasites, and the passage through a host is not an essential part of their life cycle. Most fungi capable of systemic human infection are environmental species whose opportunistic and fatal pathogenicity derives mostly from strategies developed to counteract environmental stresses [20]. Therefore, to survive in the dangerous environment of the human body – under high temperatures and constant attack by powerful immune defenses – fungi make use of virulence factors initially developed for survival in the rich ecological niche of the soil. In recent years, studies with amoeba, slime molds and worms have led to the proposal that interactions between fungi and other environmental microbes, including predators, select for characteristics that also favor survival in animal hosts [21–25]. Virulence factors allow fungi to interact with the complex and dangerous soil biota; thus, soil residence is associated with the maintenance of pathogenic fungal potential for mammalian host infection [26,27]. Interestingly, Rhame and co-workers (1995) showed that bacterial virulence factors are also versatile, with the same factors conferring pathogenicity to both plant and animal hosts [28].

Morphological changes (called fungal 'morphogenesis'), which include variations in cell size and shape, are employed by many fungal species to survive in the environment, and also within 'accidental' hosts. Morphogenesis is considered as a mechanism for fungal locomotion, is important for fungal dissemination in the environment, and may also represent a protection mechanism against amoeboid cell predation [29]. In addition, shape changes can be used as virulence factors in the establishment of fungal infections. Importantly, for most fungi, the ability for shape change is critical for human infectivity, because many pathogenic species enter the human host in the form of small, round airborne dispersal propagules, sporangiospores or conidia, which are produced from hyphal cells. Small fungal cells (1–5 µm in diameter) are ideal for dispersal and entry into hosts, and are also easily internalized by lung macrophages, which is important for pathogenesis by *Cryptococcus* spp. and *Histoplasma* spp. However, after morphogenesis, *Cryptococcus* spp. cells can reach up to 100 µm in diameter ('giant' cells), which happens in later stages of lung infection, while *Histoplasma* sp. form macroconidia of 8–16 µm in diameter [30–32]. Fungal hyphae can penetrate tissue barriers, while yeasts can disseminate more easily to distant sites [24,33]. Thus, 'dimorphic' fungal species such as *Histoplasma capsulatum*, *Paracoccidioides brasiliensis* or *Coccidioides immitis* grow in the hyphal form in the environment, for fungal dissemination and the absorption of nutrients, but shift to the yeast form in the human host. Temperature is a key host signal to trigger hyphae-to-yeast conversion during the establishment of fungal pathologies [34].

For their nutrition in the environment, fungi excrete an array of enzymes that digest complex molecules, for later absorption of their organic components (polypeptides, sugar or complex carbohydrates). In the host, the secretion of digestive enzymes degrades host tissues, generating nutritious products. Enzyme secretion also protects fungi against antimicrobial mechanisms [35,36]. Different examples of this phenomenon include the secretion of: superoxide dismutase, which protects against toxic free radicals, by detoxifying the superoxide anion [37]; phospholipases, which facilitates intracellular proliferation [38]; and urease, which favors intracellular growth and dissemination [39].

The ability to adapt rapidly to variations in the environment, via processes such as microevolution (i.e., rapid phenotypic changes driven by allelic fluctuations) [40] also provides a selective advantage to fungi during the colonization of human hosts. Rapid adaptation enables fungal pathogens to avoid recognition or destruction by the host immune system, resulting in latency and long-term pathogen persistence. In addition, rapid adaptation facilitates the emergence of resistance to antimicrobial agents, which can develop in sensitive species over time, due to improper antifungal use. Although, most acquired antifungal resistance develops in *Candida* species, resistance has also been described in other types of fungi, such as *Aspergillus* spp. [41–45].

Components of the cell wall that are exposed on the cell surface protect fungi from assaults by the environmental and the host, and some cell wall constituents are considered virulence factors [46]. The cell wall polysaccharides chitin and glucans were associated with an increase in virulence in *Blastomyces dermatitidis*, *P. brasiliensis* and *H. capsulatum*. These polysaccharides protect fungal cells against humoral immune response induction and macrophage activation [47]. On the other hand, surface molecules that play a role in adhesion – such as Gp43 from *P. brasiliensis* [48] – are important for the colonization and subsequent invasion of host tissues.

Some fungal species are also capable of modifying the pre-existing external structure of the cell wall, or producing new structures outside the cell wall, such as the polysaccharide capsule that is the hallmark of the global fungal pathogen *Cryptococcus* spp. [49]. In the environment, the capsule protects the organism against stress conditions such as dehydration and phagocytosis by environmental amoeba [50]. In the host, it contributes to infection by protecting the cell against a variety of host immune defenses, interfering with phagocytosis and suppressing both cellular and humoral immunity [51,52].

Melanins are high-molecular-weight pigments produced by a wide variety of fungal species, including *Cryptococcus* spp., *Paracoccidioides* spp., *Histoplasma* spp., *Sporothrix* spp., *Aspergillus* spp. and *Alternaria* spp. [53]. Given its chemical properties (light absorption, hydrophobicity, insolubility and negative charge), melanin protects fungal cells in the environment from the action of hydrolytic enzymes, UV and gamma radiation, extreme temperatures and toxic compounds such as heavy metals. In the host, melanin affects phagocytosis by macrophages, reduces the release of proinflammatory cytokines by host cells, and protects fungi against antifungal drugs [53–58].

The growing challenge of emerging fungal pathogens

In the last 100 years, advances in regenerative medicine, stem cell research and chemotherapy contributed to a worldwide increase in life expectancy by more than 30 years. While these medical advances reverted effectively the lethality of several diseases, they also increased the immunosuppressed population, which is particularly susceptible to uncommon fungal infections. The development of virulence factors in the environment and of thermotolerance to mammalian temperatures – which was potentially accelerated by global warming – may also have contributed significantly to the rise in opportunistic fungal infections by environmental species. Here, we will highlight some of the key species of environmental fungi that have been increasingly associated with human infections, particularly in immunocompromised individuals.

Although, some species may represent truly novel pathogens, improved diagnosis has also identified as pathogenic some species that were previously thought to be restricted to the environment. New molecular methods for the detection and identification of fungal pathogens have been developed that significantly improved our ability to distinguish between previously identified and new pathogenic species, and led to the description of new opportunistic fungal pathogens in an effective and timely manner [59].

From 2008, *Candida auris* has emerged as a new multidrug-resistant *Candida* species associated with invasive infection and high rates of mortality. The first description of human pathogenesis by *C. auris* was in 2008, of a Japanese patient with otitis [60]. Since then, *C. auris* has been reported as an agent of candidemia in Japan, South Korea, India, Kuwait, South Africa, Pakistan and the UK and, more recently, in Venezuela, Colombia and the USA [61–64]. Multidrug resistance has been widely described in *C. auris*, with the majority of clinical strains reported as resistant to the widely used antifungal fluconazole, while some strains are resistant to azoles in general, to polyenes, and also to echinocandins [41,65–67]. These strains present all the major *Candida* virulence factors, including germination, adherence, biofilm formation, and phospholipase and proteinase production [68]. *C. auris* is a major public health concern due to its rapid spread as a nosocomial infection agent in healthcare settings, with potential to cause outbreaks with high mortality rates [69].

Cryptococcosis has also grown in the last decades as a disseminated fungal infection in immunocompromised patients. Although, *Cryptococcus neoformans* and *Cryptococcus gattii* are the main agents of human cryptococcosis, opportunistic infections by environmental species, such as *C. liquefaciens*, *C. albidus* and *C. laurentii*, have been

observed recently [70]. These infections are probably acquired from environmental reservoirs such as bird excrement, trees, food (cheese and fruit), soil and water. Factors that are likely to contribute to the increased incidence of non-*neoformans* and non-*gattii* human infections include improved laboratory detection, the higher incidence of immunocompromised individuals, and the selective pressure of global warming toward the development of tolerance to human host temperatures [70,71]. Studies on cryptococcal virulence factors are typically conducted in *C. neoformans*, since other species of the genus were presumed to have low pathogenic potential. Recently, Araújo and co-workers (2017) described striking similarities in composition and structure between PS molecules of *C. liquefaciens* and *C. neoformans*, and showed that PS molecules from these species have comparable behavior in several key biological activities [72,73]. The remarkable similarities in capsule ultrastructure and virulence between single isolates of *C. neoformans* and *C. liquefaciens* described in the study of Araújo and co-workers (2017), together with the first descriptions of human fungemia by *C. liquefaciens* in immunocompromised patients [70–74], suggest that this environmental cryptococcal species may represent a novel human pathogen. Importantly, *C. liquefaciens* infections are resistant to 5-fluorocytosine, indicating that treatment of fungemia by this species may be challenging [70,71].

In general, aspergillosis is not a notifiable infection, and it often goes unreported by patients; therefore, it is difficult to determine the exact number of cases of this disease worldwide. However, it is clear that allergic bronchopulmonary aspergillosis is common worldwide, while invasive aspergillosis is considered as an uncommon form associated primarily with hospitalized immunocompromised patients [75,76]. Building works generate dust contamination and disperse large amounts of fungal spores in the environment. Invasive aspergillosis cases are, indeed, often associated with hospital renovation or building work, which can increase the amount of airborne *Aspergillus*, resulting in respiratory infections or surgical site infections in high-risk patients. Although, *Aspergillus* species are the most common causative pathogens in fungaemia outbreaks related to construction work, *Zygomycetes* and other fungi were occasionally reported [77,78]. Outbreaks of primary cutaneous and central nervous system aspergillosis associated with the use of contaminated medical devices have also been described [78].

Although, aspergillosis remains clearly the most frequent opportunistic mold infection in patients with immunodeficiency, new opportunistic pathogens have now emerged as the cause of life-threatening fungal infections worldwide. One important opportunistic fungal pathogen is *Fusarium* spp., a saprophytic soil fungus capable of causing disseminated infections in patients with hematologic malignancies [79–81]. Invasive fusarial infections are either airborne or inoculated through rupture of the skin barrier and, once established, are refractory to standard antifungal therapy [79–81].

Scedosporium spp. are soil fungi that have been implicated in numerous infections in immunocompromised and immunocompetent patients [82,83]. These mycoses are life threatening in susceptible patients and can be considered truly emerging diseases. *Scedosporium* infections are difficult to diagnose and treat. In particular, *S. apiospermum* and *S. prolificans* – emerging opportunistic pathogens associated with mycetoma and keratitis infections – are inherently resistant to most antifungal agents and, thus, considerably difficult to combat in the clinic [82–84].

An additional emerging pathogen that deserves mention is *Penicillium marneffei*, which is responsible for ‘penicilliosis marneffei’. This disseminated and progressive disease represents the third most common opportunistic infection in HIV patients in certain parts of Southeast Asia (and is also endemic in Southeast Asia and Southern China) [85]. Infections caused by *Penicillium* species other than *P. marneffei* are rare, and the spectrum of disease produced by *Penicillium* spp. is similar to that of *Aspergillus* spp.; thus, diseases by these two fungal species are difficult to distinguish clinically. Pulmonary infection is the most common presentation of penicilliosis marneffei; however, fungal transmission via trauma, surgery or prosthetic material is commonly implicated in nonpulmonary disease [86].

Future perspective

The frequency of invasive mycoses due to opportunistic fungal pathogens has increased significantly in the last two decades. Immunologically competent mammals display robust defenses against fungal diseases, but recent developments in the evolution of human societies has produced large cohorts of individuals susceptible to fungal infections. On the other hand, it is likely that environmental changes (including global warming) are selecting for fungi increasingly resistant to the harsher conditions encountered in mammalian hosts. Consequently, the number of fungal pathogens currently documented is extensive and ever increasing, and one can no longer ignore or discard environmental fungal species as nonpathogenic to humans. Also, improved diagnostic tools with increased microbial genotyping precision are likely to identify disease isolates that represent species not previously known as

pathogenic, but which had been classified incorrectly as a known pathogenic species. Growing resistance to fungal infection, due to excessive antifungal use in some countries, may also contribute to the emergence of novel fungal pathogens.

Executive summary

- Advances in regenerative medicine, transplants and chemotherapy contributed to extend life expectancy, while also increasing the number of immunocompromised hosts susceptible to unusual fungal infections.
- Environmental fungi subjected to mounting environmental pressure have developed new or more potent virulence factors.
- The emergence of saprophytic fungi as mycoses agents is a growing challenge, particularly when associated with drug resistance, highlighting the urgent need for novel antifungal agents.
- Advances in diagnostics have allowed the identification of new pathogenic fungal species previously indistinguishable from known pathogens.

Financial & competing interests disclosure

This work was supported by the Brazilian agencies Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and Fundação de Amparo à Pesquisa do Rio de Janeiro (FAPERJ). The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

The authors would like to thank WORDinSCIENCE Ltd for their expert scientific manuscript editing services.

References

Papers of special note have been highlighted as: ● of interest; ●● of considerable interest

1. Martin GS, Mannino DM, Eaton S *et al.* The epidemiology of sepsis in the United States from 1979 through 2000. *New Engl. J. Med.* 348(16), 1546–1554 (2003).
2. Holzheimer RG, Dralle H. Management of mycoses in surgical patients – review of the literature. *Eur. J. Med. Res.* 7(5), 200–226 (2002).
3. Peres-Bota D, Rodriguez-Villalobos H, Dimopoulos G *et al.* Potential risk factors for infection with *Candida* spp. in critically ill patients. *Clin. Microbiol. Infect.* 10(6), 550–555 (2004).
4. Perlroth J, Choi B, Spellberg B. Nosocomial fungal infections: epidemiology, diagnosis, and treatment. *Med. Mycol.* 45(4), 321–346 (2007).
5. Meersseman W, van Wijngaerden E. Invasive aspergillosis in the ICU: an emerging disease. *Intensive Care Med.* 33(10), 1679–1681 (2007).
6. Casadevall A, Freij JB, Hann-Soden C, Taylor J. Continental drift and speciation of the *Cryptococcus neoformans* and *Cryptococcus gattii* species complexes. *mSphere* 2(2), e00103–17 (2017).
- **Hypothesis is to stimulate thought and discussion about the mechanisms driving fungal speciation and the origins of fungal virulence.**
7. Sifuentes-Osornio J, Corzo-Leon DE, Ponce-de-Leon LA. Epidemiology of invasive fungal infections in Latin America. *Curr. Fungal Infect. Rep.* 6(1), 23–34 (2012).
8. Benedict K, Park BJ. Invasive fungal infections after natural disasters. *Emerg. Infect. Dis.* 20(3), 349–355 (2014).
9. Nakamura Y, Utsumi Y, Suzuki N *et al.* Multiple *Scedosporium apiospermum* abscesses in a woman survivor of a tsunami in northeastern Japan: a case report. *J. Med. Case Rep.* 5, 526 (2011).
10. Igusa R, Narumi S, Murakami K *et al.* *Escherichia coli* pneumonia in combination with fungal sinusitis and meningitis in a tsunami survivor after the Great East Japan Earthquake. *Tohoku J. Exp. Med.* 227(3), 179–184 (2012).
11. Schrag SJ, Wiener P. Emerging infectious disease: what are the relative roles of ecology and evolution? *Trends Ecol. Evol.* 10(8), 319–324 (1995).
12. Wookey PA, Aerts R, Bardgett RD *et al.* Ecosystem feedbacks and cascade processes: understanding their role in the responses of arctic and alpine ecosystems to environmental change. *Global Change Biology* 15(5), 1153–1172 (2009).
13. van der Putten WH. Climate change, aboveground–belowground interactions and species range shifts. *Ann. Rev. Ecol. Evolut. Syst.* 43, 365–383 (2012).
14. Greer A, Ng V, Fisman D. Climate change and infectious diseases in North America: the road ahead. *CMAJ* 178(6), 715–722 (2008).
15. Classen AT, Sundqvist MK, Henning JA *et al.* Direct and indirect effects of climate change on soil microbial and soil microbial–plant interactions: what lies ahead? *Ecosphere* 6(8), 130 (2015).

- **They explore how climatic change affects soil microbes and soil microbe–plant interactions.**
- 16. Robert VA, Casadevall A. Vertebrate endothermy restricts most fungi as potential pathogen. *J. Infect. Dis.* 200(10), 1623–1626 (2009).
- 17. Baxter M, Illston GM. Temperature relationships of fungi isolated at low temperatures from soils and other substrates. *Mycopathologia* 72, 21–25 (1980).
- 18. Marquez LM, Redman RS, Rodriguez RJ, Roossinck MJ. A virus in a fungus in a plant: three-way symbiosis required for thermal tolerance. *Science* 315, 513–515 (2007).
- 19. Garcia-Solache MA, Casadevall A. Global warming will bring new fungal diseases for mammals. *MBio* 1(1), e00061-10 (2010).
- **Hypothesis on how global warming will bring new fungal diseases for mammals.**
- 20. Brown AJ, Budge S, Kaloriti D *et al.* Stress adaptation in a pathogenic fungus. *J. Exp. Biol.* 217(Pt 1), 144–155 (2014).
- 21. Levitz SM. Does amoeboid reasoning explain the evolution and maintenance of virulence factors in *Cryptococcus neoformans*? *Proc. Natl Acad. Sci. USA* 98(26), 14760–14762 (2001).
- 22. Steenbergen JN, Casadevall A. The origin and maintenance of virulence for the human pathogenic fungus *Cryptococcus neoformans*. *Microbes Infect.* 5(7), 667–675 (2003).
- 23. Steenbergen JN, Nosanchuk JD, Malliaris SD, Casadevall A. *Cryptococcus neoformans* virulence is enhanced after growth in the genetically malleable host *Dictyostelium discoideum*. *Infect. Immun* 71(9), 4862–4872 (2003).
- 24. Steenbergen JN, Nosanchuk JD, Malliaris SD, Casadevall A. Interaction of *Blastomyces dermatitidis*, *Sporothrix schenckii*, and *Histoplasma capsulatum* with *Acanthamoeba castellanii*. *Infect. Immun* 72(6), 3478–3488 (2004).
- 25. Steenbergen JN, Shuman HA, Casadevall A. *Cryptococcus neoformans* interactions with amoebae suggest an explanation for its virulence and intracellular pathogenic strategy in macrophages. *Proc. Natl Acad. Sci. USA* 98(26), 15245–15250 (2001).
- 26. McClelland EE, Bernhardt P, Casadevall A. Estimating the relative contributions of virulence factors for pathogenic microbes. *Infect. Immun.* 74(3), 1500–1504 (2006).
- 27. Casadevall A, Pirofski L-A. Accidental virulence, cryptic pathogenesis, martians, lost hosts, and the pathogenicity of environmental microbes. *Eukaryot. Cell* 6(12), 2169–2174 (2007).
- 28. Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel FM. Common virulence factors for bacterial pathogenicity in plants and animals. *Science* 268(5219), 1899–1902 (1995).
- 29. Wang L, Lin X. Morphogenesis in fungal pathogenicity: shape, size, and surface. *PLoS Pathog.* 8(12), e1003027 (2012).
- 30. Seider K, Heyken A, Luttich A, Miramon P, Hube B. Interaction of pathogenic yeasts with phagocytes: survival, persistence and escape. *Curr. Opin. Microbiol.* 13, 392–400 (2010).
- 31. Okagaki LH, Strain AK, Nielsen JN *et al.* Cryptococcal cell morphology affects host cell interactions and pathogenicity. *PLoS Pathog.* 6, e1000953 (2010).
- 32. Zaragoza O, García-Rodas R, Nosanchuk JD *et al.* Fungal cell gigantism during mammalian infection. *PLoS Pathog.* 6, e1000945 (2010).
- 33. Neilson JB, Ivey MH, Bulmer GS. *Cryptococcus neoformans*: pseudohyphal forms surviving culture with *Acanthamoeba polyphaga*. *Infect. Immun.* 20(1), 262–266 (1978).
- 34. Sil A, Andrianopoulos A. Thermally dimorphic human fungal pathogens – polyphyletic pathogens with a convergent pathogenicity trait. *Cold Spring Harb. Perspect. Med.* 5(8), a019794 (2015).
- 35. Boelaert JR, de Locht M, Van Cutsem J *et al.* Mucormycosis during deferoxamine therapy is a siderophore-mediated infection. *In vitro* and *in vivo* animal studies. *J. Clin. Invest.* 91(5), 1979–1986 (1993).
- 36. Kronstad J, Saikia S, Nielson ED *et al.* Adaptation of *Cryptococcus neoformans* to mammalian hosts: integrated regulation of metabolism and virulence. *Eukaryot. Cell* 11(2), 109–118 (2012).
- 37. Cox GM, Harrison TS, McDade HC *et al.* Superoxide dismutase influences the virulence of *Cryptococcus neoformans* by affecting growth within macrophages. *Infect. Immun.* 71(1), 173–180 (2003).
- 38. Cox GM, McDade HC, Chen SC *et al.* Extracellular phospholipase activity is a virulence factor for *Cryptococcus neoformans*. *Mol. Microbiol.* 39(1), 166–175 (2001).
- 39. Rutherford JC. The emerging role of urease as a general microbial virulence factor. *PLoS Pathog.* 10(5), e1004062 (2014).
- 40. Boyce KJ, Wang Y, Verma S, Shakya VPS, Xue C, Idnurm A. Mismatch repair of DNA replication errors contributes to microevolution in the pathogenic fungus *Cryptococcus neoformans*. *MBio* 8(3), e00595-17 (2017).
- 41. Kumar D, Banerjee T, Pratap CB, Tilak R. Itraconazole-resistant *Candida auris* with phospholipase, proteinase and hemolysin activity from a case of vulvovaginitis. *J. Infect. Dev. Ctries* 9(4), 435–437 (2015).
- 42. Lortholary O, Desnos-Ollivier M, Sitbon K *et al.* Recent exposure to caspofungin or fluconazole influences the epidemiology of candidemia: a prospective multicenter study involving 2441 patients. *Antimicrob. Agents Chemother.* 71(9), 4862–4872 (2011).
- 43. Shah DN, Yau R, Lasco TM *et al.* Impact of prior inappropriate fluconazole dosing on isolation of fluconazole-nonsusceptible *Candida* species in hospitalized patients with candidemia. *Antimicrob. Agents Chemother.* 56, 3239–3243 (2012).

44. Ben-Ami R, Olshtain-Pops K, Krieger M *et al.* Antibiotic exposure as a risk factor for fluconazole-resistant *Candida* bloodstream infection. *Antimicrob. Agents Chemother.* 56(5), 2518–2523 (2012).
45. Arendrup MC. Update on antifungal resistance in *Aspergillus* and *Candida*. *Clin. Microbiol. Infect.* 20(Suppl. 6), 42–48 (2013).
46. Hogan LH, Klein BS. Altered expression of surface alpha-1,3-glucan in genetically related strains of *Blastomyces dermatitidis* that differ in virulence. *Infect. Immun.* 62(8), 3543–3546 (1994).
47. Levitz SM. Innate recognition of fungal cell walls. *PLoS Pathog.* 6(4), e1000758 (2010).
48. Flavia Popi AF, Lopes JD, Mariano M. GP43 from *Paracoccidioides brasiliensis* inhibits macrophage functions. An evasion mechanism of the fungus. *Cell Immunol.* 218(1–2), 87–94 (2002).
49. Casadevall A, Perfect JR. *Cryptococcus neoformans*. ASM Press, Washington, DC, USA (1998).
50. Aksenov SI, Babyeva IP, Golubev VI. On the mechanism of adaptation of micro-organisms to conditions of extreme low humidity. *Life Sci. Space Res.* 11, 55–61 (1973).
51. Chang YC, Kwon-Chung KJ. Complementation of a capsule-deficient mutation of *Cryptococcus neoformans* restores its virulence. *Mol. Cell Biol.* 14(7), 4912–4919 (1994).
52. Zaragoza O, Rodrigues ML, De Jesus M, Frases S, Dadachova E, Casadevall A. The capsule of the fungal pathogen *Cryptococcus neoformans*. *Adv. Appl. Microbiol.* 68, 133–216 (2009).
53. Taborda CP, da Silva MB, Nosanchuk JD, Travassos LR. Melanin as a virulence factor of *Paracoccidioides brasiliensis* and other dimorphic pathogenic fungi: a minireview. *Mycopathologia* 165(4–5), 331–339 (2008).
54. Huffnagle GB, Chen GH, Curtis JL, McDonald RA, Strieter RM, Toews GB. Down-regulation of the afferent phase of T cell-mediated pulmonary inflammation and immunity by a high melanin-producing strain of *Cryptococcus neoformans*. *J. Immunol.* 155(7), 3507–3516 (1995).
55. Wang Y, Aisen P, Casadevall A. *Cryptococcus neoformans* melanin and virulence: mechanism of action. *Infect. Immun.* 63(8), 3131–3136 (1995).
56. Doering TL, Nosanchuk JD, Roberts WK, Casadevall A. Melanin as a potential cryptococcal defence against microbicidal proteins. *Med. Mycol.* 37(3), 175–181 (1999).
57. Liu L, Tewari RP, Williamson PR. Laccase protects *Cryptococcus neoformans* from antifungal activity of alveolar macrophages. *Infect. Immun.* 67(11), 6034–6039 (1999).
58. Nosanchuk JD, Gómez BL, Youngchim S *et al.* Histoplasma capsulatum synthesizes melanin-like pigments *in vitro* and during mammalian infection. *Infect. Immun.* 70(9), 5124–5131 (2002).
59. Arvanitis M, Anagnostou T, Fuchs BB, Caliendo AM, Mylonakis E. Molecular and nonmolecular diagnostic methods for invasive fungal infections. *Clin. Microbiol. Rev.* 27(3), 490–526 (2014).
60. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiol. Immunol.* 53, 41–44 (2009).
61. Calvo B, Melo ASA, Perozo-Mena A *et al.* First report of *Candida auris* in America: clinical and microbiological aspects of 18 episodes of candidemia. *J. Infect.* 73, 369–374 (2016).
62. Morales-Lopez SE, Parra-Giraldo CM, Ceballos-Garzon A *et al.* Invasive infections with multidrug-resistant yeast *Candida auris*, Colombia. *Emerg. Infect. Dis.* 23(1), 162–164 (2017).
63. Schelenz S, Hagen F, Rhodes JL *et al.* First hospital outbreak of the globally emerging *Candida auris* in a European hospital. *Antimicrob. Resist. Infect. Control* 5, 35 (2016).
64. Vallabhaneni S, Kallen A, Tsay S *et al.* Investigation of the first seven reported cases of *Candida auris*, a globally emerging invasive, multidrug-resistant fungus – United States, May 2013–August 2016. *Am. J. Transplant* 17(1), 296–299 (2017).
65. Chowdhary A, Anil Kumar V, Sharma C *et al.* Multidrug-resistant endemic clonal strain of *Candida auris* in India. *Eur. J. Clin. Microbiol. Infect. Dis.* 33(6), 919–926 (2014).
66. Khillan V, Rathore N, Kathuria S, Chowdhary A. A rare case of breakthrough fungal pericarditis due to fluconazole-resistant *Candida auris* in a patient with chronic liver disease. *JMM Case Rep.* doi:10.1099/jmmcr.0.T00018 (2014) (Epub ahead of print).
67. Kim M-N, Shin JH, Sung H *et al.* *Candida haemulonii* and closely related species at 5 university hospitals in Korea: identification, antifungal susceptibility, and clinical features. *Clin. Infect. Dis.* 48, e57–e61 (2009).
68. Larkina E, Hagera C, Chandra J *et al.* The emerging pathogen *Candida auris*: growth phenotype, virulence factors, activity of antifungals, and effect of SCY-078, a novel glucan synthesis inhibitor, on growth morphology and biofilm formation. *Antimicrob. Agents Chemother.* 61(5), e02396-16 (2017).
69. Navalkhele BD, Revankar S, Chandrasekar P. *Candida auris*: a worrisome, globally emerging pathogen. *Expert Rev. Anti. Infect. Ther.* 15(9), 1–9 (2017).
70. Khawcharoenporn T, Apisarnthanarak A, Mundy LM. Non-*neoformans* cryptococcal infections: a systematic review. *Infection* 35(2), 51–58 (2007).

71. Conde-Pereira C, Rodas-Rodriguez L, Diaz-Paz M *et al.* Fatal case of polymicrobial meningitis caused by *Cryptococcus liquefaciens* and *Mycobacterium tuberculosis* complex in a human immunodeficiency virus-infected patient. *J. Clin. Microbiol.* 53(8), 2753–2755 (2015).
72. Araujo GDS, Fonseca FL, Pontes B *et al.* Capsules from pathogenic and non-pathogenic *Cryptococcus* spp. manifest significant differences in structure and ability to protect against phagocytic cells. *PLoS ONE* 7(1), e29561 (2012).
73. Araujo GRS, Freitas GJC, Fonseca FL *et al.* The environmental yeast *Cryptococcus liquefaciens* produces capsular and secreted polysaccharides with similar pathogenic properties to those of *C. neoformans*. *Sci. Rep.* 7, 46768 (2017).
74. Takemura H, Ohno H, Miura I *et al.* The first reported case of central venous catheter-related fungemia caused by *Cryptococcus liquefaciens*. *J. Infect. Chemother.* 21(5), 392–394 (2015).
75. Shah A, Panjabi C. Allergic aspergillosis of the respiratory tract. *Eur. Respir. Rev.* 23(131), 8–29 (2014).
76. Kousha M, Tadi R, Soubani A. Pulmonary aspergillosis: a clinical review. *Eur. Respir. Rev.* 20(121), 156–174 (2011).
77. Kanamori H, Rutala WA, Sickbert-Bennett EE, Weber DJ. Review of fungal outbreaks and infection prevention in healthcare settings during construction and renovation. *Clin. Infect. Dis.* 61(3), 433–444 (2015).
78. Krasinski K, Holzman RS, Hanna B, Greco MA, Graff M, Bhogal M. Nosocomial fungal infection during hospital renovation. *Infect. Control* 6(7), 278–282 (1985).
79. Ozkocaman V, Ozkalemkas F, Cilo BD *et al.* Fusariosis in patients with hematologic malignancies at a University Hospital. *Blood* 126(23), 4906 (2015).
80. Campo M, Lewis RE, Kontoyiannis DP. Invasive fusariosis in patients with hematologic malignancies at a cancer center: 1998–2009. *J. Infect.* 60(5), 331–337 (2010).
81. Nucci M, Anaissie E. Fusarium infections in immunocompromised patients. *Clin. Microbiol. Rev.* 20(4), 695–704 (2007).
82. Cortez KJ, Roilides E, Quiroz Telles F *et al.* Infections caused by *Scedosporium* spp. *Clin. Microbiol. Rev.* 21(1), 157–197 (2008).
83. Rahman FU, Irfan M, Fasih N, Jabeen K, Sharif H. Pulmonary scedosporiosis mimicking aspergilloma in an immunocompetent host: a case report and review of the literature. *Infection* 44(1), 127–132 (2016).
84. Rodriguez-Tudela JL, Berenguer J, Guarro J *et al.* Epidemiology and outcome of *Scedosporium prolificans* infection, a review of 162 cases. *Med. Mycol.* 47(4), 359–370 (2009).
85. Nittayananta W. Penicilliosis marneffeii: another AIDS defining illness in Southeast Asia. *Oral Dis.* 5(4), 286–293 (1999).
86. Lyratzopoulos G, Ellis M, Nerringer R, Denning DW. Invasive infection due to *Penicillium* species other than *P. marneffeii*. *J. Infect.* 45(3), 184–195 (2002).

Our complete Author Guidelines are available at www.futuremedicine.com

Audience

The audience for Future Medicine titles consists of research scientists, decision-makers and a range of professionals in the scientific community.

Submission

We accept unsolicited manuscripts. If you are interested in submitting an article, please do so via our online submission system, which can be found at: <http://onlinesubmissions.future-science-group.com/cms/index.php?tag=2%0A>. For presubmission enquiries, please contact the Editorial Director (l.dormer@futuremedicine.com), who will require a brief article outline and working title in the first instance. We also have an active commissioning program whereby the Editor, under the advice of the Editorial Advisory Panel, solicits articles directly for publication.

Peer review & revision

Once the manuscript has been received in-house, it will be peer-reviewed (usually 2–3 weeks). Following peer review, 2 weeks is allowed for any revisions (suggested by the referees/Editor) to be made.

In-house production

Following acceptance of the revised manuscript, it will undergo production in-house. Authors will receive proofs of the article to approve before going to print, and will be asked to sign a copyright transfer form (except in cases where this is not possible, i.e., government employees in some countries).

Article types

For a more detailed description of each article type, please view our author guidelines at: www.futuremedicine.com.

Reviews

Reviews aim to highlight recent significant advances in research, ongoing challenges and unmet needs.

Word limit: 3000 words (excluding Abstract, Executive Summary, References and Figure/Table legends). Mini-reviews are also accepted.

Required sections: (for a more detailed description of these sections go to www.futuremedicine.com):

- Practice points
- Abstract
- Conclusion
- Future perspective
- References
- Reference annotations
- Financial disclosure

Patent reviews

Patent reviews aim to highlight and critically discuss the most important, promising and recent patents granted/ filed in the chosen area. Discussions should be placed within the context of the relevant wider IP landscape, exploring the impact, significance and essential content of the inventions under discussion.

Word limit: 3000 words (excluding Abstract, Practice points, References and Figure/Table legends). Mini-reviews are also accepted.

Research articles

Types accepted:

Full research article: Primary research articles must present novel science that represents a substantial advancement in the field under investigation.

Preliminary communication

Preliminary communication articles are intended for short reports of studies that present promising improvements or developments on existing areas of research.

Methodology: Methodology articles should provide an overview of a new experimental or computational method, test or procedure. The method described may be either completely novel, or may offer a demonstrable improvement on an existing approach.

Word limit: 3000 words

Required sections: (for a more detailed description of these sections go to www.futuremedicine.com):

- Abstract
- Introduction
- Experimental
- Results and discussion
- Conclusions
- Future perspective
- References
- Reference annotations

Perspectives

Word limit: 3000

Perspectives should be speculative and very forward looking, even visionary. They offer the author the opportunity to present criticism or address controversy. Authors of perspectives are encouraged to be highly opinionated. The intention is very much that these articles should represent a personal perspective. Referees will be briefed to review these articles for quality and relevance of argument only. They will not necessarily be expected to agree with the authors' sentiments.

Special reports

Word limit: 3000

Special reports are short review-style articles that summarize a particular niche area, be it a specific technique or method.

Conference scenes

Word limit: 1500–3000

Conference scenes aim to summarize the most important research presented at a recent conference relevant to the journal's readership.

Editorials/opinions

Word limit: 1000–2000

Editorials are short articles on issues of topical importance. We encourage our editorial writers to express their opinions, giving the author the opportunity to present criticism or address controversy. The intention is very much that the article should offer a personal perspective on a topic

of recent interest. Longer **Commentaries** (word limit: 2000–5000) are also accepted.

Product reviews

Word limit: 1500–3000

Product reviews are short review-style articles that summarize a particular piece of equipment, software or book. They should include a description of the background, and the author's critical perspective on the product being reviewed.

Manuscript preparation

Spacing & headings: Please use double line spacing throughout the manuscript. No more than four levels of subheading should be used to divide the text and should be clearly designated.

Abbreviations: Abbreviations should be defined on their first appearance, and in any table and figure footnotes. It is helpful if a separate list is provided of any abbreviations. Spelling: US-preferred spelling will be used in the final publication.

Figures, tables & boxes: Future Medicine has a charge for the printing of color figures (i.e., each color page) in the print issue of the journal. We have no page charges and aim to keep our color charge to a minimum. The charge does not apply to the online version of articles, where all figures appear in color at no charge.

Copyright: If a figure, table or box has been published previously (even if you were the author), acknowledge the original source and submit written permission from the copyright holder to reproduce the material where necessary.

As the author of your manuscript, you are responsible for obtaining permissions to use material owned by others. Since the permission-seeking process can be remarkably time-consuming, it is wise to begin writing for permission as soon as possible.

Please send us photocopies of letters or forms granting you permission for the use of copyrighted material so that we can see that any special requirements with regard to wording and placement of credits are fulfilled. Keep the originals for your files. If payment is required for use of the figure, this should be covered by the author.

Key formatting points

Please ensure your paper concurs with the following article format:

Title: concise, not more than 120 characters.

Author(s) names & affiliations: including full name, address, phone & fax numbers and e-mail.

Practice points: a series of 6–10 bulleted points outlining methods of diagnosis and clinical treatment options.

Abstract: approximately 120 words and, for research papers, structured (Background, Results/Discussion/Methods, Conclusions). No references should be cited in the abstract.

Body of the article: article content under relevant headings and subheadings.

Conclusion: analysis of the data presented in the review.

Future perspective: a speculative viewpoint on how the field will evolve in 5–10 years time.

References:

- Should be numerically listed in the reference section in the order that they occur in the text.
- Should appear as a number i.e., [1,2] in the text.
- If websites or patents are included, please use a separate numbering system for them, i.e., start numbering website references at [101] and patents at [201] to allow the reader to distinguish between websites/patents and primary literature references both in the text and in the bibliography.
- Any references that are cited in figures/tables/boxes that do not appear in the text should be listed at the end of the reference list in the order they occur.
- Quote first six authors' names. If there are more than six, then quote first three *et al.*
- The Future Medicine Endnote style can be downloaded from our website at: www.futuremedicine.com/page/authors.jsp

Reference annotations: please highlight 6–8 references that are of particular significance to the subject and provide a brief (1–2 line) synopsis. Papers should be highlighted as either: * of interest or **of considerable interest.

Figures/Tables/Boxes: Summary figures/tables/boxes are very useful, and we encourage their use in reviews/perspectives/special reports. The author should include illustrations and tables to condense and illustrate the information they wish to convey. Commentary that augments an article and could be viewed as 'stand-alone' should be included in a separate box. An example would be a summary of a particular trial or trial series, a case study summary or a series of terms explained.

If any of the figures or tables used in the manuscript requires permission from the original publisher, it is the author's responsibility to obtain this. Figures must be in an editable format.