Reflections on screening blood donors for the presence of virus infections in England

Roger P Eglint^{†1} & John AJ Barbara[†]

[†]National Transfusion Microbiology Laboratories, NHS Blood & Transplant, Colindale Avenue, London, NW9 5BG, UK

[†]Author for correspondence: drreglin@btinternet.com

“These data beg the question – how safe does blood have to be?”

A recent publication by Stramer *et al.* has described the detection of US blood donors with virus infections by nucleic acid screening [1]. Traditionally, detection of HBV was by means of HBsAg and anti-HBc screening assays. Following the recent addition of nucleic acid testing (NAT), for the detection of infected donors, a higher rate of infection has been identified (1/600,000 donors). In these infected donors, the early predictions were for two to four seronegative HBV DNA positive donors, using testing of minipools of donation, plus one additional seronegative, NAT positive using single donation testing. The actual findings were nine seronegative and DNA positive donations, with a NAT-only infection undetected by minipool testing (a detection rate of 1 in 410,000 NAT-only positive donations), three conventional infections and five donors with low anti-HBs (3–36 international units [iu]/l) or anamnestic response. All these donors had been vaccinated 7–27 years previously. It should be remembered that, in the USA, universal HBV vaccination occurs, using virus strain A2. In the UK, HBV vaccination is routinely available only for specifically identified risk groups.

There was no evidence of disease associated with the viremias, which ranged from 11 to 86 iu/ml. The sub-genotype was determined to be A2, as expected – this is the most frequent genotype in the USA and is also the genotype in the vaccine. A total of six of the reported cases were HBV A2 strain and an additional five cases were non-A2 strains. There was an absence of infectivity unless the HBV DNA positive donor was anti-HBsAg positive with a titer less than 75 iu/ml when transmission occurred.

The frequencies of NAT window period donations were 1 per 1.85×10^6 for HIV and 1 per 1.85×10^6 for HCV. The incidence of detection of transfusion-transmitted infections during the 10 years of NAT testing was five HIV donations, which infected six of eight recipients; one HCV infection and no HBV infections by transfusion since 2006, when an ultrasensitive HBsAg assay was introduced. There has previously been transmission of HBV from the plasma but not from the associated red blood cell pack of a donor.

In the UK there is no universal vaccination policy for HBV. In the general population there are 600–800 new cases reported annually and an additional 100 cases in intravenous drug users. The residual risks of infection with HBV from donated blood with NAT screening in use are 3.2×10^6 in the USA; 0.67×10^6 in England; and 0.87×10^3 in South Africa. These data beg the question – how safe does blood have to be? The usual expectation of risk evaluation in all other areas of life is that a 1:1 million risk is virtually nil and the UK and USA are nearing (or are below) that risk level for several agents.

It can be considered that the risks associated with the blood donation process fall into two areas:

- There is a ‘clinical’ risk from blood when considered as a transplant, which is around 1 in 10^6 , which in reality is considered to be virtually nil. This is certainly the case when the other risks of associated healthcare are assessed. However, there is also a pharmaceutical aspect associated with EU product liability, which requires no risk to be attributable to the transfusion process. This has

Keywords

■ HBV ■ infections ■ NAT
■ nucleic acid testing ■ risk
■ testing

driven blood services into NAT with the associated escalating costs, albeit with a reducing residual risk.

- Given the current world financial situation, perhaps thought should be given to the concept of an acceptable risk and associated cost to the tax payer. It may be that the concept of zero risk at any cost is no longer sustainable and it should be amended to an agreed acceptable risk at an affordable/reasonable cost, but by whom? It is most uncertain that if in the event of the introduction of a pathogen inactivation (PI) process that is applicable to all blood products including red cells, the cost would be less than that incurred for virus marker screening. Alternatively, a mixed economy might be introduced, of virus screening for the red cells but PI for the remaining products.

How to address this situation

A revised approach to screening donors is indicated, but if NAT screening is stopped we know that some donors may become infected. What becomes of the adage "First do no harm"? What will be the accepted revised targets for the residual risks of infection? The number and therefore frequency of the NAT-only donations is very low (US data and costs: 24 NAT-only [1/153,952 donations at a cost of US\$1.69 $\times$ 10⁶]). In order that the cost of NAT per donation remains constant, the cost per minipool sample would have to be \$0.13 and the cost per individual NAT donation \$0.19.

Where is the balance of safety vs cost? For the USA the cost is \$40.6 $\times$ 10⁶ for a yield of 24 NAT 'extra' donations per year. It is salutary to reconsider a paper from Simmons *et al.* in May 2009 that first raised the question of the enormous cost of introducing NAT testing for blood services in the UK [2].

A further complication in the UK is the risk of vCJD transmission by transfusion. There are, as yet, no European Community (CE)-marked assays available for use on blood donations, although progress is being made on three possible assays for screening and confirmation. The validation process is in progress for these assays. If implemented this would be an additional burden on allocated funding from the Department of Health. Leucodepletion of components is under study to estimate the efficacy of prion removal by such methods. In addition, there will soon be a population of young adults eligible to

donate blood and who were born since the risk of infection from the food chain was removed in 1996. This opportunity will be grasped by the UK blood services.

As an additional safety measure, platelet units will be screened for evidence of bacterial contamination. The introduction of continuous culture during a product's life time will allow a shelf life of up to 7 days for platelet components. An alternative approach to a 'safe' product is the introduction of an effective PI system.

However this produces a 'catch 22' position where no one is willing to use PI until the system is perfected for all blood products. At present there is no PI system that can be applied effectively to the three major products: plasma, platelets and red cells. It is the latter component for which no truly effective PI system is yet available. Limited PI methods are available for blood products but not yet implemented by blood services, except for plasma products and only in a few laboratories for platelets. There remains the problem of producing a system that can be effectively applied to red cell components, and ideally to the whole blood unit so that inactivation is only applied once per donation.

What are the priorities for blood services? In this context, microbiologically safe products at the lowest cost possible. The potential new measures to increase the safety of products can not all be introduced without significant addition cost. Who will define the balance between ever-increasing safety (but at a low actual increment of safety yield) and escalating costs, out of all proportion to the rest of healthcare spending?

Financial & competing interests disclosure

The authors have no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.

No writing assistance was utilized in the production of this manuscript.

Bibliography

1. Stramer SL, Wend U, Candotti D *et al.*: Nucleic acid testing to detect HBV infection in blood donors. *N. Eng. J. Med.* 364, 236–247 (2011).
2. Simmons P, Kurtz J, Tedder RS: The UK blood transfusion service: over a (patent) barrel? *Lancet* 359, 1713–1714 (2009).