Research Highlights

Research highlights from the latest articles in
5-HTTLPR pharmacogenomics

The serotonin transporter gene-linked polymorphic region (5-HTTLPR), commonly reported with short and long alleles, in the promoter region of the gene, has received much attention. The short allele is associated with a lower gene-expression activity. This

month's Research Highlights aims to provide an insight into recent research regarding polymorphisms of the 5-HTTLPR variants, ranging from the pharmacogenomics (antidepressants, antipsychotics and opioids) to disease susceptibility (Type 2 diabetes).

5-HTTLPR and antidepressant efficacy in major depressive disorder

Evaluation of: Gressier F, Bouaziz E, Verstuyft C, Hardy P, Becquemont L, Corruble E: 5-HTTLPR modulates antidepressant efficacy in depressed women. *Psychiatr. Genet.* 19(4), 195–200 (2009).

More and more genetic variants associated with the antidepressant response in major depressive disorder (MDD) are being discovered [1]. The relationship between the serotonin transporter gene-linked polymorphic region (5-HTTLPR) variants of the solute carrier family 6 member 4 gene (*SLC6A4*; serotonin neurotransmitter transporter) and the drug efficacy of antidepressants in MDD has been investigated extensively. However, it remains an unresolved issue. In the Sequenced Treatment Alternatives for Depression (STAR*D) cohort of 1775 patients, it was observed that there was no association between treatment outcome and the 5-HTTLPR variant [2,3]. By contrast, a recent meta-analysis and a number of small studies have suggested that the short (S) and long (L) variants of *SLC6A4* 5-HTTLPR influence the antidepressant response [1].

In a study of Caucasian MDD in patients ($n = 103$; 74 women and 29 men), Gressier and colleagues tested the hypothesis that the 5-HTTLPR variants (S and L alleles) of the *SLC6A4* gene are associated with the treatment outcome with selective serotonin reuptake inhibitors (SSRI) and non-SSRI drugs after 4 weeks of treatment [4]. The SS genotype of the 5-HTTLPR variant was found to be associated with less

efficacy in depressed women ($p = 0.035$) [4]. Conversely, there was no trend of association found with the 5-HTTLPR variant in depressed men [4].

However, the study suffers from a low statistical power, with only 12 women having the SS genotype. When compared with 62 women carrying LL/LS genotypes, SS carriers improved less in the scores of Hamilton depression rating scale (HAMD; 45.7 vs 57.1%; $p = 0.08$) and were less likely to respond (as defined by $\geq 50\%$ HAMD reduction; 50 vs 79%; $p = 0.035$). Therefore, it is hard to conclude that this borderline significance is gender specific, as the title of the article suggests, since there are only seven men carrying the SS genotype. In addition, it is difficult to compare the report by Gressier and colleagues [4] with those of other related studies because of differences in study design or patient samples. The discrepancies among these studies may be explained by the small sample sizes (suggesting limited power), different outcome measures, treatment periods, ethnicities, analytical approaches and the use of different medications across studies [1].

Bibliography

- 1 Lin E, Chen PS: Pharmacogenomics with antidepressants in the STAR*D study. *Pharmacogenomics* 9(7), 935–946 (2008).
- 2 Hu XZ, Rush AJ, Charney D *et al.*: Association between a functional serotonin transporter promoter polymorphism and citalopram treatment in adult outpatients with major depression. *Arch. Gen. Psychiatry* 64(7), 783–792 (2007).

Hsien-Yuan Lane¹, Guochuan E Tsai²
& Eugene Lin^{3*}

*Author for correspondence:

¹Department of Psychiatry & Institute of Clinical Medical Science, China Medical University & Hospital, Taichung, Taiwan

²Los Angeles Biomedical Research Institute & Department of Psychiatry, Harbor-UCLA Medical Center, CA, USA

³Vita Genomics, Inc., 7 Fl., No. 6, Sec. 1, Jung-Shing Road, Wugu Shiang, Taipei, Taiwan

Tel.: + 886 289 769 123 Ext. 7751

Fax: + 886 289 769 523

eugene.lin@vitagenomics.com

Financial & competing interests disclosure

The authors' work is supported by National Science Council NSC-98-2627-B-039-001. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

- 3 Kraft JB, Peters EJ, Slager SL *et al.*: Analysis of association between the serotonin transporter and antidepressant response in a large clinical sample. *Biol. Psychiatry* 61(6), 734–742 (2007).
- 4 Gressier F, Bouaziz E, Verstuyft C, Hardy P, Becquemont L, Corruble E: 5-HTTLPR modulates antidepressant efficacy in depressed women. *Psychiatr. Genet.* 19(4), 195–200 (2009).

Serotonergic genes and antipsychotic-induced adverse reactions

Evaluation of: Al-Janabi I, Arranz MJ, Blakemore AI *et al.*: Association study of serotonergic gene variants with antipsychotic-induced adverse reactions. *Psychiatr. Genet.* 19(6), 305–311 (2009).

Evidence is accumulating to suggest that antipsychotic-induced adverse reactions result from a combination of a number of genetic variants [1–3]. In particular, several studies in different populations implicate the -759C/T (rs3813929) polymorphism of the 5-hydroxytryptamine receptor 2C (*HTR2C*) gene in antipsychotic-induced weight gain [1–3]. It has also been demonstrated that the TaqI A and -141C Ins/Del variants of the dopamine D2 receptor (*DRD2*) gene and the Ser9Gly polymorphism of the dopamine D3 receptor (*DRD3*) gene were strongly associated with drug-induced tardive dyskinesia [1–3].

In a study of Caucasian patients (n = 427) receiving antipsychotic treatment, Al-Janabi and colleagues determined whether serotonergic genes are associated with antipsychotic-induced

adverse reactions, such as extrapyramidal symptoms, tardive dyskinesia, obesity and diabetes [4]. There were nine variants, including 5-hydroxytryptamine receptor 2A (*HTR2A*; 102T/C [rs6313], His452Tyr [rs6314]), *HTR2C* (-995[G/A] [rs3813928], -759C/T [rs3813929], Cys23Ser [rs6318]), tryptophan hydroxylase 2 (*TPH2*; -8933A/G [rs17110451], -366C/T [rs4448731]) and *SLC6A4* (5-HTTLPR and -15370[A/G] [rs25531]) that were investigated in their study [4]. Al-Janabi *et al.* reported that the *HTR2A* 102T/C polymorphism was associated with BMI (p = 0.005), and they also found that the *TPH2* -366C/T polymorphism was significantly associated with the presence of diabetes (p = 0.01) [4]. However, they failed to detect other statistically significant associations between the genetic variants and antipsychotic-induced adverse reactions [4].

The study is limited by its cross-section design, incomplete coverage of the gene variants and sample size. Predicting the antipsychotic-induced metabolic syndrome, such as obesity, diabetes and hyperlipidemia, is of paramount importance for

clinical psychiatry [1–3]. Al-Janabi *et al.* suggest a moderate role of serotonergic polymorphisms for the development of these side effects. However, these findings are marginal and it is still premature to have genetic tests for the pretreatment prediction of antipsychotic response in patient care. It may become a reality after prospective large-series clinical trials to validate clinical factors and genetic markers [1–3].

Bibliography

- 1 Lane HY, Liu YC, Huang CL *et al.*: Risperidone-related weight gain: genetic and nongenetic predictors. *J. Clin. Psychopharmacol.* 26(2), 128–134 (2006).
- 2 Arranz MJ, de Leon J: Pharmacogenetics and pharmacogenomics of schizophrenia: a review of last decade of research. *Mol. Psychiatry* 12(8), 707–747 (2007).
- 3 Arranz MJ, Kapur S: Pharmacogenetics in psychiatry: are we ready for widespread clinical use? *Schizophr. Bull.* 34(6), 1130–1144 (2008).
- 4 Al-Janabi I, Arranz MJ, Blakemore AI *et al.*: Association study of serotonergic gene variants with antipsychotic-induced adverse reactions. *Psychiatr. Genet.* 19(6), 305–311 (2009).

Serotonin transporter gene and the analgesic response to opioids

Evaluation of: Kosek E, Jensen KB, Lonsdorf TB, Schalling M, Ingvar M: Genetic variation in the serotonin transporter gene (5-HTTLPR, rs25531) influences the analgesic response to the short acting opioid remifentanyl in humans. *Mol. Pain* 5, 37 (2009).

With the rapid advancement of pharmacogenomics, increasing literature demonstrating that an individual's polymorphic genotype may affect their therapeutic response to analgesics [1]. In particular, the cytochrome P450, family 2, subfamily D, polypeptide 6 (*CYP2D6*) gene has been reported to

modulate the pharmacokinetics of opioid medications in humans [1]. It has also been reported that the pharmacodynamics of analgesic drugs is affected by some genes, including catechol-O-methyltransferase (*COMT*), melanocortin-1 receptor (*MC1R*) and opioid receptor, $\mu 1$ (*OPRM1*) [1].

In a small study of Caucasian healthy participants ($n = 43$; 31 women and 12 men), Kosek and colleagues investigated the hypothesis that the 5-HTTLPR and rs25531 variants of the *SLC6A4* gene are associated with the analgesic response to remifentanyl, a rapid and potent opioid drug [2]. Both polymorphisms (5-HTTLPR and rs25531) are commonly referred as the triallelic 5-HTTLPR genotypes that differentiate individuals with a high (L_A/L_A), intermediate (L_A/L_G , L_A/S_A) or low (S_A/S_A , L_G/S_A) transcriptional efficacy. Kosek and colleagues found that remifentanyl had a significantly better analgesic effect ($p < 0.02$) in subjects with a genotype coding for low

expression (S_A/S_A and L_G/S_A) compared with subjects with high expression (L_A/L_A) after heat pain stimulation [2].

There were several limitations to the study by Kosek and colleagues, as follows. First, the sample size was very small, thereby preventing definite conclusions from being drawn [2]. Moreover, there was no clear dose–response curve, with the intermediate expression group experiencing a similar analgesic effect as the high expression group, which was likely a result of the lack of power. In the absence of a placebo control, it could not be determined whether the observed associations were directly attributable to opioid treatment.

Bibliography

- 1 Lacroix-Fralish ML, Mogil JS: Progress in genetic studies of pain and analgesia. *Annu. Rev. Pharmacol. Toxicol.* 49, 97–121 (2009).
- 2 Kosek E, Jensen KB, Lonsdorf TB, Schalling M, Ingvar M: Genetic variation in the serotonin transporter gene (5-HTTLPR, rs25531) influences the analgesic response to the short acting opioid remifentanyl in humans. *Mol. Pain* 5, 37 (2009).

5-HTTLPR and Type 2 diabetes

Evaluation of: Iordanidou M, Tavridou A, Petridis I *et al.*: The serotonin transporter promoter polymorphism (5-HTTLPR) is associated with Type 2 diabetes. *Clin. Chim. Acta* 411(3–4), 167–171 (2009).

Type 2 diabetes (T2D) is a chronic disorder characterized by high blood glucose in the context of insulin insensitivity and pancreas β -cell dysfunction [1–4]. The disease affects more than 150 million individuals worldwide [1–4]. Owing to the high prevalence of T2D, extensive research is still being carried out to identify disease-susceptibility loci using candidate gene approaches, genome-wide association studies, gene-expression profiling and family linkage studies [1–4]. Evidence is accumulating to suggest that T2D is genetically heterogeneous, as more and more genetic components associated with T2D are being discovered [1–4]. Common T2D genes include adiponectin, *CIQ* and collagen domain containing (*ADIPOQ*), ectonucleotide pyrophosphatase/phosphodiesterase 1 (*ENPP1*), growth hormone secretagogue receptor (*GHSR*),

peroxisome proliferator-activated receptor γ (*PPAR\gamma*) and transcription factor 7-like 2 (*TCF7L2*) [1–4].

In a study of Caucasian patients ($n = 463$; including 252 T2D patients and 211 nondiabetic controls), Iordanidou and colleagues examined whether the 5-HTTLPR variant (S and L alleles) of the *SLC6A4* gene is associated with T2D [5]. They found that the frequency of SS and SL genotypes of 5-HTTLPR was significantly higher ($p < 0.001$) in the T2D group (77.0%) than in the nondiabetic group (61.6%), and they further suggested that the S allele of 5-HTTLPR was strongly associated with the presence of T2D [5]. Moreover, the odds ratio analysis demonstrated that the genetic risk for subjects carrying at least one S allele was increased (odds ratio: 2.14; 95% CI: 1.41–3.25) compared with nondiabetic subjects after adjustment for age, sex and BMI [5].

One limitation of the study by Iordanidou and colleagues is the failure to investigate whether such an association is found in other populations [5]. In addition, the contributions of other markers in the serotonin neurotransmitter transporter gene could be further explored in future work. Finally, although this study

has the best power out of the four studies discussed, independent replications in large sample sizes are needed to confirm the role of the 5-HTTLPR polymorphism found in their study in T2D.

Bibliography

- 1 Owen KR, McCarthy MI: Genetics of Type 2 diabetes. *Curr. Opin. Genet. Dev.* 17(3), 239–244 (2007).
- 2 Doria A, Patti ME, Kahn CR: The emerging genetic architecture of Type 2 diabetes. *Cell Metab.* 8(3), 186–200 (2008).
- 3 Lin E, Pei D, Huang YJ, Hsieh CH, Wu LS: Gene–gene interactions among genetic variants from obesity candidate genes for nonobese and obese populations in Type 2 diabetes. *Genet. Test. Mol. Biomarkers* 13(4), 485–493 (2009).
- 4 Wu LS, Hsieh CH, Pei D, Hung YJ, Kuo SW, Lin E: Association and interaction analyses of genetic variants in *ADIPOQ*, *ENPP1*, *GHSR*, *PPAR\gamma* and *TCF7L2* genes for diabetic nephropathy in a Taiwanese population with Type 2 diabetes. *Nephrol. Dial. Transplant.* 24(11), 3360–3366 (2009).
- 5 Iordanidou M, Tavridou A, Petridis I *et al.*: The serotonin transporter promoter polymorphism (5-HTTLPR) is associated with Type 2 diabetes. *Clin. Chim. Acta* 411(3–4), 167–171 (2009).